


Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.





BULLETIN No. 209

Contribution from the Bureau of Plant Industry
WM. A. TAYLOR, Chief



Washington, D. C.

PROFESSIONAL PAPER

August 6, 1915.

**TESTING GRAPE VARIETIES IN THE VINIFERA
REGIONS OF THE UNITED STATES.**

By GEORGE C. HUSMANN,

Pomologist in Charge of Viticultural Investigations, Office of Horticultural and Pomological Investigations.

CONTENTS.

	Page.		Page.
Introduction.....	1	Adaptation to soil, climatic, and other con- ditions.....	12
Cooperative experiment vineyards and their nature.....	2	Hybrids.....	16
Acreage in the California experiment vine- yards.....	10	Growth ratings of resistant vines and direct producers.....	16
General plan of plantings in the experiment vineyards.....	10	Congeniality and adaptability of vines.....	26
Phenological records.....	10	Behavior of grapes grafted and on their own roots.....	27
Destruction of vineyards.....	11	Conclusions and suggestions.....	155
Factors in resistance.....	12		

INTRODUCTION.

A résumé of the viticultural investigations in the Vinifera regions of the United States up to 1910 was reported in Bulletin 172 of the Bureau of Plant Industry. The present publication supplements that bulletin and gives additional data on the investigations then under way, as well as reports upon researches started since the date of that publication. The fundamental problems of the Vinifera region, as determined by the early surveys, were found to require (1) a comprehensive test of the resistant varieties of vines to determine their adaptability to the different soils and climatic conditions; (2) a study of the congeniality of Vinifera varieties to the different resistant-stock varieties; (3) a study of the behavior of fruiting varieties to determine those best adapted to the different localities; and (4) a consideration of all classes of grapes with reference to their resistance to destructive insects and diseases and, if found necessary, the origination of an entirely new class of grapes better adapted to

Pacific coast conditions. These questions still remain the broad cardinal problems, but the facilities for their solution have been much enlarged and a number of subordinate problems that developed in the prosecution of the work have been taken up as far as the means and facilities of the Department of Agriculture permitted.

COOPERATIVE EXPERIMENT VINEYARDS AND THEIR NATURE.

To afford facilities for solving these problems, the Bureau of Plant Industry has established 12 experiment vineyards on the Pacific coast. One of these is at the Plant Introduction Field Station, Chico, Cal., and 11 are located in various other grape-growing centers in cooperation with growers.

A brief description of the purpose, location, soil, and climatic conditions at or near each of these vineyards follows. (Fig. 1.) Those desiring correlation and mechanical analyses of the soils and fuller climatic data are referred to Bulletin 172 of the Bureau of Plant Industry. The soil descriptions are from data furnished by the Bureau of Soils, while the weather data are taken from records furnished by the San Francisco office of the Weather Bureau, through Prof. McAdie, and from observations made in the experiment vineyards.

MAIN VINEYARDS.

Three primary vineyards of 20 acres each are located near Oakville, near Fresno, and at Guasti, Cal.

At the Oakville, Fresno, and Guasti experiment vineyards viticultural material introduced from foreign countries is tested. In these vineyards the adaptability to different localities and the value of grape varieties for different uses is determined; the relative resistance of grape varieties to destructive insects and diseases is inquired into; the congeniality of grape varieties to the different resistant sorts is determined; and grape varieties not now grown in the *Vinifera* regions of this country are tested, with a view to the possibility of their supplanting some of the varieties now grown. Experiments to determine how the various varieties should be propagated, grafted, pruned, trained, and otherwise cared for are under way. These vineyards offer some opportunity for the broad viticultural research and experimental work that is needed, and furnish practical object lessons in viticulture and facilities for solving some of the many commercial problems of the industry.

OAKVILLE EXPERIMENT VINEYARD.

The Oakville Experiment Vineyard (Pl. I, fig. 1) was established in the spring of 1903, and is located 1 mile west of Oakville, Napa Co., Cal., on the property of the To-Kalon Vineyard Company, at an elevation of 161 feet above sea level. The soil is a dark-brown or black

gravelly clay loam or heavy loam, containing a large quantity of organic matter formed in a swamp or lagoon extending in past geological ages up Napa Valley from San Pablo Bay, typical of the greater part of the soils in the valley floor. On weathering, the shales, sandstones, limestones, lime conglomerates, and large quantities of usually lenticular or angular gravel with little erosion of edges are washed down from the steep hills or mountains surrounding Napa Valley on




FIG. 1.—Map of California, showing (by large black dots) the location of the twelve experiment vineyards of the Bureau of Plant Industry.

all sides except the south, and tend to form a heavy or clayey soil with only small quantities of sand. No hardpan or alkali appears. The surface is undulating, affording a fairly rapid run-off of surplus rain-water, though in places the subsoil is quite wet during the spring months. No irrigation is necessary. The clay and silt in the subsoils greatly aid in retaining moisture in spite of the large quantity of

gravel (20 to 40 per cent) they contain. Cultivation reduces the surface to a good mulch. When grape culture in Napa Valley and the adjoining foothills became important, a reputation for the superior qualities of its dry wines was rapidly made, especially for the excellence of the white wines. This reputation has been sustained, and Napa County has remained one of the leading dry-wine sections of the State.

FRESNO EXPERIMENT VINEYARD.

The Fresno Experiment Vineyard (Pl. I, fig. 2) was established in the spring of 1903. It is located 3 miles east of Fresno on the property of the Fresno Vineyard Company, at an elevation of 290 feet above sea level. The soil is the San Joaquin sandy loam. The fact that it is an outlying isolated portion of soil of this character accounts for the increased depths to hardpan and the sandier subsoil immediately above. The San Joaquin sandy loams are confined to lands adjacent to the lower foothills on the eastern side of the San Joaquin and Sacramento Valleys, where 75,000 acres near Fresno, 6,000 acres near Stockton, and 265,000 acres about Sacramento have already been mapped. The soil is light red in color, granitic in origin, and composed largely of sharp, angular particles. The surface is rolling and generally covered with hog wallows and small mounds.

In the experiment-vineyard plat two varieties of soil were recognized, namely, adhesive sandy loam, closely approaching a true loam, and a friable sandy loam. The former retains moisture longer than the latter, which is a deeper soil of lighter texture. Leveling the plat disturbed the natural soil conditions, decreasing the depth of the sticky, adhesive sandy loam in spots and exposing free sandy loam in others, causing the hardpan underlying the plat to occur at depths varying from scarcely 20 inches to over 6 feet, whereas the average depth at which it occurs is $3\frac{1}{2}$ to 4 feet below the surface. This hardpan, which always accompanies San Joaquin sandy-loam soil, is a red iron-sandstone substance cemented by hydrates of iron and alumina combined with clay. When this occurs at 2 feet or less below the surface, blasting is necessary. Trees and vines thrive when the hardpan is broken or where it lies at a sufficient depth below the surface.

The soils of the plat above the hardpan contain alkali varying from less than 0.05 to more than 20 per cent; but in the lowest grade soil no alkali is visible. Of the salts, over 90 per cent are chlorids, as follows: Calcium, 50; magnesium, 25; sodium, 15; and potassium, about 2 per cent. The remainder consists of calcium sulphate and bicarbonate of soda. The depth of the water table on the tract averages 3 feet.

Fresno is in the center of the raisin industry of the country, and is also one of the most important wine and brandy producing districts of California.

At Oakville there are being assembled and tested on resistant stocks all the grape varieties of the world thought to be of value to the Pacific slope, while at Fresno similar tests are made of raisin, currant, and fleshy varieties.

GUASTI EXPERIMENT VINEYARD.

In view of the entirely different conditions prevailing south of the Tehachapi Pass, especially in the desert region, the Department of Agriculture in the spring of 1904 established another experiment vineyard of like importance and acreage in the San Bernardino desert, at Guasti, Cal., 950 feet above sea level, on the property of the Italian Vineyard Company. (See Pl. I, fig. 3.) The soil mapped as Maricopa gravelly sand is a gray-brown gravelly sand of very uniform texture to an unknown depth. The surface is compact when untilled, because the sharp, angular sand composing the soil becomes somewhat cemented by the organic matter occurring in the topsoil. At a depth of 3 feet the soil is more concentrated and often yellowish from the oxidation of iron in the soil. The soil is almost entirely granitic, and is washed from the Sierra Madres. It contains quantities of undecomposed potash, feldspar particles, which should insure abundant potash for the maturing of grapes. The soil covers most of the San Bernardino Valley floor, and when thoroughly cultivated holds moisture well, the fine sand and silt giving the soil capillary power to bring water up from below. Two of the largest vineyards of the world are in this valley, on similar soil. As the phylloxera is not known to exist here, the plantings in the experiment vineyard are principally Vinifera varieties.

CHICO VARIETAL VINEYARD.

The Chico Varietal Vineyard is located at the plant-introduction field station, 3 miles east of Chico, Butte Co., Cal., and is about 196 feet above sea level (Pl. II, fig. 1). The soil, an alluvium, is composed of material brought down from the mountains and hills on the east and is from 8 to 12 feet deep. It is underlain by a body of sandy water-worn gravel and boulders, which always carry water. The soil is of light texture, varying from light loam to heavy, fine sandy loam, the heaviest being loam. It is well drained and easily cultivated. The heavy, fine sandy loam consists of from 30 to 36 inches of fine sandy loam, underlain by very fine sandy loam, usually containing some gravel. The light loam has from 10 to 15 inches of fine sandy loam or sandy loam underlain by a heavier structure closely approaching loam. The largest area of this soil is found about Chico, but similar soil occurs in the Feather and the Bear River Valleys.

At the Chico vineyard are being assembled and maintained two plants each of grape varieties that prove of special value for specific

purposes, together with grape immigrants from all parts of the world introduced by the Office of Foreign Seed and Plant Introduction. The first plantings were made in the spring of 1906, and the collection here already comprises 308 resistant and direct-producing sorts and 141 *Vinifera* varieties grafted on resistant stocks.

SMALLER VINEYARDS.

In addition to the main plantations, outlying vineyards of 10 acres each have been established to test varieties at different altitudes, at varying distances from the ocean, bays, and other bodies of water, under different climatic and other conditions, and on the leading types of vineyard soils not found at the Oakville, Fresno, and Guasti vineyards.

BRAWLEY EXPERIMENT VINEYARD.

The Brawley Experiment Vineyard was established in the spring of 1911 on the New River, 1 mile west of Brawley, Imperial Co., Cal., on the property of Mrs. Mabel Oakley, about 110 feet below sea level. (See Pl. II, fig. 2.) The soil, Imperial loam, is sediment brought down by the Colorado River and deposited in strata either while the area was still submerged or from the overflow waters as they spread over the plain. Before irrigation these strata, varying from 1.1 to 2 or 3 inches in thickness, are quite hard and look like shale, but water softens them readily. This soil after irrigation is a sticky reddish loam, a little heavier than a silt loam, having a depth of 4 to 6 feet and resting on a clay or clay-loam subsoil, which in turn is underlain by alternate strata of lighter and heavier material to an indeterminate depth. The surface is usually smooth and level and almost devoid of vegetation. The soil often contains considerable organic matter and when irrigated is productive. Alkali is found in all of it and often greatly in excess of what even the most resistant plants can stand, thus making much reclamation work necessary. This soil is by far the most extensive type in the Imperial Valley. It extends from about the middle of the eastern boundary across in a northwesterly direction into the Salton Sink. A large part of the area west and southwest of Imperial and large tracts between Mesquite Lake and the Mexican line are of this type of soil.

The Imperial Valley is said to be the earliest fruit-ripening district in the United States, and as no phylloxera have been located in it the plantings of 280 grape varieties in this vineyard are all *Vinifera*.

COLFAX EXPERIMENT VINEYARD.

The Colfax Experiment Vineyard was established in the spring of 1906, on the property of Mr. Louis Cortopassi, in the Sierra Nevada, 2,412 feet above sea level, 1½ miles southwest of Colfax, Placer Co., Cal. (See Pl. II, fig. 3.)

The soil, hilly, usually fairly deep, and well drained, originated in the decomposition of the Mariposa formation, consisting of dark shales or slates, sandstones or quartzite sandstones, and conglomerates. The large amount of iron present from decomposing volcanic-rock material, where exposed to perfect weathering, gives the soil a deep red color. Dark, shallow, conglomerate rocks sometimes outcrop in spots, and rock fragments occur. The first few inches are often dark red from the accumulation of organic matter. The first 8 to 18 inches are usually brownish red clay or clay loam, underlain by 3 to 6 feet of red clay or clay loam, with partially decomposed and weathered rock formations, giving the subsoil a yellow appearance.

Rock outcrops of conglomerates, chert, and slate occur in the higher portions. The native vegetation is manzanita, chaparral, live oak and yellow pine.

The Colfax district is unique in the diversity of the fruit grown on sidehill locations.

GEYSERVILLE EXPERIMENT VINEYARD.

The Geyserville Experiment Vineyard was established in the spring of 1904, on the property of John D. Bosch, just east of Geyserville, Sonoma Co., Cal., against a range of high hills (Pl. III, fig. 1), 236 feet above sea level. To a depth of $2\frac{1}{2}$ to 3 feet the soil consists of a uniform dark gravelly loam with a subsoil of light or yellowish brown color, similar in texture to the topsoil. The soil is very mellow and carries considerable humus, which enables it to retain moisture well. This type of soil extends over considerable areas along the streams and the floor of the Sonoma Valley, having been washed down from the shale, schist, and conglomerate hills.

Soils of this type produce some of the choicest dry wines, both red and white, of the State.

LIVERMORE EXPERIMENT VINEYARD.

The Livermore Experiment Vineyard was established in the spring of 1904, on the property of Mr. C. H. Wentz, 3 miles south of Livermore, Alameda Co., Cal., at an elevation of about 450 feet above sea level. (See Pl. III, fig. 2.) The vineyard has a very uniform, level, alluvial soil, derived from decomposed shales and schists, and is full of rounded gravel washed down from the surrounding mountains.

The surface soil is a dark-brown gravelly loam; the second, third, and fourth feet gravelly sandy loam, replaced by gravelly sand in the fifth foot. The humus decreases with the depth, while the gravel increases, varying from 30 to 59 per cent. The proportion of clay is greater than that of silt. This gives the soil a very heavy appearance, the gravel sticking together very tightly when dry or packed. No alkali exists, but ground water is encountered at 5 or 6 feet in some

places. These soils are common over the Livermore Valley and produce a superior white wine of the sauterne type. This experiment vineyard was discontinued on July 1, 1914.

LODI EXPERIMENT VINEYARD.

The Lodi Experiment Vineyard was established in the spring of 1904, on the Lawrence & Murray property, about one-fourth of a mile northeast of Lodi, San Joaquin Co., Cal., about 55 feet above sea level. (See Pl. III, fig. 3.) A large body of this soil exists between Lodi and Acampo.

There are two variations on the plat. Phase No. 1 is a brown, free, sandy loam, underlain below 4½ feet by a more adhesive light-brown or yellowish sandy loam. Occasional iron concretions give the subsoil a mottled color. The soil has good capillarity and the water table occurs at 5 to 6 feet. Phase No. 2, an adhesive sand, was formed by an old stream channel. This is light-brown sand to a depth of 3 feet, the subsoil water-washed sand, much looser in texture and lighter in color, and dry to a depth of more than 6 feet, as the soil texture is too loose to exert much capillary force. There is no hardpan or alkali. The soils are, however, deficient in lime; otherwise, they are very productive, comparatively level, unirrigated, and easily tilled. This locality is well known for its table grapes and as a table-grape shipping point.

MOUNTAIN VIEW EXPERIMENT VINEYARD.

The Mountain View Experiment Vineyard was established in the spring of 1904, on the property of Mrs. Caroline Distel, 2 miles west of Mountain View, on the west side of the Santa Clara Valley, 76 feet above sea level. (See Pl. IV, fig. 1.)

The soil is a gravelly Placentia sandy loam. The first 12 inches is a dark-brown, gravelly sandy loam, dark from humus; below this, to 4 feet, the subsoil becomes redder, sandier, and more gravelly until sand is encountered. It is well drained, but inclined to become too dry in summer and fall. The surface soil at times becomes quite compact, and when plowed breaks up into hard clods. When tilled at the right time it works into a very mellow condition. These soils are from washings of granitic sandy shales and schist rocks. Before the destruction of vineyards by phylloxera and other agencies the Santa Clara Valley was the banner dry-wine producing section of California. The following areas of Placentia sandy loam have been surveyed in California: San Jose, 61,500; lower Salinas, 74,000; Los Angeles, 66,000; San Bernardino, 87,000; San Gabriel, 48,800; and Santa Ana, 16,800 acres. Soils of this series occur through the coast range of mountains from San Francisco to the Mexican line, occupying undulating portions of valleys close to the hills. The Mountain View Experiment Vineyard was discontinued on July 1, 1912.



FIG. 1.—OAKVILLE EXPERIMENT VINEYARD.



FIG. 2.—FRESNO EXPERIMENT VINEYARD.



FIG. 3.—GUASTI EXPERIMENT VINEYARD.



FIG. 1.—CHICO VARIETAL VINEYARD.



FIG. 2.—BRAWLEY EXPERIMENT VINEYARD.



FIG. 3.—COLFAX EXPERIMENT VINEYARD.



FIG. 1.—GEYSERVILLE EXPERIMENT VINEYARD.



FIG. 2.—LIVERMORE EXPERIMENT VINEYARD.



FIG. 3.—LODI EXPERIMENT VINEYARD.



FIG. 1.—MOUNTAIN VIEW EXPERIMENT VINEYARD.



FIG. 2.—SONOMA EXPERIMENT VINEYARD.



FIG. 3.—STOCKTON EXPERIMENT VINEYARD.

SONOMA EXPERIMENT VINEYARD.

The Sonoma Experiment Vineyard was established in the spring of 1904, on the property of the Gundlach-Bundschu Wine Company, about 2 miles south of Sonoma, Sonoma Co., Cal., about 110 feet above sea level. (See Pl. IV, fig. 2.) The soil is of rather poor quality, and to a depth of 8 or 10 inches is a gray loam more easily tilled than its texture indicates. The subsoil to 6 feet or more in depth is clay, changing at 4 feet, with an increase of sand, from a light-brown to a yellowish brown color. The soil is found near where weathered shales from the surrounding hills have been partially broken down and transported into the valleys, where they decompose. The soil usually occupies small, undulating ridges, or elevations, and is surrounded by the dark-brown, alluvial clay loam of the valley floor. The surface drainage is good, and no injurious quantity of alkali exists. This soil occupies extensive areas in the Sonoma Valley and in the adjacent bay regions and produces superior white wines of the Riesling, Chasselas, and Traminer types.

STOCKTON EXPERIMENT VINEYARD.

The Stockton Experiment Vineyard was established in the spring of 1907, on the property of the San Joaquin Valley Realty Company, a little over a mile southeast of Stockton, about 15 feet above sea level, on Stockton clay-loam adobe. (See Pl. IV, fig. 3.) This type, locally known as black adobe, was laid down in a swamp or tidal marsh in quiet water, the decomposing vegetation giving it a black color. It is a clay loam in texture, adhesive and sticky when wet and very hard when dry, cracking into large, cubical blocks full of small, cubical fractures. Sufficient rain slacks the clods readily. Cultivated when neither too wet nor too dry, the soil is friable and pulverizes well. The subsoil is a light-yellow silt loam, usually separated from the surface soil at a depth of $2\frac{1}{2}$ feet by a thin stratum, about one-half inch thick, of rather soft marly or calcareous hardpan, which is not always continuous and is often broken or disintegrated. Roots and water readily penetrate the subsoil, often passing through the hardpan. The depth to the water table varies from $3\frac{1}{2}$ to 6 feet for wet seasons and from 6 to 10 feet for dry seasons. This variation is influenced by a thin, marly hardpan which appears to hold the water down under pressure. It is somewhat difficult to establish vineyards on these soils, but when established they are very productive and lasting. Grapes for diverse purposes are grown on them. One of the largest sweet-wine establishments in the world is located near Stockton, and heavy shipments of table grapes grown on these soils are made. Soils of this type have been mapped in California as follows: Stockton, 53,312; Hanford, 5,470; and Fresno, 5,664 acres. It covers many thousand acres between the Marysville Buttes and about North Durham in the Sacramento Valley.

ACREAGE IN THE EXPERIMENT VINEYARDS IN CALIFORNIA.

The plantings and graftings in the California experiment vineyards now comprise the areas shown in Table I.

TABLE I.—Size of the twelve experiment vineyards in California of the Bureau of Plant Industry and number of varieties planted in each.

Vineyard.	Area.	Number of varieties.		Vineyard.	Area.	Number of varieties.			
		Resistants and direct producers.	Vinifera.			Resistants and direct producers.	Vinifera.		
			On their own roots.				On resistant stocks.	On their own roots.	On resistant stocks.
Brawley.....	<i>Acres.</i> 2.1		209	Livermore.....	<i>Acres.</i> 1.4	109			
Chico.....	1.8	308	141	Lodi.....	3.3	112	80		
Colfax.....	3.5	122	74	Mountain View.....	2	124			
Fresno.....	9	187	137	Oakville.....	16.6	306	262		
Geyserville.....	1.2	94	1	Sonoma.....	2	117			
Guasti.....	7	83	317	Stockton.....	1.4	91			

GENERAL PLAN OF PLANTINGS IN THE EXPERIMENT VINEYARDS.

All the plantings for comparative tests and study are made in regular checks of 10 vines of each variety, and all the larger plantings of resistant varieties are grafted in regular checks, usually of 10 vines each, and, where only preliminary readings are desired, the number of grafts of each variety put in have been a divisor of 10 in a check of 10 vines.

PHENOLOGICAL RECORDS.

Each vine or graft receives its block, row, and vine number. A complete history and accurate records are kept of all varieties from the time they are planted or grafted. Their behavior is closely noted, detailed descriptions are made of the vines and their respective fruits, and their value for specific uses and adaptability to different conditions are recorded. Table II summarizes the climatic observations made at the different vineyards.

TABLE II.—Temperature and rainfall at the twelve experiment vineyards of the Bureau of Plant Industry in California.

MAXIMUM TEMPERATURES (° F.).

Vineyard.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Annual.
Brawley ¹	80	85	95	106	118	115	115	112	111	105	93	81	118
Chico.....	74	78	83	94	104	111	114	116	106	96	82	74	116
Colfax.....	73	88	89	86	98	110	107	108	99	91	82	78	110
Fresno.....	73	77	84	101	110	107	115	113	108	98	84	74	115
Geyserville.....	79	83	91	102	108	116	116	110	113	103	98	79	116
Guasti.....	83	87	88	98	101	105	108	106	105	98	89	80	108
Livermore.....	77	79	88	95	108	108	113	107	108	99	87	75	108
Lodi.....	70	72	80	91	104	104	110	104	105	91	78	67	110
Mountain View.....	76	76	85	94	104	106	111	99	109	95	84	75	111
Oakville.....	77	75	86	95	106	109	110	105	110	98	84	74	110
Sonoma.....	77	76	82	91	104	109	106	101	111	97	82	72	111
Stockton.....	67	70	80	89	102	105	110	103	104	90	84	66	110

¹ Data for four years.

TABLE II.--Temperature and rainfall at the twelve experiment vineyards of the Bureau of Plant Industry in California--Continued.

MINIMUM TEMPERATURES (° F.).

Vineyard.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Annual.
Brawley ¹	24	29	35	38	50	52	57	60	55	40	26	20	20
Chico.....	18	20	25	30	33	40	46	48	40	34	21	22	18
Colfax.....	14	19	24	25	30	34	44	38	34	24	18	16	14
Fresno.....	24	25	30	34	40	42	51	52	42	36	31	24	24
Geyserville.....	21	21	29	30	33	37	40	39	35	30	26	23	20
Guasti.....	26	26	30	32	36	39	42	42	44	39	30	26	26
Livermore.....	23	24	30	30	34	39	41	41	40	34	25	23	23
Lodi.....	22	24	30	33	38	43	45	44	40	31	25	21	21
Mountain View.....	25	23	29	29	34	36	40	39	37	30	27	24	23
Oakville.....	20	24	25	26	30	34	32	34	35	27	25	17	17
Sonoma.....	23	27	29	27	32	34	41	34	35	32	26	23	23
Stockton.....	24	24	32	36	40	40	48	42	42	36	25	20	20

MEAN TEMPERATURES (° F.).

Brawley ¹	54.2	54.3	62.2	69.2	76.3	84.2	89.5	89.3	83.4	71.3	60.9	51.1	70.4
Chico.....	46.3	49.6	53.0	58.9	65.3	73.0	79.4	77.5	71.4	63.9	52.8	45.8	61.4
Colfax.....	42.6	44.1	46.3	52.9	59.4	68.6	75.1	73.6	65.7	59.0	51.6	43.5	56.9
Fresno.....	47.6	50.9	55.4	60.8	66.4	74.5	81.8	80.2	73.1	64.4	54.3	45.7	62.9
Geyserville.....	47.4	50.3	52.8	57.8	62.4	67.3	69.8	68.2	67.0	62.2	53.9	47.7	58.9
Guasti.....	49.9	51.7	54.7	58.3	60.7	67.2	75.2	73.2	69.9	64.4	56.5	51.3	60.9
Livermore.....	48.0	51.0	53.5	57.2	60.5	66.0	70.9	69.8	68.3	62.8	54.5	48.8	59.3
Lodi.....	46.5	49.4	53.1	57.6	63.0	68.8	72.9	70.6	67.0	60.0	51.5	45.0	58.8
Mountain View.....	49.1	51.2	53.3	56.1	59.0	62.7	66.0	65.1	64.3	60.2	54.0	48.1	57.4
Oakville.....	45.4	48.3	50.4	55.1	59.7	64.1	66.8	65.8	65.1	60.8	52.3	45.5	56.6
Sonoma.....	47.3	50.1	51.7	55.5	59.4	63.8	66.2	64.5	64.2	60.6	52.9	46.3	56.8
Stockton.....	45.7	48.8	52.5	56.9	62.1	68.3	73.2	71.1	67.8	61.1	51.6	44.2	58.6

PRECIPITATION (INCHES²).

Brawley ³													
Chico.....	6.41	3.77	5.46	1.14	0.96	0.29	T	0.01	1.03	0.79	2.03	3.50	25.39
Colfax.....	13.12	7.78	11.02	3.00	2.59	.95	T	.002	1.14	2.39	4.98	6.39	53.36
Fresno.....	2.11	1.30	2.24	.77	.58	.03	.001	T	.30	.56	.72	1.26	9.87
Geyserville.....	11.08	6.85	9.49	1.40	1.30	.45	T	T	.96	1.59	3.88	5.24	42.25
Guasti.....	6.05	4.30	6.40	1.29	.93	.12	.02	.01	.44	.56	.87	3.14	24.13
Livermore.....	4.55	2.00	3.75	.65	.57	.19	T	.03	.31	.43	.90	2.47	15.86
Lodi.....	5.02	2.68	4.71	.80	1.56	.26	T	.003	.40	.55	1.14	2.94	20.06
Mountain View.....	4.47	2.46	4.45	.80	.54	.18	T	.04	.52	.51	.96	2.49	17.42
Oakville.....	8.07	4.42	6.57	1.10	.78	.26	T	.01	.78	.77	2.10	3.95	28.81
Sonoma.....	6.80	4.91	6.02	.74	.87	.25	.34	T	.82	.83	2.75	4.00	28.33
Stockton.....	4.24	2.12	3.62	.72	.68	.14	T	.01	.48	.42	.99	2.31	15.73

¹ Data for four years.

² T=Trace.

³ No data available.

In the care and maintenance of the California experiment vineyards and in prosecuting researches in them, Mr. George C. Husmann, Pomologist in Charge of Viticultural Investigations, is assisted by Mr. Fred L. Husmann, Viticultural Superintendent, and Mr. Richard Schmidt, Assistant in Viticulture.

DESTRUCTION OF VINEYARDS.

In Bulletin 172 of the Bureau of Plant Industry the writer gives an account of the havoc wrought by the phylloxera insect to the vineyards of the Old World as well as to those of the Vinifera regions of this country. Since that bulletin was issued additional heavy losses have occurred.

The destruction in California of once flourishing vineyards covering more than 200,000 acres, through the so-called California vine disease and other agencies, but most largely through phylloxera, has already occurred. All that has been suggested to combat and eradi-

cate the phylloxera has been tried in this and other countries, but it is conceded that, with the exception of vineyards which can be flooded cheaply and sufficiently to kill the insect, the only way successfully to reestablish Vinifera vineyards is by growing the vines on phylloxera-resistant stocks.

FACTORS IN RESISTANCE.

The resistance of vines to phylloxera depends upon two factors: (1) The inherent resistant character of the vine and (2) its adaptation to soil, climatic, and other conditions.

The inherent or natural characteristics of the plant repel or invite the attacks of the phylloxera. The number of swellings, nodosities, and tuberosities from insect punctures and the rotting of the root occasioned by them progress more or less rapidly and deeply in accordance with the texture and character of the root attacked. The weakening and ultimate death of the vine are determined by the extent of the punctures and the progress of the rot upon the roots.

The nodosities rot more or less rapidly in the different grape species. In the Vinifera they are larger and usually rot in a very short time. In the American species the nodosities are smaller and do not rot so quickly, various species differing in this respect. The number of nodosities varies greatly on the different vine varieties, and when the insects have multiplied sufficiently to cover the smaller rootlets with them, eventually the larger roots are attacked, and if cancerous patches of decomposition are found on the more developed roots tuberosities occur.

Some varieties have nodosities on practically all rootlets, while some of the American species are not injured by the phylloxera except that a few nodosities form on their roots. In fact, the resistant ratings of different species are based on a determination of the relative number and size of the nodosities found on their roots.

In order to indicate with some degree of definiteness the resistance to phylloxera (not the value of the stock), scientists have provisionally adopted an arbitrary scale of ratings. In this scale the maximum of resistance or immunity is taken as 20, while absence of resistance (or no resistance) is reckoned as zero.

The necessary degree of resistance for the production of good crops varies with the character of the soil, stocks rating above 16 being considered sufficient for all soils, 14 to 16 for fairly good soils, and 10 to 14 for rich, moist, sandy soils.

ADAPTATION TO SOIL, CLIMATIC, AND OTHER CONDITIONS.

The ability of a vine to withstand or resist the attacks of the phylloxera without serious injury is greatly influenced by the adaptability of the vine to the climatic and soil conditions under which it is grown. These may increase or diminish the vigor of the plant, retard or favor the reparation of the insect injuries, and affect

the resistance of the plant by favoring or hindering the dissemination or activity of the insect. For instance, climatic conditions affect the multiplication of the insect and it can travel but little in sandy soils of a certain fineness. Then again, a variety which in one locality has splendid resistant qualities perishes in another locality having a similar soil but a different climate or in another locality having a similar climate but a different soil.

This is due to the fact that some species are adapted to moist soils, while others are variously adapted to dry soils, deep soils, or shallow soils, and also to the fact that the root systems of different species vary in habit of growth, some assuming a horizontal and others a vertical position; some roots are thick and fleshy and others small and wiry; some are soft, while others are firm. (See Pl. V.) Thus, a moisture-loving variety or one having a horizontal root system would not thrive in a dry, hot climate; neither would a variety with a deep root system thrive in a shallow, hard soil, or one adapted to a dry location thrive in a wet soil. For such reasons, a variety grown under congenial soil and climatic conditions will often prove more resistant than one of greater natural resistance grown under adverse soil and climatic conditions.

Resistance to phylloxera is also influenced by the congeniality existing between vine varieties when grafted on other vine varieties. Causes like these, and there are many others, affect the resistant qualities of vines, and it is with the study of the adaptation of varieties to varying conditions and the congeniality existing between vine varieties that the researches reported in this bulletin are particularly concerned.

The European work in the selection and breeding of resistant stocks has been determined largely by the necessity that such stocks be adapted to calcareous-soil conditions rarely encountered in the present Vinifera regions of this country. This renders it necessary for us to undertake our own researches and determinations. The varying soil, climatic, and other conditions in California make the selecting of the right grape species in the reestablishing of vineyards on resistant stocks rather a complex matter.

Of the 23 species of grapes native to North America, the 14 shown in Table III have been found sufficiently resistant to merit the attention of the viticulturist, and they are under test in the experiment vineyards of the United States Department of Agriculture.

Table III shows their natural habitat, the locations, sites, and the character of soil they prefer, the habits of the vines, their relative season of leafing (Pls. VI and VII), their root systems (Pl. V), the flowering and ripening of their fruit, the ease or difficulty of propagating them, their suitability for either bench or field grafting, and their relative resistance to phylloxera, cold, dampness, heat, and drought.

TABLE III.—Cultural data of fourteen American species of grapes whose varieties or hybrids are under test as resistant stocks in twelve experiment vineyards in California.¹

Name and region of nativity.	Preferred location.	Vine.	Roots.	Season of leafing, flowering, and ripening.	Percentage of cuttings taking root.	Grafting adaptation.	Resistance to—				
							Phylloxera (out of a possible 20).	Cold.	Dampness.	Heat.	Drought.
<i>Vitis labrusca</i> (northern fox grape): Allegheny Mountains, from New England to South Carolina. <i>V. candeliana</i> (mustang grape): Oklahoma, Texas, and New Mexico.	Wet thickets; granitic soils Black waxy lands or adobe	Vigorous, medium-sized climber. Moderately vigorous, medium climber.	Large, fleshy. Vigorous, tender.	Very late. Medium early.	85 30	BF ¹ F...	5 15	VG..... F... F..	F.. G..	F.. G..	F.. G..
<i>V. aestivalis</i> (summer grape): Southern New York to Florida; westward to the Mississippi and Missouri Rivers. <i>V. incocornii</i> (post-oak or turkey grape): Texas	High, warm, gravelly, moist soils. High, well drained timber lands, granitic gravelly clay, compact deep, rich river-bank soils.	Vigorous, medium-sized climber. Vigorous, good-sized climber.	Rather large, hard, plunging. Medium size, hard, and long.	Medium late. do.	50 40	F... F...	14 14	VG..... F.....	G.. G..	G.. G..	G.. G..
<i>V. monticola</i> (sweet mountain grape): Texas	Low limestone hills; does moderately well in sandy soils.	Rather small; good grower.	Bushy, plunging.	do.	65	FB	18	F.....	G..	G..	G..
<i>V. berlandieri</i> (little mountain grape): Texas and Mexico	Tops, sides, and bottoms of limestone hills.	Slender; medium grower.	Strong and plunging.	Late.	40	F...	19	F.....	G..	G..	G..
<i>V. cordifolia</i> (frost or sour winter grape): Great Lakes to Florida, abundant in Illinois, Tennessee, Missouri, and Arkansas. <i>V. cinerea</i> (sweet winter or ashly grape): Illinois to Texas.	Deep, rich, loose soils on river banks. do.	Vigorous, strong climber. Vigorous, strong grower.	Strong, hard, carnosus. Large, fleshy, plunging	do. Very late.	25 25	F... F...	18 15	VG. F.. F... G..	G.. G..	G.. G..	G.. G..
<i>V. champini</i> (adobe-land grape): Texas	Limestone hills; adapts itself to a variety of soils.	Vigorous, spreading grower.	Large, ramified, plunging.	Early to medium.	80	FB	12	F... G..	G..	G..	G..

<i>V. doaniana</i> (Texas Panhandle large grape): Texas.....	Sandy, limy soils.....	Slender, fair grower...	Numerous, thick, deeply penetrating.	Early.....	60	F...	12	F...	G...	G.....
<i>V. longii</i> (Solonis, bush or gulch grape): Texas Panhandle, New Mexico, Kansas, and Colorado.	Ravines along streams. Creaceous, generally rich, sandy, moist, always fresh soils.	Bushy upright, vigorous grower.	Large, ramified, horizontal, hard.	do.....	60	F...	14	F...	G...	G... G.
<i>V. rupestris</i> (sand, sugar, or rock grape): From the Rio Grande in Texas northwesterly into Oklahoma, Missouri, Kentucky, and Tennessee; Cumberland Mountain in north to Pennsylvania.	Open places in poor soils and along gravelly banks and ravines.	Vigorous, short, bushy grower.	Long, slender, or strong, plunging.	Very early.....	80	B...	19	G.....	G...	G... G.
<i>V. vulpina</i> (riparia or riverside grape): From Salt Lake east and from Texas north in all the States as far as 90 miles north of Quebec.	Moist, loose, sandy soils along creeks and river bottoms.	Vigorous; medium size.	Long, thin, slender, hard, wiry, ramified.	do.....	85	FB.	19	YG.	G...	G.....
<i>V. bicolor</i> (blue grape): Northern Missouri, Illinois, Wisconsin, Indiana, Michigan, Kentucky, Pennsylvania, New York, New Jersey, Maryland, and Ontario.	Black sandy and red siliceous soils.	Fair grower.....	Rather hard, large, plunging.	Late.....	40	F...	16	YG.	G...	G... G.

1 Abbreviations used in table: Under grafting, B for bench, F for field; under resistance, F for fair, G for good, and V for very.

HYBRIDS.

In the attempts to secure resistant stocks suited to soil, climatic, and other conditions which at the same time would prove congenial, lasting, and productive stocks on which to graft the *Vinifera* varieties, many difficulties were encountered. For instance, the stock may be adapted to the soil, but it may be so hard to root as to make its commercial use impracticable. Again, the stock may be suited to the soil and it may root easily and be resistant, but not congenial to or make a lasting junction with *Vinifera* varieties; or the congeniality of the variety may be good, but the fruitfulness of the graft may be impaired.


In many cases also no resistant species are exactly suited to the soil and climatic conditions. To overcome such difficulties and others of like nature, hybrids have been and are being produced, in the breeding of which such of the native American species were selected as possess the various qualities desired. (See Pls. VIII and IX.) In this work some remarkable successes have been achieved, such, for instance, as *Riparia* × *Rupestris*, No. 101; *Riparia* × *Rupestris*, No. 3306; *Riparia* × *Rupestris*, No. 3309; *Solonis* × *Othello*, No. 1616; *Rupestris* × *Cordifolia*, Nos. 107-11; *Riparia* × (*Cordifolia* × *Rupestris*), Nos. 106-8; *Rupestris* × *Berlandieri*, No. 301A; *Berlandieri* × *Riparia*, No. 420A; and *Monticola* × *Riparia*, No. 18808.

Efforts have also been made to produce hybrids between the *Vinifera* and American native-grape varieties which would be resistant to phylloxera and at the same time give satisfactory crops of fruit of desirable character and quality. (See Pl. X, fig. 2.) By having such direct producers, the cost of grafting would not only be avoided, but congeniality would not have to be reckoned with. Some remarkable strides are being made along this line. A number of these hybrids are under test in the experiment vineyards, but so far it is not possible to say that any of them are better than or equal to some of our finer varieties of native American grapes or that they have as good phylloxera-resistant qualities.

GROWTH RATINGS OF RESISTANT VINES AND DIRECT PRODUCERS.


In Table IV the upper numbers after each name in the columns headed "Experiment vineyard" show the years when the vines were planted; the lower numbers show the growth ratings, which in all cases were made in the autumn of 1913.

The growth or adaptability of each variety at each vineyard where it is under test is expressed in the form of a percentage rating on a scale in which the growth of the variety under conditions for which it is well adapted is taken as the standard of excellence, 100 per cent.




VARIOUS TYPES OF GRAPE ROOT SYSTEMS.

FIG. 1.—Roots of the fleshy type. FIG. 2.—Roots of the shallow or spreading type. FIG. 3.—Roots of the deep-striking type. FIG. 4.—Roots of the oblique type.




LEAVES OF FOUR NATIVE AMERICAN SPECIES OF GRAPES USED AS STOCKS ON WHICH TO GRAFT VINIFERA VARIETIES.

FIG. 1.—*Vitis champini*, upper and lower side of leaf, one-fourth natural size. FIG. 2.—*Vitis doaniana*, upper and lower side of leaf, one-sixth natural size. FIG. 3.—*Vitis candidans*, upper and lower side of leaf, one-fourth natural size. FIG. 4.—*Vitis berlandieri*, upper and lower side of leaf, one-sixth natural size.




LEAVES OF FOUR NATIVE AMERICAN SPECIES OF GRAPES EXTENSIVELY USED AS STOCKS ON WHICH TO GRAFT VINIFERA VARIETIES.

FIG. 1.—*Vitis rupestris*, upper and lower side of leaf, one-fourth natural size. FIG. 2.—*Vitis aestivalis*, upper and lower side of leaf, one-fourth natural size. FIG. 3.—*Vitis labrusca*, upper and lower side of leaf, one-seventh natural size. FIG. 4.—*Vitis riparia*, upper and lower side of leaf, two-fifths natural size.



LEAVES OF FOUR GRAPE HYBRIDS USED AS STOCKS ON WHICH TO GRAFT VINIFERA VARIETIES.

FIG. 1.—*Monticola* × *Riparia* No. 18808, upper and lower side of leaf, one-sixth natural size. FIG. 2.—*Cordifolia* × *Riparia* No. 125-1, upper and lower side of leaf, one-eighth natural size. FIG. 3.—*Berlandieri* × *Riparia* No. 420A, upper and lower side of leaf, one-sixth natural size. FIG. 4.—*Linsecomii* × (*Labrusca* × *Vinifera*), upper and lower side of leaf, one-eleventh natural size.



LEAVES OF FOUR GRAPE HYBRIDS ORIGINATED IN FRANCE AND EXTENSIVELY USED AS STOCKS ON WHICH TO GRAFT VINIFERA VARIETIES.

FIG. 1.—*Riparia* × *Rupestris* No. 3306, upper and lower side of leaf, one-eighth natural size.
FIG. 2.—*Mourvedre* × *Rupestris* No. 1202, upper and lower side of leaf, five-sixteenths natural size.
FIG. 3.—*Riparia* × *Rupestris* No. 101, upper and lower side of leaf, five-fourteenths natural size.
FIG. 4.—*Riparia* × *Rupestris* No. 3309, upper and lower side of leaf, one-third natural size.



FIG. 2.—A DIRECT PRODUCER ON ITS OWN ROOTS (ALICANTE GANZIN).



FIG. 1.—A VINIFERA ON RESISTANT STOCK (HUNISA GRAFTED ON DOG RIDGE).

These adaptability ratings therefore represent the behavior of each variety under the conditions existing at the several vineyards expressed in terms that permit comparison with its behavior elsewhere. They are not based on a comparison with other varieties in the same vineyard. Each variety is therefore rated on a scale based on its own standard of excellence rather than on any arbitrary scale formulated for application to all varieties. It is believed that this method renders possible a truer expression of the reaction of each variety to different soil and climatic conditions than would be possible by using any arbitrary scale of measurement of growth.

To illustrate, Aramon × Rupestris Ganzin, No. 2, planted at each of three experiment vineyards in 1904, at Fresno was rated 97; at Livermore, 83; and at Lodi, only 46. This shows that at Fresno the growth was very satisfactory, and it was therefore rated at 97. At Livermore the growth was good, but not nearly so good as at Fresno; it was therefore rated at 83. At Lodi it made a very poor growth, which as compared with the growth made at Fresno was as 46 to 97, or when compared to perfect growth as 46 to 100, i. e., 46 per cent.

TABLE IV.—Resistant and direct-producing varieties of grapes in eleven experiment vineyards in California, showing the year of planting in each vineyard and the relative growth rating.

Variety.	Experiment vineyard.										
	Chico.	Collax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Adobe Giant:											
Year of planting.....	1906	1906	1903	1904					1903	1907	1907
Growth rating.....	89	93	85	85					90	90	87
(Aestivalis × Monticola) × (Riparia × Rupestris, No. 534-5):											
Year of planting.....	1906	1906	1904	1904			1904		1904	1904	1908
Growth rating.....	96	87	93	92			92		95	89	89
(Aestivalis × Rupestris) × Riparia, No. 227:											
Year of planting.....	1906							1905	1905	1905	
Growth rating.....	82							81	95	89	
Aramon × Rupestris Ganzin, No. 1:											
Year of planting.....	1907	1906	1913	1904		1904	1904	1904	1904	1904	1908
Growth rating.....	99	94		98		96	98	93	98	90	85
Aramon × Rupestris Ganzin, No. 2:											
Year of planting.....	1907	1906	1904	1904	1907	1904	1904	1904	1904	1904	1908
Growth rating.....	93	93	97	97	87	83	46	94	98	92	83
Aramon × Rupestris Ganzin, No. 9:											
Year of planting.....	1906		1904	1904		1904	1905	1904	1904	1904	1907
Growth rating.....	98		95	71		88	94	67	78	94	95
Aramon × Riparia, No. 143A:											
Year of planting.....			1904						1904		1907
Growth rating.....			96						87		95
Arizonica Phoenix:											
Year of planting.....	1912				1905						
Growth rating.....	20				94						
Australis:											
Year of planting.....	1906	1906	1903	1904		1904	1904	1907	1903	1904	1908
Growth rating.....	91	90	97	92		89	90	60	89	92	89
Barnes:											
Year of planting.....	1906	1906	1903		1907	1907		1907	1903		1907
Growth rating.....	95	88	94		95	87		51	99		91
Berlandieri, No. 1:											
Year of planting.....	1906	1906	1904	1904		1904	1904	1904	1904	1904	
Growth rating.....	88	81	88	85		60	72	78	83	84	

TABLE IV.—Resistant and direct-producing varieties of grapes in eleven experiment vineyards in California, showing the year of planting in each vineyard and the relative growth rating—Continued.

Variety.	Experiment vineyard.										
	Chico.	Collax.	Fresno.	Geyserville.	Guasli.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Berlandieri, No. 2:											
Year of planting.....	1906	1906	1904	1906					1909		
Growth rating.....	89	79	65	76					85		
Berlandieri Lafoni, No. 9:											
Year of planting.....	1906		1904	1904		1905	1905	1904	1904	1904	
Growth rating.....	88		71	93		83	66	72	86	77	
Berlandieri × Riparia, No. 33 E. M.:											
Year of planting.....	1907	1906	1904				1904	1904	1904		1907
Growth rating.....	92	83	77				68	70	93		74
Berlandieri × Riparia, No. 34 E. M.:											
Year of planting.....	1906	1906	1903	1904		1907		1904	1903	1907	1907
Growth rating.....	90	90	87	91		85		66	93	87	86
Berlandieri × Riparia, No. 157-11:											
Year of planting.....	1907	1906	1904	1904			1904	1904	1904		1907
Growth rating.....	99	86	81	89			76	75	92		82
Berlandieri × Riparia, No. 420A:											
Year of planting.....	1907	1907	1903	1904	1905	1910	1904	1904	1903	1905	1907
Growth rating.....	97	95	93	75	91	33	93	88	98	92	87
Berlandieri × Riparia, No. 420B:											
Year of planting.....	1912	1907	1904			1907	1907	1904	1904		
Growth rating.....	10	90	85			85	87	64	88		
(Bourisquon × Rupestris, No. 601) × Calicola, No. 13205:											
Year of planting.....	1906		1913	1907		1905		1907	1905		
Growth rating.....	95		91	93		91			94		
Cabernet × Berlandieri, No. 333:											
Year of planting.....			1903	1904		1904		1904	1903		1907
Growth rating.....			86	91		77		91	94		84
Cabernet × Rupestris Ganzin, No. 33A:											
Year of planting.....	1909		1904	1904	1907	1904	1904	1904	1904	1905	
Growth rating.....	85		91	93	85	87	82	83	92	72	
Chasselas × Berlandieri, No. 41B:											
Year of planting.....	1906		1904	1904				1904	1906	1907	1907
Growth rating.....	89		81	94				69	87	84	91
(Cinerea × Rupestris) × Riparia, No. 229:											
Year of planting.....	1906		1905						1905		1908
Growth rating.....	88		87						90		82
Columbaud × Riparia, No. 2502:											
Year of planting.....	1906		1904			1907	1907	1907	1904	1907	1907
Growth rating.....	99		89			74	86	80	89	89	92
Columbaud × Rupestris:											
Year of planting.....						1907			1903	1907	
Growth rating.....						94			96	90	
Constantia:											
Year of planting.....	1907	1907	1905		1905				1904		1907
Growth rating.....	79	94	96		98				99		96
Cordifolia × Riparia, No. 125-1:											
Year of planting.....	1906		1904			1906	1906		1904	1907	
Growth rating.....	90		98			81	84		95	84	
Cordifolia × Rupestris:											
Year of planting.....	1906		1905			1905		1905			
Growth rating.....	89		77			88		60			
Dog Ridge:											
Year of planting.....	1906	1906	1903	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating.....	97	96	89	97	97	94	94	98	97	97	85
Hotporup:											
Year of planting.....	1907		1903	1904		1904	1904	1904	1903	1904	
Growth rating.....	97		83	89		84	93	74	90	92	
Judge:											
Year of planting.....	1906	1906	1903	1907	1907	1906	1907	1907	1903	1907	
Growth rating.....	85	85	93	77	84	66	83	80	90	86	
Lenoir:											
Year of planting.....	1906	1906	1904	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating.....	100	93	77	95	97	86	96	94	98	95	90
Monticola × Riparia, No. 554:											
Year of planting.....	1912		1904	1904		1904	1904	1904	1904	1904	1907
Growth rating.....	40		88	82		84	87	80	98	87	88

TABLE IV.—Resistant and direct-producing varieties of grapes in eleven experiment vineyards in California, showing the year of planting in each vineyard and the relative growth rating—Continued.

Variety.	Experiment vineyard.										
	Chico.	Collax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Monticola × Riparia, No. 18304:											
Year of planting.....	1906	1906	1904	1905	1904	1904	1904	1904	1907
Growth rating.....	95	84	94	90	91	81	89	91	85
Monticola × Riparia, No. 18808:											
Year of planting.....	1907	1906	1904	1905	1904	1904	1904	1904	1907
Growth rating.....	97	90	94	90	88	86	91	93	93
Monticola × Riparia, No. 18815:											
Year of planting.....	1906	1906	1904	1904	1905	1904	1904	1904	1908
Growth rating.....	96	92	95	95	90	86	93	95	92
Motley:											
Year of planting.....	1906	1906	1904	1906	1906	1904	1903	1906
Growth rating.....	90	92	81	76	96	90	86	88
Mourvedre × Rupestris, No. 1202:											
Year of planting.....	1906	1906	1903	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating.....	93	97	100	99	94	98	97	95	100	92	95
Mourvedre × Rupestris, No. 1203:											
Year of planting.....	1906	1907	1904	1906	1906	1906	1906	1906
Growth rating.....	99	91	98	88	81	91	98	80
No name, No. 1:											
Year of planting.....	1905
Growth rating.....	95
No name, No. 2:											
Year of planting.....	1905
Growth rating.....	95
No name, No. 3:											
Year of planting.....	1905
Growth rating.....	94
Pinot Bouschet × Riparia, No. 3001:											
Year of planting.....	1907	1904	1904	1907	1907
Growth rating.....	91	96	90	82	83
Pinot × Rupestris, No. 1305:											
Year of planting.....	1907	1907	1909	1904	1907	1904	1904	1904	1904	1904	1907
Growth rating.....	99	93	99	93	89	96	93	94	98	95	94
Ramsey:											
Year of planting.....	1906	1907	1903	1907	1910	1907	1906	1903
Growth rating.....	90	89	94	88	85	92	94	98
Riparia du Colorado:											
Year of planting.....	1906	1903	1904	1907	1905	1904	1904	1903	1904
Growth rating.....	82	72	83	82	51	91	66	66	71
Riparia France:											
Year of planting.....	1906	1904	1907	1904	1904	1904	1907
Growth rating.....	51	97	84	74	79	86	87
Riparia Gloire:											
Year of planting.....	1906	1906	1903	1905	1904	1907	1903	1904	1907
Growth rating.....	87	86	85	93	82	77	88	85	75
Riparia Grand Glabre:											
Year of planting.....	1909	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating.....	56	92	85	82	82	74	86	89	83
Riparia Martineau:											
Year of planting.....	1906	1904
Growth rating.....	92	86
Riparia Ramond:											
Year of planting.....	1912	1905	1906
Growth rating.....	60	92	86
Riparia Selected:											
Year of planting.....	1905
Growth rating.....	89
Riparia × Berlandieri, No. 161-49:											
Year of planting.....	1906	1907	1907	1904
Growth rating.....	95	91	87	90
Riparia × (Cordifolia × Rupestris), No. 106-8:											
Year of planting.....	1906	1906	1910	1904	1904	1904	1904	1910	1904	1907
Growth rating.....	88	86	93	86	76	85	72	92	90	91
Riparia Grand Glabre × Aramon											
Rupestris, No. 4110:											
Year of planting.....	1906	1904	1904	1904	1904	1904
Growth rating.....	73	89	92	91	91	91
Riparia × Rupestris, No. 101:											
Year of planting.....	1906	1906	1904	1904	1904	1904	1904	1904	1904	1907
Growth rating.....	85	81	95	96	91	97	65	93	89	91

TABLE IV.—Resistant and direct-producing varieties of grapes in eleven experiment vineyards in California, showing the year of planting in each vineyard and the relative growth rating—Continued.

Variety.	Experiment vineyard.										
	Chico.	Colfax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Riparia × Rupestris, No. 101-14:											
Year of planting	1906	1906	1904	1904	1905	1904	1904	1904	1904	1904	1907
Growth rating	92	92	93	90	72	90	98	86	94	90	94
Riparia × Rupestris, No. 3306:											
Year of planting	1906	1906	1904	1904	1907	1904	1904	1904	1904	1904	1907
Growth rating	94	92	74	96	89	77	80	91	93	93	93
Riparia × Rupestris, No. 3309:											
Year of planting	1906	1906	1904	1904	1907	1904	1904	1904	1903	1904	1907
Growth rating	89	95	95	97	88	79	94	88	92	91	70
Riparia × Rupestris, No. 108-103:											
Year of planting		1906		1906					1905	1906	1907
Growth rating		90							93	91	82
Riparia × Rupestris de Jaeger:											
Year of planting	1909	1906	1904		1907		1907		1904	1904	
Growth rating	67	91	94		86		70		91	87	
Riparia × (Rupestris × Aramon) Jaeger, No. 201:											
Year of planting	1906	1906	1904	1904	1907	1904	1904	1904	1904	1904	1907
Growth rating	99	94	85	95	82	95	93	67	96	89	95
Riparia × Rupestris Ramond:											
Year of planting	1906								1904		
Growth rating	87								77		
Rupestris des Caussettes:											
Year of planting	1906	1906	1903	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating	95	94	91	90	92	85	90	78	91	89	90
Rupestris des Semis, No. 81-2:											
Year of planting	1907	1906	1904			1907		1907	1904		1907
Growth rating	100	90	96			96		81	96		88
Rupestris Ganzin:											
Year of planting	1906								1906		
Growth rating	86								93		
Rupestris le Reux:											
Year of planting	1906				1905				1906		
Growth rating	93				89				91		
Rupestris Martin:											
Year of planting	1906	1907	1903	1904	1905	1904	1904	1903	1903	1904	1907
Growth rating	92	83	77	95	79	92	92	75	92	90	89
Rupestris Metallica:											
Year of planting	1906	1906	1903	1904	1905	1904	1904	1904	1903	1906	
Growth rating	99	92	90	95	95	93	93	82	97	89	
Rupestris Mission:											
Year of planting	1906	1906	1904	1904		1904	1904	1904	1903	1904	1907
Growth rating	91	86	79	85		88	87	74	85	81	81
Rupestris Othello:											
Year of planting	1906		1904			1907			1904		1907
Growth rating	98		87			65			92		81
Rupestris Pillans:											
Year of planting	1906	1906	1905	1906	1905	1905	1905	1905	1905	1905	1907
Growth rating	99	92	96	83	95	92	93	80	87	82	87
Rupestris St. George:											
Year of planting	1906	1906	1903	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating	94	99	96	97	95	98	98	87	100	87	97
Rupestris × Berlandieri, No. 219A:											
Year of planting	1907	1906	1902	1904	1905	1904	1904	1904	1903	1904	1907
Growth rating	75	90	82	85	85	88	91	73	83	86	93
Rupestris × Berlandieri, No. 301A:											
Year of planting	1907	1906	1903	1904		1904		1907	1903	1904	1907
Growth rating	100	83	85	92		85		88	92	86	84
Rupestris × Berlandieri, No. 301B:											
Year of planting	1906	1907		1904			1905	1905			
Growth rating	92	94		94			97	87			
Rupestris × Berlandieri, No. 301-37-152:											
Year of planting	1906	1907	1904			1906		1907	1904		
Growth rating	91	89	90			84		82	88		
Rupestris × Chasselas Rose, No. 4401:											
Year of planting	1906	1906							1907		
Growth rating	91	96							98		
Rupestris × Cinerea:											
Year of planting	1909	1907	1904					1907	1904		1907
Growth rating	60	89	96					85	92		92

TABLE IV.—Resistant and direct-producing varieties of grapes in eleven experiment vineyards in California, showing the year of planting in each vineyard and the relative growth rating.—Continued.

Variety.	Experiment vineyard.										
	Chico.	Collax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Rupestris × Cordifolia, No. 107-11:											
Year of planting.....	1906	1907	1904	1905	1904	1905	1905	1905
Growth rating.....	71	89	89	84	94	71	89	85
Rupestris × (Cordifolia × Rupestris), No. 202:											
Year of planting.....	1907	1907	1904	1907	1904	1904	1907
Growth rating.....	68	91	90	82	85	87	77
Rupestris × (Cordifolia × Rupestris), No. 202-5:											
Year of planting.....	1906	1904	1906	1906	1907	1906
Growth rating.....	90	90	89	50	78	90
Rupestris × Hybrid Azemar, No. 215:											
Year of planting.....	1907	1907	1904	1905	1904	1904	1905	1904	1904
Growth rating.....	93	85	89	70	74	83	70	91	92
Rupestris × Petit Bouschet, No. 503:											
Year of planting.....	1906	1904
Growth rating.....	95	91
Rupestris × Petit Bouschet Jaeger, No. 504:											
Year of planting.....	1906	1904	1906	1907	1907
Growth rating.....	96	96	90	94	92
Rupestris × Riparia, No. 108-16:											
Year of planting.....	1912	1904	1904	1904	1904
Growth rating.....	75	91	78	68	81
Salt Creek:											
Year of planting.....	1906	1906	1903	1904	1905	1904	1904	1903	1904
Growth rating.....	91	87	86	85	77	92	68	90	84
Solonis Robusta:											
Year of planting.....	1906	1906	1903	1904	1905	1904	1904	1904	1903	1905	1907
Growth rating.....	84	92	92	93	92	83	88	90	92	90	87
Solonis × (Cordifolia × Rupestris), No. 202-4:											
Year of planting.....	1909	1907	1904	1906	1906	1906
Growth rating.....	60	93	84	76	84	61
Solonis × Othello:											
Year of planting.....	1906	1907	1904	1904	1904	1904	1904
Growth rating.....	99	93	96	87	88	88	95
Solonis × Othello, No. 1613:											
Year of planting.....	1906	1906	1903	1905	1906	1907	1903	1907
Growth rating.....	99	96	100	95	90	92	100	94
Solonis × Riparia, No. 1615:											
Year of planting.....	1906	1906	1904	1904	1907	1904	1904	1907
Growth rating.....	93	90	96	91	83	86	89	89
Solonis × Riparia, No. 1616:											
Year of planting.....	1906	1906	1904	1904	1907	1904	1907	1904	1904	1904	1907
Growth rating.....	90	91	94	91	86	80	86	80	98	88	94
Taylor Narbonne:											
Year of planting.....	1906	1907	1904	1904	1907	1904	1904	1904	1904	1904	1907
Growth rating.....	70	82	89	95	78	70	51	71	91	85	86
Texas:											
Year of planting.....	1905
Growth rating.....	79
Tisserand:											
Year of planting.....	1906	1904	1904	1904
Growth rating.....	93	84	78	92
Viala:											
Year of planting.....	1906	1907	1904	1904	1903	1907	1907
Growth rating.....	92	86	89	71	97	82	87
Viala × Riparia:											
Year of planting.....	1904	1904	1904	1905	1904	1905
Growth rating.....	77	70	77	78	69	82
Vitis candicans:											
Year of planting.....	1904	1904
Growth rating.....	65	89

A number of the resistant-stock varieties have been growing a sufficient length of time to show what may be expected of them under similar conditions. In the following list of stocks that are worthy of special mention as having made excellent growth ratings at each of eleven California experiment vineyards, the varieties are given in the order of their ratings, i. e., the best growers first, and so on:

Chico Varietal Vineyard.—Lenoir; Rupestris de Semis, No. 81-2; Rupestris × Berlandieri, No. 301A; Aramon × Rupestris Ganzin, No. 1; Berlandieri × Riparia, No. 157-11; Columbaud × Riparia, No. 2502; Rupestris Metallica; Rupestris Pillans; Solonis × Othello; Solonis × Othello, No. 1613; Aramon × Rupestris Ganzin, No. 9; Rupestris Othello; Berlandieri × Riparia, No. 420A; Dog Ridge; Hotporup; Monticola × Riparia, No. 18808; (Aestivalis × Monticola) × (Riparia × Rupestris, No. 554-5); Barnes.

Colfax Experiment Vineyard.—Rupestris St. George; Mourvedre × Rupestris, No. 1202; Dog Ridge; Solonis × Othello, No. 1613; Berlandieri × Riparia, No. 420A; Riparia × Rupestris, No. 3309; Aramon × Rupestris Ganzin, No. 1; Constantia; Rupestris des Caussettes; Rupestris × Berlandieri, No. 301B; Adobe Giant; Aramon × Rupestris Ganzin, No. 2; Lenoir; Solonis × (Cordifolia × Rupestris), No. 202-4; Solonis × Othello; Monticola × Riparia, No. 18815; Motley; Riparia × Rupestris, No. 101-14.

Fresno Experiment Vineyard.—Mourvedre × Rupestris, No. 1202; Solonis × Othello, No. 1613; Cordifolia × Riparia, No. 125-1; Aramon × Rupestris Ganzin, No. 2; Australis; Riparia France; Constantia; Rupestris des Semis, No. 81-2; Rupestris Pillans; Rupestris St. George; Solonis × Riparia, No. 1615; Aramon × Rupestris Ganzin, No. 9; Monticola × Riparia, No. 18815; Riparia × Rupestris, No. 101; Riparia × Rupestris, No. 3309; Barnes; Monticola × Riparia, No. 18804; Monticola × Riparia No. 18808.

Geyserville Experiment Vineyard.—Mourvedre × Rupestris, No. 1202; Aramon × Rupestris Ganzin, No. 1; Aramon × Rupestris Ganzin, No. 2; Dog Ridge; Riparia × Rupestris, No. 3309; Rupestris St. George; Riparia × Rupestris, No. 101; Riparia × Rupestris, No. 3306; Solonis × Othello; Lenoir; Rupestris Martin; Rupestris Metallica; Taylor Narbonne; Rupestris × Berlandieri, No. 301B; Berlandieri Lafont, No. 9; Solonis Robusta; (Aestivalis × Monticola) × (Riparia × Rupestris, No. 554-5); Australis.

Guasti Experiment Vineyard.—Constantia; Dog Ridge; Lenoir; Barnes; Rupestris Metallica; Rupestris Pillans; Rupestris St. George; Solonis × Othello, No. 1613; Arizonica Phoenix; Mourvedre × Rupestris, No. 1202; Riparia Gloire; Riparia Ramond; Rupestris des Caussettes; Solonis Robusta; Berlandieri × Riparia, No. 420A; Pinot × Rupestris, No. 1905; Riparia × Rupestris, No. 3306; Rupestris le Reux.

Livermore Experiment Vineyard.—Mourvedre × Rupestris, No. 1202; Rupestris St. George; Aramon × Rupestris Ganzin, No. 1; Rupestris des Semis, No. 81-2; Monticola × Riparia, No. 18815; Riparia × (Rupestris × Aramon) Jaeger, No. 201; Dog Ridge; Rupestris Metallica; Rupestris Martin; Rupestris Pillans; Riparia × Rupestris, No. 101; Monticola × Riparia, No. 18804; Monticola × Riparia, No. 18808; Riparia × Rupestris, No. 101-14; Solonis × Othello, No. 1613; Australis; Aramon × Rupestris Ganzin, No. 9; Cordifolia × Rupestris.

Lodi Experiment Vineyard.—Aramon × Rupestris Ganzin, No. 1; Riparia × Rupestris, No. 101-14; Mourvedre × Rupestris, No. 1202; Riparia × Rupestris, No. 101; Lenoir; Motley; Aramon × Rupestris Ganzin, No. 9; Dog Ridge; Riparia × Rupestris,

No. 3309; Rupestris \times Cordifolia, No. 107-11; Berlandieri \times Riparia, No. 420A; Riparia \times (Rupestris \times Aramon) Jaeger, No. 201; Rupestris Metallica; Rupestris Pillans; (Aestivalis \times Monticola) \times (Riparia \times Rupestris, No. 554-5); Ramsey; Rupestris Martin; Salt Creek.

Mountain View Experiment Vineyard.—Dog Ridge; Mourvedre \times Rupestris, No. 1202; Lenoir; Monticola \times Riparia, No. 18804; Aramon \times Rupestris Ganzin, No. 1; Aramon \times Rupestris Ganzin, No. 2; Riparia \times Rupestris, No. 3306; Motley; Solonis Robusta; Berlandieri \times Riparia, No. 420A; Riparia \times Rupestris, No. 3309; Rupestris \times Berlandieri, No. 301A; Solonis \times Othello; Riparia \times Berlandieri, No. 161-49; Rupestris St. George; Rupestris \times Berlandieri, No. 301B; Monticola \times Riparia, No. 18808; Monticola \times Riparia, No. 18815.

Oakville Experiment Vineyard.—Mourvedre \times Rupestris, No. 1202; Rupestris St. George; Solonis \times Othello, No. 1613; Barnes; Constantia; Aramon \times Rupestris Ganzin, No. 1; Aramon \times Rupestris Ganzin, No. 2; Berlandieri \times Riparia, No. 420A; Lenoir; Monticola \times Riparia, No. 554; Ramsey; Solonis \times Riparia, No. 1616; Dog Ridge; Rupestris Metallica; Viala; Rupestris des Semis, No. 81-2; (Aestivalis \times Monticola) \times (Riparia \times Rupestris, No. 554-5); (Aestivalis \times Rupestris) \times Riparia, No. 227.

Sonoma Experiment Vineyard.—Dog Ridge; Lenoir; Monticola \times Riparia, No. 18815; Solonis \times Othello; Aramon \times Rupestris Ganzin, No. 9; Monticola \times Riparia, No. 18808; Riparia \times Rupestris, No. 3306; Aramon \times Rupestris Ganzin, No. 2; Australis; Berlandieri \times Riparia, No. 420A; Mourvedre \times Rupestris, No. 1202; Monticola \times Riparia, No. 18804; Riparia \times Rupestris, No. 3309; Riparia \times Rupestris, No. 108-103; Adobe Giant; Aramon \times Rupestris Ganzin, No. 1; Riparia \times (Cordifolia \times Rupestris), No. 106-8; Riparia \times Rupestris, No. 101-14.

Stockton Experiment Vineyard.—Rupestris St. George; Constantia; Aramon \times Rupestris Ganzin, No. 9; Mourvedre \times Rupestris, No. 1202; Riparia \times Rupestris, No. 101-14; Solonis \times Othello, No. 1613; Solonis \times Riparia, No. 1616; Monticola \times Riparia, No. 18808; Columbaud \times Riparia, No. 2502; Monticola \times Riparia, No. 18815; Rupestris \times Cinerea; Barnes; Riparia \times (Cordifolia \times Rupestris), No. 106-8; Riparia \times Rupestris, No. 101; Lenoir; Riparia \times Rupestris, No. 3306; Rupestris des Causettes; Rupestris \times Berlandieri, No. 219A.

Table V gives the resistant varieties in each vineyard which are estimated to have made the best and most creditable growth records as compared to all the varieties under test. The numbers in line with each name in the vineyard columns show the relative growth rating made by the variety in the respective vineyards where it is under test. The highest rating is expressed by the figure 1, the next by 2, and so on. The ratings therefore represent the behavior of each variety under the conditions existing at the several vineyards, expressed in terms that permit comparison with its behavior elsewhere, and in comparison also with other varieties in the same vineyard. To illustrate: Of all the resistant varieties at Livermore the best record was made by Mourvedre \times Rupestris, No. 1202 (rated as 1), whereas at Stockton it was fourth best (expressed by 4), and at Sonoma eleventh best (expressed by 11).

TABLE V.—Resistant-stock varieties of grapes making the best growth records, showing their relative merits in each of eleven experiment vineyards in California.

Variety.	Chico.	Colfax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Adobe Giant		11								15	
Aramon × Rupestris Ganzin:											
No. 1	4	7		2		3	1	5	6	16	
No. 2		12	4	3				6	7	8	
No. 9	11		12			17	7			5	3
Australis				18		16				9	
Barnes	18		16		4				4		12
Berlandieri × Riparia:											
No. 420 A	13	5			15		11	1	8	10	
No. 157-11		5									
Columbaud × Riparia, No. 2502		6									9
Constantia		8			1				5		2
Cordifolia × Riparia, No. 125-1			3								
Dog Ridge	14	3		4	2	7	2	1	13	1	
Lenoir	1	13		10	3		5	3	9	2	15
Monticola × Riparia:											
No. 18804			17			12		4		12	
No. 18808	16		18			13		17		6	8
No. 18815		16	13			5		18		3	10
Motley		2					6	8			
Mourvedre × Rupestris, No. 1202		17	1	1	10	1	3	2	1	11	4
Ramsay							16		11		
Riparia Gloire					11						
Riparia × Rupestris:											
No. 101			14	7		11	4				14
No. 101-14		18				14	2			18	5
No. 3306				8	17			7		7	16
No. 3709		6	15	5			9	11		13	
Rupestris des Semis, No. 81-2	2		8			4			16		
Rupestris Martin				11		9	17				
Rupestris Metallica	7			12	5	8	13		14		
Rupestris Pillans	8		9	6	6	10	14				
Rupestris St. George		1	10	6	7	2		15	2		1
Rupestris × Berlandieri:											
No. 301A	3							12			
No. 301B		10		14				16			
Rupestris Chirezi							10				11
Rupestris × Cordifolia, No. 107-11											
Solomis Robusta				16	14			9			
Solomis × Othello	9	15		9				13		4	
Solomis × Othello, No. 1613	10	4	2		8	15			3		6

Table VI gives an alphabetical list of improved American native and Franco-American grape varieties which are being tested on their own roots in the Chico, Colfax, Fresno, Geyserville, Guasti, Livermore, Lodi, Mountain View, Oakville, Sonoma, and Stockton experiment vineyards. The plantings are too young to permit the drawing of conclusions.

TABLE VI.—Varieties of American native and Franco-American grapes under test on their own roots at eleven experiment vineyards in California.

[The locations of the tests are indicated by plus (+) marks.]

Varieties.	Chico.	Colfax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Agawam	+	+									
Albania			+								
Alexander Winter	+								+		
Alicante Ganzin	+	+				+					
Alicante × Rupestris Terrace, No. 20	+			+		+					+
Alice	+								+		
Amber Queen	+										
Ambrosia	+										
Amerbonte								+			

TABLE VI.—Varieties of American native and Franco-American grapes under test on their own roots at eleven experiment vineyards in California—Continued.

Varieties.	Chico.	Colfax.	Fresno.	Geyserville.	Guasti.	Livermore.	Lodi.	Mountain View.	Oakville.	Sonoma.	Stockton.
Lenoir.....	++										+
Lindley.....	++							+			
Linn.....	++										
Lindmar.....	++								+		
Little Blue.....	++										
Livingston.....	++										
Long John.....	++										
Louisiana.....	++	+	+								
Lucile.....	++										
Lukfata.....	++										
Manito.....	++		+					+		+	
Marguerite.....	++	+	+		+						
Martha.....	++										
Mary Favorite.....	++										
Massasoit.....	++										
Maxatawney.....	++										
Mericadel.....	++		+					+	+		
Merrimac.....	++										
Missouri Riesling.....	++										
Montefiore.....	++										
Moore.....	++	+	+	+		+	+	+	+	+	
Moyer.....	++										
Mrs. Munson.....	++				+						
Muench.....	++	+									
Niagara.....	++			+		+	+	+		+	
Noah.....	++										
Oliarato.....	++	+									
Osage.....	++										
Olita.....	++										
Paradox.....	++										
Paragon.....	++										
Pardes.....	++			+			+				+
Peabody.....	++										
Pearl.....	++								++		
Perkins.....	++										
Pierce.....	++		+	+		+		+	++	+	
Plant de Carmes.....	++		+					+			
Plant de Gounay.....	++								++	+	
Pocklington.....	++										
Fresley.....	++								+		
Ragon.....	++	+			+						
Rebecca.....	++			+			+				
Red Eagle.....	++										
Regal.....	++										
Requa.....	++										
Rockford.....	++										
Rogers:											
No. 5.....	++										
No. 13.....	++								+		
No. 32.....	++										
Rommel.....	++								+		+
Siebel No. 14.....	++			+							
Shalah.....	++						+				
Wyoming Red.....	++										

To grow vines on resistant stocks successfully, it should be borne in mind that the resistance of vines depends upon the inherent characters of the vine and its adaptation to soil, climatic, and other conditions, and that the resistant quality of the stock is very materially affected by the congeniality of the varieties grafted on it.

CONGENIALITY AND ADAPTABILITY OF VINES.

Two vine varieties are congenial to each other if both top and root flourish when one is grafted on the other. (See Pl. X, fig. 1.) The congeniality would be called perfect when one variety grafted on another behaves as if the stock were grafted with a scion of itself, the union being perfect and the behavior of the vine the same as that of an entire ungrafted plant.

The term "congeniality" as used in this discussion is limited to the relation of vine varieties to the resistant stocks upon which they are grafted. To discriminate properly between adaptability and congeniality and then to determine the congeniality, it is necessary to know the behavior of the resistant varieties as well as the Vinifera varieties on their own roots. If we have grafted vines of which both the stock and the scion varieties are known to be suited to the soil and climatic conditions and they do not respond, we know that congeniality is lacking.

The adaptability of varieties can be closely forecasted, but their congeniality must be determined by actual tests. Without knowledge of its adaptability to existing conditions, the extent to which differences in the behavior of a variety grafted on different stocks are due to congeniality or to adaptability is impossible of determination.

Saccharine and acid determinations of the fruit from grafted vines have been made for a number of years with a view to ascertain whether the quality of the fruit is influenced by the stock upon which the vine is grafted. (See Pl. X.) Such determinations contrasted with the same season's growth ratings of the same vines indicate a close correspondence between these important chemical constituents of the fruit and the congeniality of grafts and stocks as determined by observations, and they afford a useful check on the congeniality ratings.

Similar growth ratings of a variety grafted on various stocks are found to be accompanied by fairly definite percentages of sugar and acid. Under like conditions of growth the sweetness and the acidity of the fruit, as well as its time of ripening, are materially influenced by the congeniality of the scion and stock. The saccharine and acid contents are two of the leading considerations in the money value of the fruit.

In determining the relative congeniality of vine varieties on diverse resistant stocks, these and the relative quantity of fruit produced, the difference in time of ripening, the relative healthfulness and comparative durability of varieties on different resistant stocks, and the relative amount of wood produced are some of the considerations that appear most important.

BEHAVIOR OF GRAPES GRAFTED AND ON THEIR OWN ROOTS.

In Table VII, column 1 gives (1) the variety name, and indented under it (2) the name of the resistant stock on which it is grafted, or if the variety is on its own roots the fact is so stated. Column 2 shows the experiment vineyard in which the growth was tested, use being made of the following abbreviations: C for Chico, Cx for Colfax, F for Fresno, G for Geyserville, Gi for Guasti, L for Lodi, Li for Livermore, M for Mountain View, O for Oakville, S for Sonoma, St for Stockton. Column 3 shows the year in which the stock was

planted, thereby indicating its age, and column 4 gives the year of grafting. Column 5 shows the congeniality, or the growth of each variety on the different stocks, expressed in the form of a percentage rating, on a scale in which the growth of the variety when not grafted but growing as an entire plant on its own root under conditions to which it is well adapted is taken as the standard of excellence, that is, 100 per cent. These ratings therefore represent the behavior of each variety grafted on the several stocks under the conditions existing at the vineyard at which it was found, expressed in terms that permit comparison with its behavior when growing as an entire plant on its own roots under favorable conditions and not based on a comparison with other *Vinifera* varieties grafted on the same stock in the same vineyard. The rating in each case is the average rating made in different seasons to and including the autumn of 1913. To illustrate: Alicante Bouschet, grafted in the Oakville vineyard in 1906 on different resistant stocks, on Aramon \times Rupestris Ganzin, No. 1, was rated at 91; on Riparia \times Rupestris, No. 3309, at 88; on Mourvedre \times Rupestris, No. 1202, at 72; and on Riparia \times Rupestris, No. 101, at 55. This shows that Alicante Bouschet, which is well adapted to the conditions there, when grafted on these different stocks at the same time, under the same conditions, in the same vineyard, and with the same treatment, varied in growth and behavior in comparison with the same variety on its own roots in accordance with the above ratings. Column 6 gives the pruning method, s being used for spurs and c for canes. Column 7 gives the weight of prunings per vine; 8, the nodes bearing fruit; 9 and 10, the growth-starting dates in early and late seasons; 11 and 12, the blossoming dates in early and late seasons; 13 and 14, the fruit-setting dates in early and late seasons; 15 and 16, the fruit-ripening dates in early and late seasons. Columns 17 to 21 give the fruit per vine for the seasons from 1909 to 1913, inclusive; 22, the average percentage of sugar, Balling scale; 23, the average acid, as tartaric, per 100 c. c. Column 24 shows the size of the clusters, m indicating medium; m-l, medium to large; l, large; v, very; s, small. Column 25 shows the shape of the clusters, whether round (r), cylindrical (cy), long (l), or tapering (t). Column 26 designates the density of the clusters, whether compact (c), medium (m), or loose (l). Column 27 shows the size of the berry, whether large (l), medium (m), or small (s). Column 28 gives the shape of the berry, whether round (r), oval (o), or oblong (ob). Column 29 shows the color of the berry, whether black (b), red (r), or white (w). Column 30 indicates the purpose for which the fruit is used, whether for table (t), shipping (s), juice (j), wine (w), or storage (st). In this table the nomenclature of varieties has been brought into conformity with the code of the American Pomological Society in so far as it has appeared practicable.

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year planted.	Year grafted.	Congenality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Ach-I-Soumi:															
Dog Ridge.	O	1904	1910	P. ct. 92	C, S	6.5		Mar. 24		May 30	June 3				
Lenoir.	O	1904	1910							May 31	June 6	June 20			
Actoni Macaron:															
Lenoir.	O	1904	1907	89	S	2.9	3 to 4	Mar. 24		May 20	May 24	June 24			
Monticola X Rupestris.	F	1907	1907	91	S	2 to 5	2 to 5			May 20	May 24	June 29			
Mourvedre X Rupestris, No. 1202.	F	1907	1907	92	C, S	5.5	2 to 5	Mar. 21		May 20	May 24	June 24			
Solonis X Riparia, No. 1616.	F	1907	1907	90	C, S	3	2 to 5	Mar. 14	Apr. 1	May 15	May 24	June 15			
Actonky:															
Lenoir.	O	1904	1907	89	S	3.3	2 to 5	Mar. 24		May 23	June 12	June 29			
Monticola X Rupestris.	F	1907	1907	89	C, S	3.7	4 to 5	Mar. 21	Apr. 1	May 19	May 23	June 31			
Molley.	F	1907	1907	87	C, S	3	3 to 5	Mar. 23	Mar. 28	May 18	June 1	June 1			
Admirable:															
Riparia X Rupestris, No. 3309.	O	1904		91	C, S	6.5	2 to 4	Mar. 22		May 15	June 2	June 19			
Afeenthaler:															
Own roots.	GI	1904		83	S			Mar. 14	Mar. 20	May 26	June 4	June 16			
Agada:															
Riparia X Rupestris, No. 3309.	O	1904	1910	96	C, S	2		Mar. 24		May 25	June 1	June 6	June 19		
Ahmeur bon Ahmeur:															
Lenoir.	O	1904	1907	88	S	2.8	3 to 6	Mar. 24		May 20	June 7	June 25	June 30		
Ajaki Otila:															
Mourvedre X Rupestris, No. 1202.	F	1907	1907	97	C, S	4.5	4 to 6	do.	Apr. 3	May 18	May 23	May 29	Sept. 3		
Riparia X Rupestris, No. 101.	F	1907	1907	88	C, S	3	3 to 5	Mar. 20	Mar. 30	May 17	May 21	May 27	Sept. 3		
Rupestris Metallica.	F	1907	1907	85	S	3	3 to 4	do.	Apr. 2	May 15	May 23	May 24	Sept. 6		
Ajmi:															
Lenoir.	O	1906	1909	88	S	1.5		Mar. 27		June 8	June 13	June 17	June 20		
Aleatico:															
Own roots.	GI	1904	1906	90	S	3.5	2 to 5	Mar. 14	Mar. 25	May 29	May 30	June 2	June 5	Sept. 3	Sept. 23
Dog Ridge.	O	1904	1906	91	C, S	3	2 to 4	Mar. 20	Mar. 25	May 23	June 1	June 28	June 17	Sept. 24	Sept. 6
Lenoir.	O	1903	1905	88	S	3.8	2 to 4	Mar. 16	Mar. 16	May 21	June 3	May 26	June 10	Oct. 5	Oct. 5
Rupestris St. George.	O	1903	1905	93	C, S	3	2 to 4	Mar. 18	Mar. 18	May 20	do.	June 1	June 17	Sept. 22	Oct. 2
Taylor Narbonne	O	1904	1906	81	S	3.2	2 to 5	Mar. 17	Mar. 17	May 25	June 8	May 30	June 15	Oct. 4	Oct. 4

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Alexandria:															
Own roots.....	G						6								
Adobe Giant.....	F	1904		78	8	1.3	2 10 5	9	10	11	12	13	14	15	16
Aramoun × Rupestris Gazin, No. 1.....	F	1904	1906	86	8	1.5	3 10 5	Mar. 13	Apr. 1	May 23	May 26	June 2	June 2	Sept. 16	Sept. 24
Aramoun × Rupestris Gazin, No. 2.....	F	1907	1909	83	8	2.5		Apr. 1	Apr. 1	May 15	May 24	May 29	June 1	Sept. 12	Sept. 22
Australis.....	F	1907	1909	88	8	3		do.	do.	May 22	May 22	Oct. 16	Oct. 16	Sept. 15	Sept. 27
Berlandieri × Riparia, No. 420A.....	F	1907	1909	89	8	2		Mar. 24	do.	do.	do.	do.	do.	Sept. 15	Oct. 16
Constandia.....	F	1907	1909	87	8	2.5		Mar. 23	Mar. 23	May 25	May 29	May 28	June 11	Sept. 8	Sept. 20
Log Ridge.....	F	1904	1906	90	3	2.3	2 10 5	Mar. 11	Apr. 6	May 21	May 30	do.	June 8	do.	Do.
Leonor.....	F	1903	1905	82	8	1.3	3 10 5	do.	Apr. 1	May 12	May 29	May 21	June 5	do.	Sept. 15
Monticola × Riparia, No. 18808.....	F	1907	1909	86	8	3.6		Mar. 18	do.	May 20	May 20	May 24	June 8	Oct. 16	do.
Monticola × Rupestris.....	F	1907	1909	86	8	3.6		Mar. 27	Apr. 3	May 25	June 1	May 24	June 10	do.	Sept. 20
Monvredo × Rupestris, No. 1202.....	F	1904	1906	86	8	2.5	3 10 5	Mar. 13	Apr. 1	May 14	May 30	May 25	June 9	Sept. 4	Sept. 20
Riparia Gloire.....	F	1903	1905	83	8	4	3 10 6	Mar. 14	Apr. 1	May 13	May 30	May 25	June 9	Sept. 9	Sept. 20
Riparia × (Cardinalia × Rupestris, No. 106-S).....	F	1907	1909	87	8	5.5		Mar. 24	Apr. 5	May 20	May 31	May 28	June 7	Oct. 16	Sept. 20
Riparia × Rupestris, No. 101.....	F	1904	1906	89	8	2	2 10 5	Mar. 12	Apr. 5	do.	May 26	May 28	June 7	Oct. 16	Sept. 20
Riparia × Rupestris, No. 101-14.....	F	1907	1909	89	8	3.5		Mar. 25	Apr. 5	May 26	May 26	May 28	June 5	Sept. 5	Sept. 20
Riparia × Rupestris, No. 3306.....	F	1907	1909	87	8	2.5		Mar. 31	Apr. 5	May 18	May 29	May 28	June 5	Sept. 5	Sept. 20
Riparia × Rupestris, No. 3309.....	F	1904	1906	91	8	4	1 10 5	Mar. 13	Apr. 5	May 23	May 29	May 28	June 5	Sept. 5	Sept. 20
Rupestris Martin.....	F	1903	1905	85	8	1.6	2 10 5	Mar. 8	Apr. 1	May 11	May 31	May 27	do.	do.	Do.
Rupestris St. George.....	F	1903	1905	89	8	2	2 10 5	Mar. 14	do.	May 15	May 28	May 26	do.	do.	Do.
Rupestris × Berlandieri, No. 219A.....	F	1907	1908	87	8	3		Apr. 1	Apr. 1	May 22	May 29	May 21	June 3	Sept. 10	Sept. 20
Salt Creek.....	F	1903	1905	86	8	1.6	3 10 6	Mar. 7	Apr. 1	May 16	May 29	May 21	June 3	Sept. 10	Sept. 20
Solanis Robusta.....	F	1907	1909	84	8	1		Mar. 29	May 22	May 25	May 30	May 27	June 8	Oct. 16	Sept. 20
Solanis × Othello, No. 1613.....	F	1907	1909	86	8	3		Mar. 8	Apr. 5	May 22	May 29	May 27	June 4	Oct. 16	Sept. 20
Solanis × Riparia, No. 1616.....	F	1904	1906	88	8	2.3	2 10 5	Mar. 14	Apr. 1	May 13	May 29	May 20	June 4	Sept. 4	Sept. 20
Taylor Narbonne.....	F	1904	1906	85	8	1.4	3 10 5	Mar. 13	Apr. 1	May 13	May 29	do.	do.	Sept. 15	Sept. 20
Viala.....	F	1907	1911	80	8	1		Mar. 25	Apr. 1	May 20	May 29	do.	do.	Sept. 15	Sept. 20

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Aramon:															
Aramon × Rupestris Ganzin, No. 1.															
Australis.....	O	1904	1906	P. cl.	S	Lbs.	3 to 5	Mar. 19	May 22	June 3	May 28	June 17	Sept. 28	Oct. 20	
Barnes.....	F	1907	1907		C, S	3.1	3 to 5	Mar. 18	May 5	May 21	May 27	June 17	Sept. 17	Sept. 21	
Barnes.....	F	1907	1907		S	1.9	3 to 5	Mar. 18	May 14	May 22	May 27	June 27	Sept. 10	Do.	
Cornucopia.....	F	1907	1907		C, S	3	3 to 5	Mar. 12	May 10	May 21	May 23	June 23	do.	Sept. 17	
De Graisset.....	O	1907	1907		C, S	3	3 to 5	Mar. 12	May 10	May 21	May 23	June 23	Sept. 20	Oct. 17	
Dog Ridge.....	O	1903	1905		S	3.1	3 to 6	Mar. 18	May 22	June 10	May 27	June 16	Sept. 29	Oct. 25	
Herbmont.....	O	1903	1905		S	3.3	3 to 6	Mar. 18	May 22	June 10	May 27	June 16	Oct. 5	Oct. 12	
Hotopop.....	O	1907	1907		C, S	3.3	3 to 6	Mar. 18	May 22	June 10	May 27	June 16	Oct. 17	Oct. 15	
Lenoir.....	O	1907	1907		S	3.7	3 to 5	Mar. 22	May 10	May 21	May 27	June 15	Sept. 27	Oct. 11	
Monticola × Rupestris.....	O	1907	1907		C, S	3.5	3 to 5	Mar. 15	May 11	May 21	May 28	June 20	Sept. 16	Sept. 21	
Mourvedre × Rupestris, No. 1202.....	O	1904	1906		S	2.8	3 to 5	Mar. 20	May 22	June 3	May 28	June 20	Sept. 16	Sept. 21	
Ponroy.....	O	1907	1907		S	3.07	3 to 6	Mar. 19	May 11	May 23	May 27	June 20	Sept. 16	Sept. 21	
Ponroy.....	F	1907	1907		S	3.5	3 to 5	Mar. 25	May 14	May 20	May 24	June 24	Sept. 16	Sept. 21	
Ramsey × Rupestris, No. 101.....	F	1907	1907		C, S	2.5	3 to 5	Mar. 28	May 19	May 23	May 27	June 24	Sept. 16	Sept. 21	
Do.....	F	1907	1907		S	2.3	2 to 5	Mar. 28	May 19	May 23	May 27	June 14	Sept. 16	Sept. 21	
Riparia × Rupestris, No. 3369.....	O	1904	1907		S	1.5	3 to 5	Mar. 20	May 16	June 1	May 21	June 11	Sept. 21	Oct. 8	
Riparia × Rupestris de Jaeger.....	F	1907	1907		S	3.2	3 to 5	Mar. 15	May 12	May 22	May 29	June 11	Sept. 21	Oct. 15	
Rupestris des Hautes-Alpes.....	F	1907	1907		C, S	3	3 to 5	Mar. 28	May 14	May 22	May 29	June 11	Sept. 21	Oct. 13	
Rupestris des Semis, No. 81-2.....	F	1907	1907		S	3.5	3 to 5	Mar. 26	May 14	May 22	May 29	June 13	Sept. 21	Oct. 10	
Rupestris Martin.....	F	1907	1907		S	3	3 to 5	Mar. 26	May 14	May 22	May 29	June 13	Sept. 21	Oct. 10	
Do.....	O	1903	1905		S	3.08	3 to 6	Mar. 15	May 12	May 21	May 28	June 11	Sept. 21	Oct. 11	
Rupestris St. George.....	F	1907	1907		S	3.7	3 to 6	Mar. 18	May 12	May 23	May 29	June 19	Sept. 21	Oct. 15	
Rupestris × Rupestris de Berlandieri, No. 219 A.....	F	1907	1907		C, S	3.2	3 to 5	Mar. 11	May 23	June 3	May 29	June 19	Sept. 21	Oct. 13	
Suff. Creek.....	O	1903	1905		S	3.1	3 to 5	Mar. 18	May 17	May 21	May 28	June 20	Sept. 21	Oct. 15	
Do.....	O	1903	1905		S	2.1	3 to 5	Mar. 18	May 17	May 21	May 28	June 20	Sept. 21	Oct. 15	
Solonis × Othello.....	F	1907	1907		S	1.9	3 to 7	Mar. 25	May 14	May 21	May 28	June 20	Sept. 21	Oct. 15	
Solonis × Rupestris, No. 1616.....	F	1907	1907		S	2.8	3 to 7	Mar. 30	May 21	June 2	May 28	June 20	Sept. 21	Oct. 15	
Do Taylor Narbonne.....	O	1904	1906		S	3.3	3 to 6	Mar. 20	May 21	June 5	May 26	June 20	Sept. 21	Oct. 15	
Do.....	O	1904	1906		S	4	3 to 6	Mar. 20	May 21	June 5	May 26	June 20	Sept. 21	Oct. 15	
Valencia.....	F	1907	1907		S	2.6	3 to 6	Mar. 16	May 22	May 29	May 23	June 23	Sept. 21	Oct. 5	
Vermorel.....	F	1907	1907		C, S	2.6	3 to 5	Mar. 10	May 15	May 23	May 29	June 23	Sept. 21	Oct. 5	
Viala.....	F	1907	1907		C, S	5	3 to 5	Mar. 20	May 14	May 25	May 24	June 24	Sept. 16	Oct. 17	

Ascert.	Own roots.	Gi	1904	1907	1906	1905	5	Mar. 24	Apr. 1	May 23	May 30	June 6	Sept. 20
Aspiran Noir.	Own roots.	Gi	1904	1907	1906	1905	S	Mar. 12	Mar. 30	May 24	May 29	June 6	Sept. 20
Augliato:	Adobe Giant.	F	1907	1907	1906	1905	C, S	3.3	3 to 5	Mar. 29	Mar. 17	May 23	May 4	June 15	Sept. 30
	Lenoir.	O	1904	1907	1906	1905	S	3.3	3 to 5	Mar. 17	Mar. 17	May 26	June 4	June 15	Sept. 30
Baba:	Own roots.	Gi	1904	1906	1906	1905	S	3.3	2 to 3	Mar. 18	Mar. 28	May 24	May 30	June 5	Sept. 24
	Dog Ridge.	O	1904	1906	1906	1905	S	3.3	2 to 4	Mar. 24	Mar. 24	May 25	June 7	June 21	Sept. 26
	Lenoir.	O	1903	1906	1906	1905	S	3.3	2 to 4	Mar. 21	Mar. 21	May 23	June 5	June 20	Sept. 25
	Rupestris St. George.	O	1903	1906	1906	1905	S	3.5	3 to 5	Mar. 21	Mar. 21	May 23	June 5	June 20	Sept. 25
	Taylor Narbonne	O	1906	1906	1906	1905	S	3	2 to 5	Mar. 21	Mar. 21	May 23	June 8	June 20	Sept. 26
Bakalar:	Own roots.	Gi	1904	1906	1906	1905	S	75	Mar. 14	Mar. 28	May 25	June 4	June 10	Sept. 25
	Aramon X Rupestris Ganzin, No. 1.	O	1904	1906	1906	1905	S	3.5	2 to 4	Mar. 20	Mar. 20	May 26	June 2	June 2	Sept. 22
	Dog Ridge.	O	1904	1906	1906	1905	S	1.4	June 18	Sept. 23
	Lenoir.	O	1903	1908	1908	1905	S	2.5	3 to 4	Mar. 19	Mar. 19	May 22	May 28	June 15	Sept. 21
	Riparia X Rupestris, No. 101.	O	1904	1904	1904	1905	C, S	4.4	3 to 4	Mar. 20	Mar. 20	May 24	June 3	June 16	Sept. 23
	Riparia X Rupestris, No. 3369.	O	1904	1906	1906	1905	S	3.5	3 to 4	Mar. 18	Mar. 18	May 26	June 2	June 14	Sept. 20
	Rupestris St. George.	F	1903	1905	1905	1905	S	3.2	2 to 5	Mar. 13	Mar. 30	May 17	May 22	June 24	Sept. 19
	Solenis X Riparia, No. 1616.	O	1904	1906	1906	1905	S	3.4	3 to 5	Mar. 18	Mar. 18	May 23	June 10	June 28	Sept. 22
	Taylor Narbonne.	O	1904	1906	1906	1905	S	4.3	3 to 4	Mar. 21	Mar. 21	May 26	June 5	June 19	Sept. 22
Barbarosa:	Own roots.	Gi	1905	1906	1906	1905	C, S	1.9	Mar. 14	Mar. 14	May 25	May 27	June 3	Sept. 24
	Adobe Giant.	F	1904	1906	1906	1905	C, S	4.1	3 to 5	Mar. 9	Mar. 9	May 13	May 16	May 22	Aug. 25
	Aramon X Rupestris Ganzin, No. 1.	F	1904	1906	1906	1905	C, S	2.7	2 to 6	Mar. 14	Mar. 30	May 13	May 20	May 26	Sept. 20
	Dog Ridge.	F	1904	1906	1906	1905	C, S	5.3	2 to 4	Mar. 12	Mar. 12	May 15	May 25do	Do.
	Do	F	1904	1906	1906	1905	C, S	1dodo
	Lenoir.	O	1903	1905	1905	1905	S	2.8	3 to 5	Mar. 18	Mar. 28	May 15	May 30	June 21	Sept. 27
	Do	F	1903	1905	1905	1905	S	2.7	3 to 5	Mar. 19	Mar. 19	May 15	May 25	June 21	Sept. 27
	Riparia Gloire.	F	1903	1905	1905	1905	S	2.2	3 to 6	Mar. 11	Apr. 1	May 10	May 16	May 20	Sept. 25
	Riparia X Rupestris, No. 3369.	F	1904	1906	1906	1905	S	2.9	3 to 6	Mar. 8	Apr. 5	May 13	May 18	May 22	Sept. 20
	Rupestris St. George.	F	1903	1905	1905	1905	S	4.9	3 to 6	Mar. 13	Mar. 30	May 14	May 21	May 29	Sept. 19
	Do	O	1903	1905	1905	1905	C, S	1.9	3 to 6	Mar. 19	Mar. 28	May 13	May 18	May 23	Sept. 20
	Solenis X Riparia, No. 1616.	F	1904	1906	1906	1905	S	2	2 to 5	Mar. 20	Mar. 28	May 13	May 18	May 25	Sept. 2
	Taylor Narbonne.	O	1904	1906	1906	1905	S
Barbarosa Finebourgo:	Own roots.	Gi	1901	1906	1906	1905	C, S	Mar. 13	Apr. 3	May 27	May 30	June 5	Sept. 23
Barbara:	Own roots.	Gi	1904	1907	1906	1905	S	2.3	2 to 4	Mar. 18	Mar. 30	May 22	June 3dodo
	Dog Ridge.	O	1904	1907	1906	1905	S	2.6	3 to 4	Mar. 18	Mar. 18	May 19	June 5	June 15	Sept. 24
	Lenoir.	O	1903	1905	1905	1905	S	1.9	3 to 5	Mar. 19	Mar. 19	May 26	June 2	June 14	Sept. 20
	Rupestris St. George.	O	1903	1905	1905	1905	Sdodododo
	Taylor Narbonne.	O	1904	1906	1906	1905	Sdo
Barducci:	Adobe Giant.	F	1907	1907	1906	1905	S	1.5	Mar. 12	Mar. 25	May 12	May 21	June 23	Sept. 15
	Lenoir.	O	1904	1907	1906	1905	S	2	2 to 5	Mar. 22	Mar. 22	May 21	June 5	June 17	Sept. 26

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	Low pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Bastardo:				<i>P. ct.</i>		<i>Lbs.</i>		9	10	11	12	13	14	15	16
Own roots.....	F	1904	1906	86	S	4.5	3 to 4	Mar. 15	Mar. 20	May 26	June 3	May 31	June 10	Aug. 25	Sept. 20
Dog Ridge.....	O	1904	1906	92	S	5.3	2 to 3	Mar. 17	Mar. 25	May 27	June 5	June 3	June 20	Sept. 23	Oct. 2
Lenoir.....	O	1903	1905	90	S	4.5	2 to 3	Mar. 15	Mar. 19	May 21	May 21	May 24	June 19	Sept. 20	Sept. 24
Rupestris St. George.....	O	1903	1905	93	S	4.5	2 to 3	Mar. 23	Mar. 23	May 22	June 1	May 28	June 15	Sept. 25	Oct. 10
Taylor Narbonne.....	O	1904	1906	92	S	4.8	2 to 3	Mar. 19	Mar. 19	May 28do.....	June 4do.....do.....	Oct. 8
Beclan:															
Own roots.....	F	1904	1906	90	S	3.6	2 to 4	Mar. 24	Mar. 30	May 24	May 30	May 29	June 7	Sept. 16	Sept. 25
Dog Ridge.....	O	1904	1906	90	S	2.8	2 to 4	Mar. 19	Mar. 19	May 20	June 8	May 24	June 20	Sept. 28	Oct. 15
Lenoir.....	O	1903	1905	88	S	2.6	2 to 5	Mar. 16	Mar. 16	May 22do.....	May 27	June 22do.....	Do.
Rupestris St. George.....	O	1903	1905	82	S	2.6	2 to 5	Mar. 16	Mar. 16	May 22do.....do.....do.....do.....	Oct. 10
Taylor Narbonne.....	O	1904	1906	82	S	1.3	2 to 5	Mar. 18	Mar. 18	May 23	June 1	May 28	June 11	Sept. 26	Do.
Bellino:															
Own roots.....	F	1904	1906	88	S	1.2	3 to 5	Mar. 17	Apr. 1	May 27	May 29	May 20	May 31	Aug. 25	Sept. 21
Adobe Giant.....	F	1904	1906	91	S	4.5	3 to 6	Mar. 13	Mar. 28	May 15do.....	May 29	June 7do.....	Sept. 20
Aramon X Rupestris Ganzin, No. 1.....	F	1904	1906	90	S	5.1	3 to 5do.....	Apr. 2	May 15do.....	May 25	June 7do.....	Do.
Dog Ridge.....	F	1904	1906	87	S	3.3	3 to 4	Mar. 14	Mar. 24	May 21	June 8	May 29	June 13	Sept. 26	Oct. 8
Do.....	F	1903	1905	88	S	3.3	3 to 6	Mar. 7	Mar. 30	May 12	June 16	May 21	June 24	Aug. 25	Sept. 13
Lenoir.....	F	1903	1905	92	S	4.2	3 to 6	Mar. 18	Mar. 18	May 26	June 1	June 2	June 20	Sept. 25	Do.
Do.....	F	1904	1906	88	S	4.2	3 to 6	Mar. 6	Apr. 1	May 20	May 26	May 21	June 1	Aug. 25	Sept. 20
Mourvedre X Rupestris, No. 1202.....	F	1904	1906	88	S	3.1	3 to 5do.....do.....	May 15	May 22	May 28	May 28	Oct. 17	Sept. 10
Pomroy.....	F	1907	1907	93	S	3.1	3 to 6	Mar. 15	Mar. 15	May 12	May 20	May 24	May 27do.....	Sept. 20
Riparia X Rupestris, No. 101.....	F	1904	1906	92	S	2	3 to 6	Mar. 7	Apr. 1	May 8do.....	May 24	May 27do.....	Sept. 19
Riparia X Rupestris, No. 3409.....	F	1904	1906	92	S	2	3 to 5	Mar. 20	Mar. 28	May 10	May 20	May 22	June 13	Aug. 25	Sept. 30
Rupestris des Causseilles.....	F	1907	1907	83	S	3.4	2 to 5	Mar. 14	Mar. 14	May 12	May 22	May 28	June 26	Aug. 25	Sept. 20
Rupestris St. George.....	F	1903	1905	89	S	1.8	2 to 3	Mar. 7	Mar. 25	May 16	May 25	May 25	May 30do.....	Do.
Do.....	F	1903	1905	83	S	3.7	3 to 5	Mar. 18do.....	May 13	May 25	May 22	May 27do.....	Do.
Solonis X Riparia, No. 1616.....	F	1904	1906	87	S	3.7	2 to 3	Mar. 18do.....	May 13	June 1	May 26	June 14do.....	Oct. 8
Taylor Narbonne.....	F	1904	1906	90	S	3	2 to 3	Mar. 22	Mar. 22	May 21	June 1	May 26	June 14do.....	Do.
Do.....	F	1904	1906	90	S	3.5	2 to 3	Mar. 22	Mar. 20	May 23	May 23	May 26	June 14do.....	Oct.
Vernorel.....	F	1907	1907	90	S	3.5	2 to 3	Mar. 20	Mar. 20	May 23	May 23	May 26	June 14do.....	Oct.

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Blanc d'Ambre:															
Lenoir.....	O	1904	1907	<i>P. cl.</i> 86	s	2.7	3 to 5	Mar. 17	May 18	May 31	June 14	Sept. 25	Sept. 20	
Monticola X Rupestris.....	F	1907	1907	86	s	1.8	3 to 5	Mar. 15	Mar. 25	May 14	May 22	June 14	Sept. 25	Sept. 23	
Mourvedre X Rupestris, No. 1202.....	F	1907	1907	83	s	3.5	Mar. 14	May 13	Sept. 15	Oct. 2	
Sali Creek.....	F	1907	1907	92	s	5	4 to 5	Mar. 24	Apr. 1	May 12	May 22	Oct. 4	
Blauer Portugieser:															
Own roots.....	Gi	1904	1904	s	Mar. 17	Mar. 29	May 24	May 28	June 7	Aug. 27	Aug. 28	
Dog Ridge.....	O	1906	1906	92	s	5.2	3 to 4	Mar. 18	May 25	June 1	June 13	Sept. 23	Sept. 23	
Lenoir.....	O	1903	1905	92	s	4	3 to 5	May 19	May 31	June 4	Sept. 10	Oct. 2	
Rupestris St. George.....	O	1903	1905	89	s	2.7	2 to 4	Mar. 19	May 25	June 5	June 15	Sept. 15	Oct. 4	
Boal de Madere:															
Own roots.....	Gi	1904	1907	89	s	3	May 24	June 1	June 7	Aug. 27	Sept. 23	
Dog Ridge.....	O	1903	1907	89	s	2 to 4	2 to 4	Mar. 22	Oct. 6	
Lenoir.....	O	1903	1905	95	s	4.9	3 to 4	Mar. 19	May 26	June 2	June 16	Sept. 15	Sept. 24	
Rupestris St. George.....	O	1903	1905	95	c, s	5.1	2 to 4	May 21	May 30	June 13	Sept. 24	Oct. 6	
Taylor Narbonne.....	O	1904	1906	92	s	4.1	2 to 4	Mar. 20	May 25	June 1	Oct. 10	
Rocheh:															
Own roots.....	Gi	1904	1904	s	Mar. 14	Mar. 28	May 26	May 30	June 1	
Bolyuno:															
Own roots.....	Gi	1904	1906	88	s	3.9	2 to 6	Apr. 1	May 20	June 6	June 13	Sept. 18	Sept. 24	
Dog Ridge.....	O	1903	1905	91	s	3.4	3 to 5	Mar. 17	May 20	June 8	June 17	Sept. 20	Oct. 1	
Lenoir.....	O	1903	1905	90	s	4.5	2 to 4	Mar. 19	May 22	June 6	June 20	Sept. 24	Oct. 8	
Rupestris St. George.....	O	1903	1906	88	c, s	4.6	2 to 4	Mar. 20	May 23	June 5	Do.	
Taylor Narbonne.....	O	1904	1906	88	s	
Bomarda:															
Own roots.....	Gi	1904	1904	s	Mar. 11	Apr. 2	June 3	Sept. 5	Sept. 23	
Boudales:															
Own roots.....	Gi	1907	1905	83	s	3.8	Mar. 98	Apr. 1	May 27	June 6	June 17	Sept. 20	Oct. 15	
Lenoir.....	O	1903	1905	93	s	4.3	2 to 4	Mar. 22	May 25	June 5	Oct. 8	
Rupestris St. George.....	O	1903	1905	93	s	
Bowood Muscat:															
Own roots.....	Gi	1904	1904	s	Mar. 25	Apr. 5	June 5	June 6	June 14	Sept. 19	Sept. 24	

Region	Variety	Year	Planting	Harvest	Yield	Quality	Planting	Harvest	Yield	Quality	Planting	Harvest	Yield	Quality
Brustiano:	Adobe Giant	1907	Mar. 20	Mar. 30	May 15	May 23	May 29	June 4	Sept. 21	Oct. 17				
	Lenoir	1904	do.	do.	May 15	May 23	May 28	June 4	Sept. 21	Oct. 17				
	Monticola	1907	Mar. 18	Mar. 28	May 15	May 23	May 24	June 4	Sept. 21	Oct. 17				
	Mourvedre	1907	Mar. 11	do.	May 14	May 28	May 28	do.	do.	do.				
	Bucarelli:	1904	Mar. 18	Mar. 18	May 23	June 7	May 30	June 22	Sept. 25	Oct. 2				
	Lenoir	1907	Mar. 19	Mar. 28	May 20	May 26	June 2	June 2	do.	do.				
	Rupestris St. George	1905	Mar. 15	Apr. 3	May 23	June 5	May 27	June 14	Sept. 22	Oct. 25				
	Buckland:	1905	Mar. 22	Mar. 28	May 23	June 5	May 27	June 14	Sept. 22	Oct. 25				
	Owln roots:	1904	Mar. 15	Mar. 28	May 21	June 1	May 26	June 6	Aug. 29	Oct. 25				
	Burger:	1904	Mar. 15	Mar. 28	May 21	June 1	May 26	June 6	Aug. 29	Oct. 25				
O	(Aestivals X Monticola) X (Riparia X Rupestris No. 554-5)	1909	Mar. 20	Mar. 30	May 23	June 10	May 28	June 16	Sept. 30	Oct. 11				
	Aramon X Rupestris Ganzin, No. 1.	1904	Mar. 21	Mar. 21	June 1	June 5	June 6	June 17	Oct. 1	Oct. 15				
	Aramon X Rupestris Ganzin, No. 2.	1907	Mar. 24	Mar. 24	June 2	June 3	do.	June 20	Oct. 10	Oct. 15				
	Beflandieri X Riparia, No. 420A	1907	Mar. 19	Mar. 19	May 20	June 1	June 6	June 15	Oct. 10	Oct. 15				
	Constantia.	1903	Mar. 17	Mar. 17	May 22	June 5	May 27	June 15	Sept. 20	Oct. 15				
	Colombaud X Rupestris No. 2502.	1906	Mar. 17	Mar. 17	May 22	June 5	May 27	June 15	Sept. 20	Oct. 15				
	Dog Ridge.	1905	Mar. 19	Mar. 19	May 23	June 8	May 29	June 17	Sept. 27	Oct. 12				
	Herboment.	1903	Mar. 23	Mar. 23	May 24	June 5	do.	June 17	Sept. 27	Oct. 12				
	Leonor.	1905	Mar. 21	Mar. 21	May 28	May 31	June 5	June 15	Oct. 10	Oct. 15				
	Monticola X Riparia, No. 1880K.	1907	Mar. 20	Mar. 20	May 25	June 5	May 30	June 23	Sept. 29	Oct. 11				
	Mourvedre X Rupestris, No. 1202.	1904	do.	do.	do.	do.	do.	do.	do.	do.				
	Riparia X Rupestris, No. 101.	1904	Mar. 19	Mar. 19	May 21	June 6	May 26	June 19	Oct. 5	Oct. 20				
	Riparia X Rupestris, No. 101-14.	1907	Mar. 21	Mar. 21	June 1	June 5	June 4	June 19	Oct. 5	Oct. 20				
	Riparia X Rupestris, No. 3306.	1907	Mar. 20	Mar. 20	June 2	June 8	June 6	June 23	Oct. 10	Oct. 11				
	Riparia X Rupestris, No. 3309.	1904	Mar. 20	Mar. 20	May 23	do.	May 29	June 27	Sept. 27	Oct. 6				
	Rupestris Martin.	1905	Mar. 18	Mar. 18	May 22	do.	do.	June 20	Sept. 30	Oct. 10				
	Rupestris Metaltica.	1907	do.	do.	May 31	do.	do.	June 25	do.	do.				
	Rupestris Mission.	1907	Mar. 19	Mar. 19	June 2	June 3	June 6	June 25	do.	do.				
	Rupestris St. George.	1903	Mar. 16	Mar. 16	May 22	May 31	May 28	June 10	do.	do.				
	Salt Creek.	1905	Mar. 17	Mar. 17	May 23	June 3	May 28	June 20	do.	do.				
	Solonis Robusta.	1907	Mar. 21	Mar. 21	June 1	June 10	June 6	June 24	do.	do.				
	Solonis X Othello, No. 1613.	1907	Mar. 18	Mar. 18	June 3	June 5	do.	June 18	do.	do.				
	Solonis X Riparia, No. 1616.	1907	Mar. 19	Mar. 19	May 21	June 8	May 26	June 23	do.	do.				
	Taylor Narbonne.	1904	Mar. 17	Mar. 17	May 22	June 5	May 29	June 18	do.	do.				
	Cabernet Sauvignon:	1906	Mar. 17	Mar. 17	May 22	June 5	May 29	June 18	do.	do.				
Owln roots:	1904	Mar. 20	Apr. 3	May 27	June 5	May 29	June 22	Sept. 23	Oct. 5					
Aramon X Rupestris Ganzin, No. 1.	1906	do.	do.	May 24	May 23	do.	June 20	Sept. 23	Oct. 10					
Dog Ridge.	1905	Mar. 25	Mar. 25	May 23	do.	do.	June 20	Sept. 23	Oct. 10					
Herboment.	1903	Mar. 24	Mar. 24	do.	do.	do.	June 21	Sept. 15	Oct. 11					
Leonor.	1905	Mar. 26	Mar. 26	May 24	June 7	May 28	June 20	Sept. 15	Oct. 5					
Mourvedre X Rupestris, No. 1202.	1904	Mar. 20	Mar. 20	May 24	June 10	May 27	June 27	Sept. 25	Oct. 6					
Riparia X Rupestris, No. 101.	1904	Mar. 21	Mar. 21	May 22	June 8	May 27	June 24	do.	do.					
Riparia X Rupestris, No. 3309.	1906	Mar. 20	Mar. 20	May 26	do.	June 2	June 22	do.	do.					
Rupestris Martin.	1905	do.	do.	May 29	June 5	May 25	do.	do.	do.					
Rupestris St. George.	1903	Mar. 24	Mar. 24	May 20	June 2	May 28	June 10	Sept. 30	Oct. 11					
Salt Creek.	1905	Mar. 25	Mar. 25	May 20	June 4	May 28	June 17	do.	do.					
Solonis X Riparia, No. 1616.	1904	Mar. 20	Mar. 20	May 24	June 2	May 28	June 19	do.	do.					
Taylor Narbonne.	1906	Mar. 20	Mar. 20	May 24	June 2	May 28	June 19	do.	do.					
O	1904	Mar. 24	Mar. 24	May 26	June 15	do.	June 23	Sept. 22	Oct. 10					

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of prunings per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Californian:															
Own roots.															
Australis.	F	1907	1907	88	C, S	4.5	3 to 5	Mar. 26	Mar. 29	May 31	June 4	June 9	June 9	June 9	Sept. 15
Do.	F	1907	1907	86	S	3.2	4 to 6	Mar. 14	Apr. 1	May 18	May 20	May 31	May 31	May 31	Sept. 5
Lendzi.	F	1907	1907	89	S	3.4	3 to 5	Mar. 21	Apr. 1	May 23	May 25	May 27	June 3	June 3	Aug. 28
Monticola × Rupestris.	F	1907	1907	91	C, S	3.3	3 to 5	Mar. 25	Apr. 1	May 14	May 23	May 28	June 19	June 19	Aug. 24
Riparia Gloire.	F	1907	1907	89	C, S	3.3	2 to 6	Mar. 12	Apr. 3	May 20	May 20	May 24	June 2	June 2	Sept. 25
Rupestris des Caussesteven.	F	1907	1907	85	C, S	3.7	3 to 4	Mar. 24	Apr. 3	May 16	May 24	June 1	June 2	June 2	Aug. 23
Rupestris St. George.	F	1907	1907	94	C, S	4.2	3 to 6	Mar. 14	Mar. 31	May 12	May 16	May 21	May 27	May 27	Aug. 23
Calmette:															
Adobe Giant.	F	1907	1907	82	S	2 to 4	2 to 4	Mar. 17	Apr. 5	May 18	May 22	June 2	June 2	June 2	Sept. 19
Aranson × Rupestris Gaucin, No. 1.	F	1907	1907	85	S	1.7	2 to 5	Mar. 18	Apr. 6	May 17	June 8	June 5	June 5	June 5	Sept. 21
Do.	F	1907	1907	92	C, S	4.9	2 to 5	Mar. 20	Apr. 5	May 23	June 8	June 28	June 20	June 20	Sept. 15
Berlandieri × Riparia, No. 420B.	F	1907	1907	81	S	5.8	3 to 5	Mar. 24	Apr. 5	May 13	May 21	May 31	June 2	June 2	Oct. 11
Constantia.	F	1907	1907	96	C, S	2.6	2 to 5	Mar. 23	Apr. 2	June 1	May 23	May 30	June 2	June 2	Oct. 17
Herbement.	F	1907	1907	82	S	2.6	2 to 4	Mar. 20	Apr. 2	June 1	June 5	June 8	June 20	June 20	Oct. 17
Lenoir.	O	1907	1907	89	S	2.6	2 to 4	Mar. 19	Apr. 2	May 12	June 5	May 18	June 22	June 22	Oct. 8
Monticola × Riparia, No. 18604.	F	1907	1907	86	C, S	3	2 to 5	Mar. 20	Apr. 5	May 10	May 22	May 29	May 30	May 30	Oct. 17
Monticola × Riparia, No. 18815.	F	1907	1907	88	C, S	3	2 to 4	Mar. 19	Apr. 5	May 12	May 23	May 29	May 30	May 30	Oct. 17
Monticola × Rupestris.	F	1907	1907	85	C, S	1.2	2 to 4	Mar. 24	Apr. 5	May 15	May 25	May 30	June 3	June 3	Sept. 17
Pomroy.	F	1907	1907	85	C, S	1.2	2 to 4	Mar. 24	Apr. 5	May 15	May 25	May 30	June 3	June 3	Sept. 5
Ramsay.	F	1907	1907	90	C, S	2.2	3 to 5	Mar. 21	Apr. 5	May 16	May 22	May 29	June 5	June 5	Sept. 14
Riparia × Rupestris, No. 101.	F	1907	1907	90	C, S	2.2	3 to 5	Mar. 18	Apr. 5	May 16	May 22	June 4	June 5	June 5	Sept. 21
Do.	F	1907	1907	87	C, S	2.2	2 to 6	Mar. 20	Apr. 5	May 12	June 5	May 31	June 5	June 5	Sept. 21
Riparia × Rupestris, No. 3309.	F	1907	1907	87	C, S	3.1	2 to 5	Mar. 20	Apr. 5	May 21	June 5	May 26	June 30	June 30	Oct. 16
Do.	O	1907	1907	88	C, S	2.8	2 to 5	Mar. 25	Apr. 5	May 17	June 4	May 29	June 19	June 19	Sept. 16
Riparia × Rupestris de Jaeger.	F	1907	1907	80	C, S	1.4	2 to 5	Mar. 21	Apr. 6	May 16	May 22	June 3	June 4	June 4	Oct. 11
Rupestris × Berlandieri, No. 218A.	F	1907	1907	81	C, S	1.7	2 to 5	Mar. 21	Apr. 6	May 16	May 24	June 3	June 4	June 4	Sept. 14
Rupestris × Berlandieri, No. 301-37-152.	F	1907	1907	85	C, S	.7	2 to 5	Mar. 20	Apr. 6	May 14	May 21	May 30	June 2	June 2	Sept. 20
Salt Creek.	F	1907	1907	82	S	2.2	3 to 5	Mar. 25	Apr. 1	May 17	May 21	May 29	June 2	June 2	Sept. 21
Solonis Robusta.	F	1907	1907	83	S	2.5	3 to 5	Mar. 22	Apr. 3	May 20	June 20	June 1	June 1	June 1	Sept. 21
	F	1907	1907	76	S	1.5	3 to 5	Mar. 23	Apr. 3	May 23	June 20	June 1	June 1	June 1	Sept. 21

F	Solomis × Othello	1907	85	c, s	2, 3	2 to 5	Mar. 21	Apr. 5	May 15	May 23	May 29	June 3	Sept. 8	Sept. 21
F	Solomis × Riparia, No. 1616	1907	88	s	4	2 to 5	Mar. 27	Mar. 23	Mar. 23	June 4	May 28	June 20	Sept. 20	Oct. 17
O	Do	1907	56	s	1, 2	2 to 5	Mar. 25	Mar. 23	May 2	May 22	May 2	June 20	Sept. 28	Oct. 9
F	Taylor Narbonne	1907	88	s	1, 7	2 to 4	do	Apr. 1	do	do	May 29	do	Oct. 17	
F	Vernorel	1907	85	s	4	2 to 4	do	do	do	do	do	do	do	
Carrigane:														
Gi	Own roots	1904		s			Mar. 18	Mar. 27	May 27	June 3	June 30	June 10	Sept. 20	Sept. 26
O	Aranon × Rupestris Ganzin, No. 1	1906	92	s	3	2 to 5	Mar. 19	Mar. 27	May 25	June 5	June 1	June 21	Sept. 28	Oct. 20
O	Dog Ridge	1903	95	s	3, 3	2 to 5	Mar. 23	Mar. 25	May 25	June 7	June 7	June 23	do	Oct. 10
O	Herbemout	1903	92	s	3, 9	1 to 6	do	do	May 24	June 5	May 30	June 19	Sept. 24	Oct. 11
O	Lenoir	1903	93	s	4, 3	3 to 6	Mar. 26	Mar. 25	May 24	June 7	do	June 29	Sept. 28	Oct. 11
O	Mourvèdre × Rupestris, No. 1202	1904	96	s	2, 9	3 to 6	Feb. 19	do	do	June 13	do	June 29	do	Oct. 11
O	Riparia × Rupestris, No. 101	1904	92	s	2, 2	3 to 6	Mar. 17	do	do	June 3	May 29	June 24	Sept. 25	Oct. 10
O	Riparia × Rupestris, No. 3309	1904	92	s	2, 8	2 to 5	Mar. 17	do	do	June 3	May 29	June 24	Sept. 27	Oct. 6
O	Rupestris Martin	1903	95	c, s	4	3 to 6	Mar. 22	Mar. 22	do	June 5	do	June 21	do	Oct. 8
O	Rupestris St. George	1903	85	s	4, 2	2 to 6	do	do	May 23	June 8	May 28	June 21	do	Oct. 8
O	Salt Creek	1903	95	c, s	2	3 to 6	Mar. 23	Mar. 23	May 23	June 12	May 30	June 24	do	Oct. 11
O	Solomis × Riparia, No. 1616	1904	90	s	2, 1	3 to 6	Mar. 17	Mar. 17	May 23	June 10	May 28	June 24	do	Oct. 11
O	Castizar	1904	90	c, s	3	3 to 6	Mar. 20	do	May 26	June 5	June 1	June 20	Sept. 29	Oct. 10
O	Riparia × Rupestris, No. 3309	1910	90	s	2, 1		do	do	June 3	June 15	June 7	June 29	do	do
O	Catarrato a la Porta	1910		s			Mar. 25	Mar. 30	May 28	June 10	June 2	June 15	Sept. 15	Sept. 20
Chaouch:														
Gi	Own roots	1904		s			Mar. 13	Mar. 25	May 29	June 3	June 10	do	Aug. 25	Do.
Gi	Own roots	1904		s			Mar. 15	do	May 18	June 9	May 23	June 13	Sept. 24	do.
O	Chaouch Rose	1910	91	s	4, 5	2 to 3	do	do	do	do	do	do	do	do
O	Lenoir	1907	97	c, s	4, 3	3 to 5	Mar. 18	do	do	June 1	May 22	June 20	Sept. 29	do.
Charbono:														
Gi	Own roots	1907		s			do	Apr. 5	May 27	May 27	May 29	do	Sept. 25	do.
O	Dog Ridge	1904	92	c, s	5, 5	2 to 4	do	Apr. 5	May 24	May 24	May 29	do	Oct. 3	do.
O	Lenoir	1903	93	c, s	3, 9	2 to 5	do	do	May 21	May 21	May 30	do	Sept. 28	do.
O	Rupestris St. George	1903	91	c, s	4	2 to 5	do	do	May 25	May 25	May 30	do	Sept. 27	do.
O	Riparia × Rupestris, No. 101	1904	90	s	2, 5	3 to 5	Mar. 18	do	May 24	June 8	do	June 20	Sept. 28	Sept. 30
O	Riparia × Rupestris, No. 3309	1904	90	s	2, 7	2 to 5	Mar. 20	do	May 26	do	do	June 21	Sept. 27	Oct. 6
O	Solomis × Riparia, No. 1616	1904	84	s	2	2 to 5	do	do	May 23	June 5	May 27	June 20	Sept. 24	Do.
O	Taylor Narbonne	1904	87	s	2, 3	2 to 5	Mar. 17	do	May 24	do	May 29	do	Sept. 25	Oct. 11
O	Chasselas Bouches du Rhone	1906	87	s	2, 3	2 to 5	do	do	May 24	do	May 29	do	Sept. 26	Oct. 10
Chasselas Bouches du Rhone:														
Gi	Own roots	1905		c, s			Mar. 18	Apr. 1	May 28	June 1	May 29	June 15	Sept. 5	Sept. 25
O	Dog Ridge	1904	91	c, s	3, 5	3 to 5	Mar. 15	do	May 24	do	May 29	June 15	Sept. 9	Oct. 9
O	Lenoir	1903	90	c, s	3, 3	2 to 5	Mar. 13	do	May 20	do	May 24	June 15	do	Oct. 9
O	Rupestris St. George	1903	93	s	5, 2	3 to 4	Mar. 17	do	May 21	June 2	May 26	June 21	do	Oct. 10
O	Taylor Narbonne	1904	70	s	5	3 to 5	Mar. 18	do	May 20	June 5	May 25	June 10	do	Sept. 21
Chasselas Bulberry:														
O	Dog Ridge	1904	92	s	4	3 to 5	Mar. 15	do	May 23	June 1	May 29	June 13	Sept. 21	Oct. 10
O	Lenoir	1903	87	s	4, 2	2 to 5	do	do	May 20	June 8	May 23	June 21	Sept. 23	Sept. 25
O	Rupestris St. George	1903	94	s	4, 4	2 to 5	Mar. 14	do	May 24	June 1	May 30	June 13	Sept. 9	Sept. 30
O	Taylor Narbonne	1904	75	s	1, 6	3 to 5	Mar. 18	do	do	do	May 29	June 9	Sept. 23	Oct. 1

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congenality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Chasselas Cloufard:															
Own roots.....	Gi	1907													
Dog Ridge.....	O	1904	1906	91	S	3.3	2 to 5	Mar. 17	Apr. 10	May 28	June 2	May 26	June 15	Sept. 9	Oct. 10
Lenoir.....	O	1903	1905	87	S	2.9	2 to 5	do.	do.	do.	June 6	May 21	June 14	do.	Oct. 1
Rupestris St. George.....	O	1903	1905	88	S	3.9	2 to 5	do.	do.	do.	June 1	May 25	June 20	do.	Oct. 2
Chasselas Dore:															
Own roots.....	Gi	1905													
Dog Ridge.....	O	1904	1906	91	S	2.8	3 to 5	Mar. 20	Mar. 26	May 29	June 2	June 7	June 15	Aug. 24	Sept. 21
Lenoir.....	O	1903	1905	94	S	3.5	3 to 5	Mar. 14	do.	May 25	June 31	May 30	June 12	Sept. 9	Oct. 10
Rupestris St. George.....	O	1903	1905	94	S	6.1	2 to 5	Mar. 15	do.	May 23	June 3	May 24	June 20	do.	Oct. 2
Chasselas Duhamel:															
Aramon X Rupestris Ganzin, No. 1.....	O	1904	1906	92	S	5	2 to 4	Mar. 21	do.	May 31	June 1	June 2	June 13	Sept. 29	Oct. 8
Dog Ridge.....	O	1904	1906	91	S	4	3 to 4	Mar. 15	do.	May 21	June 3	May 26	June 19	Sept. 9	Oct. 10
Lenoir.....	O	1903	1905	90	S	3.1	3 to 5	do.	do.	May 20	do.	May 28	June 14	do.	Oct. 2
Rupestris St. George.....	O	1903	1905	94	S	4.1	2 to 5	Mar. 21	do.	May 21	June 8	May 26	June 24	do.	Oct. 2
Taylor Narbonne.....	O	1904	1906		S	2.5									
Chasselas de Fontainebleau:															
Rupestris St. George.....	F	1903	1905	85	S	1.5	2 to 6	Mar. 14	Mar. 30	May 15	May 17	May 20	May 24	Aug. 19	Sept. 19
Chasselas Florence:															
Own roots.....	Gi	1905													
Dog Ridge.....	O	1904	1906	91	S	4.7	2 to 5	Mar. 8	Apr. 21	May 27	June 1	June 7	June 12	Sept. 23	Oct. 10
Lenoir.....	O	1903	1905	91	S	3.1	3 to 5	Mar. 16	do.	May 24	do.	May 30	June 15	Sept. 9	Oct. 9
Rupestris St. George.....	O	1903	1905	91	S	3.9	2 to 4	Mar. 17	do.	May 25	June 1	May 27	June 18	do.	Do.
Taylor Narbonne.....	O	1904	1906	82	S	1	2 to 3	Mar. 16	do.	May 22	do.	May 27	June 12	Sept. 21	Do.
Chasselas Montauban:															
Own roots.....	Gi	1905													
Lenoir.....	O	1903	1905	95	S	2.5	2 to 6	Mar. 18	Apr. 1	do.	May 27	May 26	June 17	Sept. 9	Sept. 25
Rupestris St. George.....	O	1903	1907	91	S	4.7	2 to 5	Mar. 24	do.	May 23	June 3	May 28	June 20	do.	Oct. 12
Chasselas Musque Vert:															
Own roots.....	Gi	1907													
Dog Ridge.....	O	1904	1906	91	S	3.2	2 to 5	Mar. 15	Mar. 20	May 25	June 1	May 27	June 20	Sept. 25	Oct. 10
Lenoir.....	O	1903	1905	88	S	3.9	2 to 5	Mar. 20	do.	May 22	June 1	May 24	June 18	do.	Oct. 9
Rupestris St. George.....	O	1903	1905	94	S	6.2	1 to 4	Mar. 17	do.	May 23	do.	May 28	June 15	do.	Oct. 10

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California. (Continued.)

Variety and stock (on own roots, if so stated).	Experiment vineyard	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Chenin Blanc:															
Own roots.....	Gi	1905													
Dog Ridge.....	O	1904	1906	91	c, s			Mar. 15	Mar. 28	May 26	May 30	May 30	June 5	Aug. 28	Sept. 25
Lenoir.....	O	1903	1905	93	s	3.6	2 to 4	Mar. 20	Mar. 18	May 24	May 29	May 29	June 15	Sept. 23	Oct. 17
Rupestris St. George.....	O	1903	1906	90	s	5.8	2 to 5	Mar. 18	Mar. 19	May 21	May 29	May 29	June 20	Sept. 15	Sept. 28
Child of Hall:															
Lenoir.....	O	1904	1907	89	s	5.6	2 to 4	Mar. 18	Mar. 18	May 23	May 29	May 29	June 17	Sept. 20	Oct. 1
Chassac:															
Own roots.....	Gi	1904													
Lenoir.....	O	1903	1905	81	s	3.2	3 to 5	Mar. 21	Mar. 29	May 26	June 4	June 2	June 9	Aug. 28	Sept. 20
Rupestris St. George.....	O	1903	1905	95	s	2.2	2 to 3	Mar. 17	Mar. 17	May 24	June 1	June 1	June 10	Sept. 28	Oct. 4
Cipro Nero:															
Own roots.....	Gi	1904													
Charlot & Gros Grain:															
Lenoir.....	O	1904	1907	88	s	3.9	2 to 4	Mar. 20	Mar. 30	May 25	May 29	May 29	June 20	Sept. 30	Oct. 8
Monticola X Rupestris.....	F	1907	1907	82	c, s	2.3	3 to 6	Mar. 14	Mar. 30	May 11	May 22	May 30	June 20	Sept. 21	Oct. 21
Charette Blanche:															
Own roots.....	Gi	1904													
Charette Mazed:															
Own roots.....	Gi	1905													
Lenoir.....	O	1906	1912	85	c, s	4	3 to 4	Mar. 22	Mar. 30	May 11	May 22	May 30	June 9	Sept. 20	Oct. 7
Riparia Gloire.....	O	1904						Mar. 20	Apr. 1	May 27	June 4	June 4	June 9	Sept. 20	Oct. 28
Cornua Nostra:															
Own roots.....	Gi	1907	1907	88	s	4	4	Mar. 20	Apr. 1	May 29	June 2	June 2	June 7	Aug. 28	Sept. 20
Algerais.....	F	1907	1907	84	s	4.8	4 to 5	Mar. 20	Mar. 20	May 25	May 31	June 4	June 7	Sept. 5	Oct. 7
Lenoir.....	O	1904	1907	84	s	2.8	2 to 4	Mar. 19	Mar. 19	May 30	June 8	June 6	June 23	Sept. 22	Do.
Coltura Cusco Bianco:															
Own roots.....	F	1907	1907	90	c, s	5.7	2 to 5	Mar. 23	Mar. 23	May 27	May 27	May 30	June 13	Sept. 30	Oct. 3
Commandeur:															
Own roots.....	Gi	1904													
Riparia X Rupestris, No. 3306.....	O	1904		85	s	6.5	2 to 3	Mar. 20	Mar. 30	May 24	June 6	May 29	June 12	Aug. 29	Sept. 26
										May 12	June 1	May 17	June 15	Sept. 25	Oct. 1

Year	Number	Origin	Roots	Height	Flowering	Harvest	Quality	Notes					
1904	1904	Corbeau Noir Shar.	F	91	5	Mar. 18	Mar. 30	May 26	June 3	June 8	June 9	Sept. 17	Sept. 25
1906	1907	Own roots.	F	96	5	Mar. 17	Mar. 3	May 18	June 23	June 8	June 8	June 25	Sept. 23
1904	1907	Corinthe Blanc.	F	96	5	Mar. 17	Mar. 3	May 18	June 23	June 8	June 8	June 25	Sept. 23
1904	1907	Corinthe & Gros Grain.	F	96	5	Mar. 17	Mar. 3	May 18	June 23	June 8	June 8	June 25	Sept. 23
1904	1907	Adobis Giant.	F	98	9.6	Mar. 20	Apr. 1	May 12	May 24	May 24	May 24	Sept. 21	Sept. 15
1904	1907	Canada.	F	98	5	Mar. 20	Apr. 1	May 12	May 24	May 24	May 24	Sept. 21	Sept. 15
1904	1907	Concanta.	F	96	7.7	Mar. 14	Apr. 28	May 16	May 22	May 23	May 23	Sept. 10	Sept. 15
1904	1907	Lenoir.	F	99	3.1	Mar. 23	Apr. 3	May 25	June 4	May 29	May 29	Aug. 19	Aug. 13
1904	1907	Monticola X Rupestris.	F	95	3.3	Mar. 23	Apr. 3	May 25	June 4	May 29	May 29	Aug. 19	Aug. 13
1904	1907	Mourvedre X Rupestris, No. 1202.	F	96	4.1	Mar. 15	Apr. 1	May 13	May 22	May 25	May 25	Sept. 10	Sept. 15
1904	1907	Riparia Grand Glaibre.	F	97	9.7	Mar. 15	Apr. 1	May 13	May 22	May 25	May 25	Sept. 10	Sept. 15
1904	1907	Riparia X Rupestris, No. 101.	F	97	9	Mar. 23	Apr. 1	May 14	May 22	May 25	May 25	Sept. 10	Sept. 15
1904	1907	Vernorel.	F	93	7.1	Mar. 19	Mar. 28	May 18	May 25	May 25	May 25	Sept. 10	Sept. 15
1904	1907	Coristano.	F	97	7.5	Mar. 23	Apr. 1	May 16	May 24	June 1	June 1	Sept. 17	Oct. 17
1904	1907	Australis.	F	96	4.3	Mar. 25	Apr. 1	May 16	May 21	May 29	May 29	Sept. 17	Oct. 26
1904	1907	Barnes.	F	96	2.2	Mar. 20	Apr. 1	May 26	June 7	May 30	May 30	Sept. 17	Oct. 17
1904	1907	Lenoir.	F	96	6.3	Mar. 28	Apr. 1	May 19	May 25	June 4	June 4	Sept. 17	Oct. 17
1904	1907	Monticola X Rupestris.	F	96	5.5	Mar. 21	Apr. 1	May 17	May 22	May 30	May 30	Sept. 17	Oct. 17
1904	1907	Rupestris Martin.	F	96	7	Mar. 25	Apr. 1	May 20	May 23	June 3	June 3	Sept. 10	Do.
1904	1907	Rupestris St. George.	F	96	7	Mar. 25	Apr. 1	May 20	May 23	June 3	June 3	Sept. 10	Do.
1904	1907	Own roots.	G1	93	3.5	Mar. 17	Mar. 27	May 30	June 6	June 4	June 4	Sept. 19	Oct. 1
1904	1907	Aramon X Rupestris Ganzin, No. 1.	O	94	4.1	Mar. 24	Mar. 27	May 22	June 10	May 28	May 28	Sept. 27	Oct. 1
1904	1907	Lenoir.	O	90	6	Mar. 22	Apr. 5	May 23	June 8	June 20	June 20	Oct. 13	Oct. 13
1904	1907	Riparia Gloire.	O	90	6	Mar. 22	Apr. 5	May 23	June 8	June 20	June 20	Oct. 13	Oct. 13
1904	1907	Rupestris X Borlandfrei, No. 301A.	F	90	4.5	Mar. 12	Apr. 6	May 23	May 24	May 29	May 29	Sept. 15	Sept. 15
1904	1907	Solonis Robusta.	F	92	4.7	Mar. 12	Apr. 6	May 18	May 24	May 29	May 29	Sept. 15	Sept. 15
1904	1907	Crabbs Burgundy.	F	92	4.7	Mar. 12	Apr. 6	May 18	May 24	May 29	May 29	Sept. 15	Sept. 15
1904	1907	Own roots.	O	87	3.8	Mar. 18	Apr. 10	May 25	June 6	May 24	May 24	Sept. 23	Oct. 12
1904	1907	Aramon X Rupestris Ganzin, No. 1.	O	94	4	Mar. 20	Mar. 20	May 19	June 2	June 21	June 21	Sept. 28	Oct. 8
1904	1907	Dog Ridge.	O	95	3.9	Mar. 17	Mar. 15	May 26	June 5	June 2	June 2	Sept. 28	Oct. 8
1904	1907	Lenoir.	O	91	3.2	Mar. 15	Mar. 15	May 20	June 17	June 20	June 20	Sept. 28	Oct. 12
1904	1907	Riparia X Rupestris, No. 101.	O	94	2.7	Mar. 20	Mar. 20	May 21	June 4	May 27	May 27	Sept. 28	Oct. 11
1904	1907	Riparia X Rupestris, No. 3399.	O	94	2.8	Mar. 20	Mar. 20	May 21	June 4	May 27	May 27	Sept. 28	Do.
1904	1907	Rupestris St. George.	O	88	2.8	Mar. 16	Mar. 16	May 23	June 10	May 29	May 29	Sept. 29	Oct. 8
1904	1907	Solonis X Riparia, No. 1616.	O	94	1.1	Mar. 20	Mar. 20	May 19	June 10	May 24	May 24	Sept. 29	Oct. 11
1904	1907	Taylor Narbonne.	O	85	2.1	Mar. 20	Mar. 20	May 21	June 12	May 26	May 26	Sept. 28	Oct. 10
1904	1907	Own roots.	G1	94	2.1	Mar. 19	Apr. 1	May 24	June 3	May 29	May 29	Aug. 29	Sept. 27
1904	1907	Crudero.	G1	94	2.1	Mar. 17	Mar. 26	May 28	June 6	June 2	June 2	Sept. 22	Oct. 13
1904	1907	Own roots.	G1	94	2.1	Mar. 17	Mar. 26	May 28	June 6	June 2	June 2	Sept. 22	Oct. 13

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year planted was	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
I	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Danague:				<i>P. cl.</i>	<i>s</i>	<i>Lbs.</i>									
Own roots.....	Gi	1904		86	<i>s</i>	2	2 to 5	Mar. 24	Mar. 28	June 2	June 4	June 7	Sept. 24
Dog Ridge.....	O	1904		89	<i>s</i>	3.1	3 to 3	Mar. 17	May 23	May 31	May 30	June 12	Sept. 9	Sept. 25
Lenoir.....	O	1903		82	<i>c, s</i>	3.0	3 to 4	Mar. 17	May 20	June 6	May 29	June 14	Sept. 11	Do.
Rupestris St. George.....	O	1903		81	<i>s</i>	1.8	2 to 4	Mar. 20	May 23	June 2	May 29	June 7	Do.
Taylor Narbonne.....	O	1904		81	<i>s</i>	1.8	2 to 4	May 19	June 9	May 21	June 15
Dattler de Beyrouth.....	O	1903	1912	58	<i>s</i>	2.5	Mar. 16	June 4	June 9
Deacon Steyer.....	Gi	1907	<i>c, s</i>	Mar. 10	Apr. 1	May 26	Sept. 19
Own roots.....	Gi	1905	<i>s</i>	Mar. 23	Sept. 25
Dordrelat:					<i>s</i>										
Own roots.....	Gi	1904	1909	<i>s</i>	3 to 4	3 to 4	Mar. 18	Mar. 28	May 2	June 1	May 31	June 7	Sept. 20	Sept. 26
Amarin X Rupestris Ganzin, No. 1.....	O	1904	1907	94	<i>s</i>	3.5	3 to 4	Mar. 16	May 2	June 8	June 8	June 22	Oct. 10	Oct. 20
Dog Ridge.....	O	1901	1909	89	<i>s</i>	3.1	3 to 4	Mar. 20	May 25	June 5	May 26	June 20	Oct. 15	Oct. 15
Flacumont.....	O	1903	1905	90	<i>s</i>	2.7	2 to 4	Mar. 17	June 3	June 8	June 9	June 25	Oct. 2	Oct. 10
Lenoir.....	O	1903	1905	89	<i>s</i>	3.0	3 to 4	Mar. 17	June 3	June 8	June 9	June 25	Oct. 10	Oct. 15
Mourvèdre X Rupestris, No. 1202.....	O	1904	1909	95	<i>s</i>	3.3	3 to 4	Mar. 21	June 3	June 6	June 8	June 20	Oct. 10	Oct. 12
Rupestris St. George.....	O	1903	1905	89	<i>s</i>	3.3	3 to 4	Mar. 19	June 22	June 2	May 28	June 21	Sept. 28	Oct. 10
Solonis X Riparia, No. 1616.....	O	1904	1909	93	<i>s</i>	4.2	3 to 4	Mar. 23	May 25	June 1	June 8	June 15	Oct. 12	Oct. 12
Taylor Narbonne.....	O	1904	1906	87	<i>s</i>	2.2	3 to 4	Mar. 18	May 27	June 1	June 2	June 13	Sept. 30	Oct. 4
Draculus:					<i>s</i>										
Own roots.....	Gi	1904	<i>s</i>	Mar. 14	Mar. 24	May 28	June 5	June 10	Aug. 29	Sept. 25
Dronkane:					<i>s</i>										
Lenoir.....	O	1904	1907	88	<i>s</i>	3	3 to 5	Mar. 26	May 26	June 15	June 1	June 30	Sept. 25	Oct. 25
Rupestris des Causseottes.....	F	1907	88	<i>s</i>	2	4 to 5	Mar. 21	Apr. 1	May 16	May 22	May 27	Oct. 17
Duc de Magenta.....	F	1903	1905	86	<i>s</i>	2.3	5 to 7	Mar. 9	Mar. 15	May 10	May 16	May 20	May 26	Aug. 5	Sept. 13
Rupestris St. George.....	F	1903	1905	86	<i>s</i>	2.3	5 to 7	Mar. 9	Mar. 15	May 10	May 16	May 20	May 26	Aug. 5	Sept. 13
Duc de Malakoff:					<i>c, s</i>										
Own roots.....	Gi	1905	<i>c, s</i>	Mar. 18	Mar. 27	May 29	May 31	June 6	Aug. 28	Sept. 25

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California.—(continued.)

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year planted was	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-settling date.		Fruit-ripening date.			
								Early sea- son.	Late sea- son.	Early sea- son.	Late sea- son.	Early sea- son.	Late sea- son.	Early sea- son.	Late sea- son.		
Flame Tokay—Continued. Monticola × Riparia, No. 534..... Monticola × Riparia, No. 18804..... Monticola × Riparia, No. 18808..... Monticola × Riparia, No. 18815..... Monticola × Riparia, No. 18845..... Riparia 3 Grandes Feuilles..... Riparia × Rupestris, No. 101..... Riparia × Rupestris, No. 3309..... Do..... Rupestris Martin..... Salt Creek..... Taylor Narbonne..... Folle Blanche: Rupestris St. George..... Foster:	F	1907	1907	96	c, s	6	3 to 5	Mar. 22	Apr. 1	May 17	May 22	May 30	Sept. 29	Sept. 29	Late sea- son.		
	F	1907	1907	95	c, s	7	3 to 5	Mar. 23	do.	May 18	May 26	May 28	do.	do.	do.	Early sea- son.	
	F	1907	1907	97	s	7	3 to 6	Mar. 24	Apr. 6	May 20	May 25	May 30	do.	do.	do.	do.	
	F	1907	1907	97	s	6.9	3 to 6	do.	do.	do.	do.	May 22	do.	do.	do.	do.	
	F	1907	1907	90	s	4.9	3 to 5	Mar. 23	Apr. 2	May 14	May 23	May 24	May 24	Sept. 29	Sept. 29	do.	
	F	1907	1907	94	s	3.1	2 to 4	Mar. 21	Mar. 21	May 25	June 3	June 1	June 1	June 14	Sept. 29	do.	
	O	1904	1907	94	s	4.2	3 to 5	Mar. 22	Apr. 2	May 21	May 24	May 30	June 1	June 1	Sept. 29	do.	
	O	1904	1907	94	s	4.1	3 to 5	Mar. 25	Apr. 2	May 19	May 24	May 30	June 1	June 1	Sept. 29	do.	
	F	1907	1907	83	c, s	2.7	3 to 4	Mar. 23	Apr. 1	May 23	May 24	May 24	May 24	May 24	Sept. 25	Sept. 25	
	F	1907	1907	91	c, s	4.2	3 to 4	Apr. 1	Apr. 2	May 14	May 24	May 24	May 24	May 24	Sept. 12	Sept. 12	
	F	1907	1907	95	c, s	4.4	3 to 4	Mar. 23	Apr. 2	May 14	May 24	May 24	May 24	May 24	Sept. 12	Sept. 12	
	F	1903	1905	79	s	.9	3 to 6	Mar. 8	Mar. 30	May 15	May 20	May 20	June 5	June 5	Sept. 13	Sept. 13	
	Own roots..... Lenoir..... Riparia Gloire..... Rupestris St. George..... Frankenthal Precocoe:	G	1905	1905	88	s	1.4	3 to 6	Mar. 15	Mar. 28	May 24	May 27	May 30	June 5	June 5	Sept. 5	Sept. 20
		F	1903	1905	84	s	1.8	3 to 5	Mar. 11	Mar. 25	May 13	May 18	May 22	May 28	May 28	Aug. 24	Sept. 19
		F	1903	1905	90	s	3	2 to 6	Mar. 15	do.	May 12	May 19	May 23	May 25	May 25	Aug. 23	Sept. 19
F		1903	1905	90	s	3	2 to 6	Mar. 15	do.	May 15	May 20	May 21	June 4	June 4	Aug. 24	Sept. 10	
Herbemont..... Lenoir..... Monticola × Riparia, No. 18808..... Monticola × Riparia, No. 18815..... Rupestris St. George..... Solonis × Othello..... Solonis × Riparia, No. 1616..... Frederickton:	O	1904	1909	72	s	1.1	3 to 5	Mar. 24	Mar. 24	June 2	June 7	June 7	June 7	June 20	Sept. 23	Oct. 5	
	O	1903	1905	90	c, s	2.2	3 to 5	Mar. 18	Mar. 18	May 20	June 1	May 29	June 14	June 14	Sept. 23	Oct. 17	
	F	1907	1907	90	c, s	3.3	4 to 6	Mar. 20	Mar. 30	May 14	May 24	May 28	do.	do.	Aug. 30	Do.	
	F	1907	1907	87	c, s	2.7	4 to 6	do.	do.	May 16	May 23	do.	do.	do.	Aug. 30	Do.	
	O	1903	1905	92	c, s	2.2	3 to 4	do.	do.	do.	do.	do.	do.	do.	do.	Sept. 30	
	F	1907	1907	90	c, s	3	3 to 4	Mar. 24	Mar. 24	May 22	May 22	May 28	do.	do.	Oct. 17	Sept. 30	
	F	1907	1907	89	s	3.7	3 to 5	Mar. 25	Apr. 2	May 15	May 22	May 28	do.	do.	Sept. 1	Sept. 21	
	F	1907	1907	92	c, s	5	3 to 4	do.	do.	May 18	May 23	May 23	May 19	June 17	Sept. 25	Sept. 28	
O	1904	1907	91	c, s	2.8	3 to 4	Mar. 17	Apr. 1	May 25	June 5	May 30	June 4	June 17	Sept. 25	Sept. 28		
F	1907	1907	92	s	5	3 to 4	Mar. 31	Apr. 1	May 21	May 23	June 3	June 3	June 17	Sept. 25	Sept. 28		

Year	Code	Origin	S	Quality	Mar. 16	Mar. 29	May 26	May 30	June 6	Sept. 23
1904	GI	Fresa de Monferat: Own roots.....	s							
1907	GI	Gamay de Bourgogne: Own roots.....	s		Mar. 8	Apr. 8	May 20	May 28	June 15	Sept. 15
1904	O	Dog Ridge.....	s	3.6	Mar. 20	May 19	May 19	May 31	June 11	Sept. 9
1903	O	Lenoir.....	s	2.4	Mar. 16	do.	do.	May 30	May 23	Sept. 8
1903	O	Rupestris St. George.....	s	4.3	Mar. 18	do.	do.	June 1	May 24	Sept. 24
1904	GI	Gamay Teinturier: Own roots.....	s		do.	Apr. 3	May 27	June 3	June 6	Sept. 20
1904	O	Dog Ridge.....	c, s	3.6	Mar. 16	May 19	May 19	May 31	June 9	Sept. 28
1903	O	Lenoir.....	s	2.2	Mar. 15	May 24	May 24	do.	June 1	Oct. 5
1903	O	Rupestris St. George.....	s	3.3	Mar. 19	May 25	May 25	June 3	June 10	Oct. 4
1904	O	Taylor Narbonne.....	s	.8	do.	May 19	May 19	May 31	June 8	Oct. 1
1904	GI	Gewirtz Traminer: Own roots.....	c, s		Mar. 12	Apr. 1	May 29	May 30	June 6	Sept. 22
1905	GI	Golden Champion: Own roots.....	c, s		Mar. 27	Mar. 30	May 28	June 3	June 2	Sept. 19
1904	GI	Do.....	s		Mar. 12	Apr. 13	May 10	May 24	May 29	Sept. 6
1904	O	Dog Ridge.....	s	4	Mar. 25	May 24	May 24	June 5	June 20	Sept. 28
1903	O	Lenoir.....	s	3.5	Mar. 20	do.	do.	do.	do.	Oct. 25
1903	O	Rupestris St. George.....	s	3.7	Mar. 17	do.	do.	do.	do.	Oct. 9
1906	O	Golden Hamburg: Lenoir.....	s	.5	Mar. 16	do.	June 5	June 8	June 9	Sept. 30
1906	O	Golden Queen: Lenoir.....	c, s	.7	Mar. 19	do.	May 21	do.	June 6	Oct. 8
1904	O	Goebacie: Own roots.....	c, s	4.5	Mar. 18	do.	May 12	May 31	May 18	Sept. 22
1907	GI	Grarduska: Own roots.....	s		Mar. 12	Apr. 10	May 26	June 1	June 29	Sept. 25
1904	O	Dog Ridge.....	s	2.7	Mar. 22	May 23	May 23	June 8	June 17	Sept. 23
1903	O	Lenoir.....	s	2.4	Mar. 20	May 19	May 19	June 1	June 23	Oct. 8
1903	O	St. George.....	s	2.9	Mar. 24	May 21	May 21	June 3	June 16	Oct. 10
1906	O	Taylor Narbonne.....	s	1.4	Mar. 20	May 21	May 21	June 8	June 26	Oct. 4
1904	GI	Green Hungarian: Own roots.....	s		Mar. 12	Apr. 1	May 27	May 30	June 5	Sept. 25
1907	O	(Aestivallis X Monticola) X (Riparia X Rupestris, No. 554-5).	s	2	Mar. 22	June 7	June 7	June 8	June 12	Sept. 19
1906	O	Aramon X Rupestris Ganzin, No. 1.....	s	3	Mar. 21	May 22	May 22	June 8	May 28	Oct. 20
1907	O	Berlandieri X Riparia, No. 2.....	s	.5	Mar. 19	do.	do.	do.	June 24	Oct. 14
1909	O	Constantia.....	s	.9	Mar. 20	do.	do.	June 5	June 6	do.
1903	O	Dog Ridge.....	s	3.8	Mar. 19	May 28	May 28	June 3	June 4	do.
1903	O	Herberton.....	s	1.3	Mar. 18	May 25	May 25	June 5	June 21	Sept. 30
1903	O	Lenoir.....	c, s	2.1	do.	May 25	May 25	June 7	June 21	Oct. 17
1907	O	Monticola X Riparia, No. 18808.....	s	1.6	Mar. 19	June 2	June 2	June 5	June 20	Oct. 10
1906	O	Monvredre X Rupestris No. 1202.....	s	1.5	Mar. 22	May 20	May 20	June 5	May 29	Sept. 30
1904	O	Riparia X Rupestris No. 101.....	s	3	Mar. 21	June 1	June 1	do.	June 26	Oct. 15
1907	O	Riparia X Rupestris, No. 3306.....	s	1.9	Mar. 22	June 1	June 1	do.	June 8	Oct. 10
1904	O	Riparia X Rupestris, No. 3369.....	s	3	Mar. 21	June 1	June 1	do.	June 26	Oct. 15
1907	O	Rupestris Martin.....	s	.2	Mar. 21	June 1	June 1	June 5	June 5	Oct. 6

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California.—Continued.

Variety and stock (on own roots, if so stated).	Experiment yard.	Year planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Modes bearing.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Green Hungarian—Continued.															
Rupprecht's Mission.....	O	1907	1909	67	S	1.4	2 to 5	Mar. 18	June 2	June 5	June 2	June 12	Sept. 29	Oct. 10	16
Rupprecht's St. George.....	O	1903	1905	79	S	1.4	2 to 5	do	May 20	June 1	June 2	June 12	Oct. 1	Oct. 11	
Salt Creek.....	O	1903	1905	90	S	2	3 to 5	Mar. 21	May 21	June 1	June 6	June 20	Oct. 5	Oct. 15	
Solanis Rubuska.....	O	1907	1909	82	S	1.9	2 to 3	Mar. 20	June 1	June 1	June 10	June 22	Sept. 30	Oct. 15	
Solanis X Othello, No. 1013.....	O	1907	1909	87	S	1.9	2 to 5	Mar. 19	June 1	June 1	June 10	June 22	Sept. 30	Oct. 15	
Solanis X Riparian, No. 1616.....	O	1901	1903	68	S	1.9	3 to 5	Mar. 22	May 23	June 1	June 1	June 12	Sept. 29	Do.	
Solator Narbonne.....	O	1901	1901	81	S	1.7	3 to 5	Mar. 22	May 23	June 1	June 1	June 12	Sept. 29	Do.	
Grenache.															
Own roots.....	Cl	1901	1901		S			do	Mar. 28	May 31	June 1	June 7	Sept. 20	Oct. 24	
Dog Ridge.....	O	1904	1905	91	C, S	4.2	2 to 4	Mar. 24	May 20	June 2	May 26	June 18	Sept. 29	Oct. 15	
Lenoir.....	O	1903	1905	91	C, S	4	2 to 4	Mar. 20	May 21	June 2	May 27	June 18	Sept. 25	Oct. 6	
Rupprecht's St. George.....	O	1903	1905	90	S	4.7	2 to 4	Mar. 19	do	do	do	do	do	do	
Taylor Narbonne.....	O	1904	1906	88	S	2.8	2 to 4	Mar. 22	May 21	June 1	May 30	June 19	Sept. 29	Oct. 10	
Gros thine de Ladassane.															
Lenoir.....	O	1904	1907	86	S	2	3 to 6	Mar. 20	May 20	June 10	May 25	June 20	Sept. 28	Oct. 7	
Solanis X Othello.....	F	1907	1907	85	C, S	2.5	3 to 4	Mar. 16	May 10	May 21	May 29	June 25	Aug. 25		
Gros Guillaume.															
Lenoir.....	O	1904	1911	90	S	2.5			Mar. 16	May 26	May 30	June 5	Sept. 15	Sept. 25	
Gros Marzane.															
Own roots.....	O	1904	1904		S				Mar. 13	May 18	May 29	June 4	Sept. 25		
Gros Marce.															
Own roots.....	Cl	1905	1905		S				Mar. 13	May 18	May 29	June 4	Sept. 25		
Gros Verdlet.															
Own roots.....	Cl	1907	1905		S				Mar. 20	May 25	May 25	June 4	do		
Dog Ridge.....	F	1903	1906	92	S	4	3 to 6	Mar. 13	May 16	May 25	May 26	June 15	Aug. 24	Sept. 20	
Do.....	O	1901	1906	90	S	4.5	3 to 5	Mar. 17	May 23	May 31	May 30	June 15	Sept. 28	Oct. 5	
Lenoir.....	O	1903	1905	90	S	2.8	2 to 4	Mar. 18	May 23	June 3	May 27	June 14	Sept. 27	Do.	
Do.....	F	1903	1905	80	S	2.8	2 to 4	Mar. 14	May 23	June 3	May 27	June 14	Sept. 27	Do.	
Do.....	F	1903	1905	88	S	1.7	2 to 5	Mar. 12	May 14	May 20	May 23	June 29	Aug. 24	Sept. 19	
Rupprecht's St. George.....	F	1903	1905	88	S	1.7	2 to 5	Mar. 12	May 14	May 20	May 24	June 20	do	do	
Do.....	O	1903	1905	89	S	3.7	2 to 5	Mar. 20	May 20	May 31	May 26	June 30	Sept. 30	Oct. 10	
Taylor Narbonne.....	O	1904	1906	86	S	2.1	2 to 4	do	May 23	June 4	June 3	do	Sept. 27	Oct. 6	

Region	Year	Origin	Parentage	S	3, 4	2 to 4	Mar. 21	Apr. 8	May 25	June 3	May 30	June 16	Sept. 29	Oct. 12	
Hebron:	1904	Lenoir.	Rupestris St. George.	89	2.7	2 to 4	Mar. 21	Apr. 8	May 15	June 3	May 30	June 16	Sept. 29	Oct. 12	
	1907	F	Solonis X Othello, No. 1613.	91	5.7	2 to 4	Mar. 22	Apr. 8	May 20	June 3	June 2	June 4	Sept. 21		
	1907	F	Own roots.	95	7.5	2 to 4	Mar. 21	Apr. 8	May 20	June 3	June 4	June 4	Sept. 21		
	1904	Gi	Own roots.				Mar. 21	Apr. 3	May 24	June 10	May 28	June 16	Sept. 14		
	1904	Fi	Aramon X Rupestris Ganzin, No. 1.	88	3	2 to 5	Mar. 13	Apr. 3	May 15	June 8	May 28	June 16	Sept. 8	Sept. 22	
	1904	O	Do.	91	3	3 to 5	Mar. 26	Apr. 3	May 15	June 8	May 28	June 16	Sept. 5	Sept. 20	
	1909	F	Aramon X Rupestris Ganzin, No. 2.	85	3	3 to 5	Mar. 8	Apr. 3	May 27	June 10	May 28	June 16	Sept. 5	Sept. 20	
	1907	F	Australis.	90	2	2 to 5	Apr. 5	Apr. 5	May 23	June 6	May 28	June 16	Sept. 5	Sept. 20	
	1907	F	Berlandieri X Riparia, No. 420 A.	84	1	2 to 5	Apr. 1	Apr. 1	May 12	June 2	May 23	May 30	Sept. 10	Oct. 16	
	1903	F	Monticola X Riparia, No. 18808.	86	1.4	2 to 5	Apr. 2	Apr. 1	May 17	June 2	May 23	May 30	Sept. 10	Sept. 15	
1907	F	Mourvedre X Rupestris, No. 1202.	89	4.5	2 to 5	Mar. 22	Apr. 5	May 14	June 2	May 30	June 12	Sept. 16	Sept. 20		
1904	F	Riparia Gloire.	80	2.8	2 to 5	Mar. 23	Apr. 5	May 14	June 2	May 29	June 11	Sept. 4	Sept. 19		
1903	F	Riparia X (Cordifolia X Rupestris, No. 109-8).	85	4.4	2 to 5	Mar. 12	Apr. 1	May 21	June 4	May 29	June 11	Sept. 4	Sept. 19		
1907	F	Riparia X Rupestris, No. 101.	88	2.5	2 to 5	Apr. 1	Apr. 1	May 27	June 6	May 29	May 26	Sept. 5	Sept. 20		
1904	F	Do.	88	2.1	3 to 5	Mar. 24	Mar. 26	May 14	June 16	May 29	June 7	Sept. 4	Sept. 20		
1904	O	Riparia X Rupestris, No. 3390.	90	1.1	2 to 5	Mar. 24	Mar. 26	May 14	June 16	May 29	June 7	Sept. 4	Sept. 20		
1903	F	Rupestris St. George.	88	3.4	2 to 5	Mar. 23	Apr. 5	May 13	June 6	May 27	June 3	Sept. 28	Sept. 20		
1907	F	Solonis Robusta.	90	2.3	2 to 5	Mar. 12	Apr. 5	May 20	June 3	May 30	June 1	Sept. 4	Do.		
1907	F	Solonis Robusta.	88	5	2 to 5	Apr. 2	Apr. 3	May 24	June 6	May 30	June 1	Sept. 3	Sept. 19		
1904	F	Solonis X Riparia, No. 1616.	84	3	2 to 6	Mar. 7	Mar. 29	May 17	June 4	May 30	June 6	Sept. 10	Sept. 20		
1904	F	Taylor Narbonne.	92	3.1	2 to 6	Mar. 7	Mar. 29	May 17	June 4	May 30	June 6	Sept. 10	Sept. 20		
1907	F	Viala.	89	2	2 to 6	Apr. 1	Apr. 1	May 29	June 6	May 29	June 6	Sept. 17	Sept. 20		
Humata:	1904	F	Adobe Giant.	86	1.5	3 to 5	Mar. 8	Mar. 26	May 16	June 28	May 27	June 4	Sept. 5		
	1907	F	Australis.	91	4	3 to 5	Mar. 27	Mar. 26	May 21	June 16	May 27	June 4	Sept. 5		
	1907	F	Berlandieri X Riparia, No. 420 A.	90	4	3 to 5	Mar. 24	Mar. 26	May 21	June 16	May 27	June 4	Sept. 5		
	1907	F	Constantia.	92	5	3 to 5	Mar. 24	Mar. 26	May 23	June 16	May 27	June 4	Sept. 5		
	1903	F	Dog Ridge.	95	2.9	4 to 5	Mar. 23	Mar. 26	May 24	June 16	May 27	June 4	Sept. 5		
	1906	F	Riparia X Rupestris, No. 3390.	97	3	4 to 5	Mar. 5	Mar. 28	May 20	June 25	May 28	June 5	Sept. 5		
	1907	F	Solonis Robusta.	98	3	3 to 5	Mar. 30	Apr. 5	May 19	June 25	May 28	June 5	Sept. 5		
	1909	F	Solonis X Riparia, No. 1616.	98	3	3 to 5	Mar. 21	Mar. 28	May 23	June 25	May 28	June 5	Sept. 25		
	1907	F	Solonis (S. P. I. No. 6124).	96	8.3	3 to 5	Mar. 23	Apr. 1	May 16	June 25	May 30	June 5	Sept. 25		
	1904	O	Own roots.		8		Mar. 23	Apr. 1	May 16	June 25	May 30	June 5	Sept. 25		
Humata:	1904	O	Humata (S. P. I. No. 8583).		6.7	2 to 4	Mar. 16	Mar. 16	May 15	June 12	May 21	June 27	Oct. 10		
	1904	O	Own roots.		6.7	2 to 4	Mar. 16	Mar. 16	May 15	June 12	May 21	June 27	Oct. 10		
	1904	Gi	Own roots.				Mar. 29	Apr. 1	June 1	June 6	June 13	June 27	Oct. 10		
	Hyalas:	1904	O	Lenoir.	88	2.2	2 to 5	Mar. 18	Apr. 5	May 19	June 5	May 21	June 17	Sept. 28	Oct. 7
		1907	F	Rupestris Mission.	97	10	1 to 4	Mar. 22	Apr. 5	May 16	June 23	May 21	June 17	Sept. 28	Oct. 7
		1907	F	Rupestris St. George.	97	4.4	2 to 4	Mar. 20	Apr. 5	May 15	June 22	May 23	June 17	Sept. 28	Oct. 7
		1907	F	Solonis X Othello, No. 1613.	90	3	1 to 4	Mar. 17	Apr. 8	May 15	June 22	May 23	June 17	Sept. 28	Oct. 7
		1907	F	Own roots.		3	1 to 4	Mar. 17	Apr. 8	May 15	June 22	May 23	June 17	Sept. 28	Oct. 7
	Imperial:	1905	Gi	Imperial.				Mar. 26	Apr. 1	May 26	June 6	May 30	June 12	Sept. 25	
		1904	O	Inzolia Bianca.	84	1.8	2 to 4	Mar. 24	Apr. 1	May 22	June 7	May 27	June 20	Sept. 23	Oct. 7
1907		F	Rupestris St. George.	90	4.1	3 to 5	Mar. 23	Apr. 1	May 21	June 7	May 27	June 20	Sept. 23	Oct. 7	

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California.—Continued.

Variety and stock (on own roots, if so stated).	Experiment yard.	Year planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Johannisberger: Own roots.....	G1	1904			c, s	Lbs.	8	9	10	11	12	13	14	15	16
Jura Muscat:															
Leonor.....	F	1903	1905	81	s	8	2 to 5	Mar. 15	Mar. 26	May 29	June 3	June 3	Sept. 5	Sept. 5	Sept. 20
Riparia Gloire.....	F	1903	1905	89	s	3.3	3 to 5	Mar. 7	Mar. 25	May 4	May 18	May 20	May 25	Aug. 10	Sept. 19
Rupestris St. George.....	F	1903	1905	91	s	2.5	3 to 5	Mar. 8	Mar. 20	May 12	May 21	May 25	May 31	Aug. 20	Do.
Kabardak:															
Own roots.....	O	1904			s	8	2 to 3	Mar. 22	Mar. 22	May 16	June 1	..do.	June 20	Sept. 29	Oct. 15
Kadarka:															
Own roots.....	G1	1904			c, s			Mar. 25	Mar. 27	May 30	June 6	June 11	June 10	Sept. 15	Sept. 25
Vog Ridge.....	V	1903	1905	93	s	4.8	2 to 6	Mar. 14	Apr. 3	May 23	May 30	May 30	June 10	Sept. 10	Sept. 20
Leonor.....	F	1903	1905	92	c, s	3.3	2 to 6	Mar. 7	Apr. 1	May 20	May 27	May 27	June 4	..do.	Sept. 19
Riparia Gloire.....	F	1903	1905	90	c, s	3.9	3 to 7	Mar. 8	Apr. 5	..do.	May 27	May 28	June 7	Sept. 5	Do.
Rupestris St. George.....	F	1903	1905	95	c, s	4	1 to 6	Mar. 13	Apr. 1	May 11	May 20	May 24	May 30	Sept. 12	Do.
Keshu-Aly-Blanc.....															
Leonor.....	O	1904	1907	84	s	2.6	3 to 5	Mar. 21	Mar. 21	May 25	June 12	May 30	June 28	Sept. 23	Oct. 5
Solais X Othello, No. 1613.....	F	1907	1907	92	s	3.5		Mar. 30	Mar. 30	May 20	May 22	June 4	June 28	Sept. 23	Oct. 20
Leonor.....															
Riparia X Rupestris, No. 3309.....	O	1904	1907	90	c, s	4	2 to 5	Mar. 22	Apr. 5	..do.	June 5	May 29	June 23	Sept. 23	Oct. 20
Leonor.....	O	1904	1907	87	s	3.5		Mar. 20	Apr. 5	May 16	May 23	June 2	June 23	Sept. 23	Oct. 20
Herbemont.....	O	1904	1909	47	s	9		Mar. 22	Apr. 5	June 5	June 12	June 12	June 17	..do.	Do.
Leonor.....	O	1904	1906	83	s	2	1 to 5	Mar. 24	Apr. 5	May 27	June 13	June 1	June 22	Sept. 30	Do.
Kleinburger.....															
Own roots.....	G1	1904			s			Mar. 10	Apr. 5	May 22	May 30	May 27	June 5	Aug. 29	Sept. 23
Köber.....															
Own roots.....	G1	1904	1906	88	s	2.6	2 to 4	..do.	Mar. 30	May 27	..do.	May 31	..do.	Sept. 10	Sept. 25
Dog Ridge.....	O	1904	1906	88	c, s	2.4	2 to 6	Mar. 17	Mar. 17	..do.	June 7	June 19	..do.	Sept. 26	Oct. 3
Leonor.....	O	1903	1905	82	s	2.8	3 to 5	Mar. 15	Mar. 15	..do.	June 8	May 27	..do.	..do.	Oct. 15
Rupestris St. George.....	O	1903	1905	82	s	2.8	3 to 5	Mar. 16	Mar. 16	..do.	June 1	May 25	..do.	..do.	Oct. 10
Taylor Narbonne.....	O	1904	1906	73	s	8	2 to 4	Mar. 18	Mar. 18	May 22	..do.	..do.	June 15	..do.	Oct. 4

Kürstl. Miel:	1904	1907	81	s	2.1	3 to 5	Mar. 25	Mar. 29	May 25	June 9	May 29	June 25	Oct. 7	Oct. 10
Lennor:				s			Mar. 20	Mar. 29	May 30	June 1	June 6	June 15	Sept. 24	Oct. 10
Kürstl. s:				s			Mar. 25		May 19	June 8	May 24	June 18	Sept. 25	Sept. 25
Do.				s					May 15	May 22	May 21	May 25	Aug. 24	Sept. 19
Lahntraube:				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Do.				s			Mar. 13	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Riparia Gloire:				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Riparia St. George:				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Leani Zelo:				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Rupestris St. George:				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Lignan Blanc:				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Do.				s			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Aug. 24	Do.
Aramon X Rupestris Ganzin, No. 1:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Do.				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Dog Ridge:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Do.				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Hofortrup:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Do.				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Monvredre X Rupestris, No. 1202:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Do.				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Riparia Gloire:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Riparia X Rupestris, No. 101:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Riparia X Rupestris, No. 3309:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Rupestris St. George:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Do.				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Solomis X Othello:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Solomis X Riparia, No. 1616:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Taylor Narbonne:				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
Do.				C, S			Mar. 12	Mar. 25	May 12	May 18	May 20	May 25	Sept. 19	Do.
L'istan:				s			Mar. 18	Mar. 31	May 21	June 11	May 26	June 15	Sept. 25	Oct. 4
Do.				s			Mar. 18	Mar. 31	May 21	June 11	May 26	June 15	Sept. 25	Oct. 4
Own roots:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
(Aestivalis X Monticola) X (Riparia X Rupestris, No. 554-5):				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Aramon X Rupestris Ganzin, No. 1:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Do.				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Berlandieri X Riparia, No. 429B:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Columboid X Riparia, No. 2502:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Constantia:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Cornucopif:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
De Grassoff:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Dog Ridge:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Do.				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Herbertmont:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Do.				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Monticola X Riparia, No. 18804:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Monticola X Riparia, No. 18808:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Monvredre X Rupestris, No. 1202:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Riparia X Rupestris, No. 101:				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Do.				C, S			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23
Riparia X Rupestris, No. 401-14:				s			Mar. 16	Mar. 31	May 22	June 1	May 27	June 6	Sept. 20	Sept. 23

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment yard.	Year planted was	Year grafted.	Congeniality.	How pruned.	Weight per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Listan—Continued.															
Riparia X Rupestris, No. 3306.	O	1907	1909	P. et.	S	1.3	3 to 4	Mar. 18	Apr. 2	May 31	June 8	June 4	June 21	Sept. 25	Late season.
Riparia X Rupestris, No. 3369.	F	1907	1907		C, S	3.2	4 to 5	Mar. 21	Apr. 1	May 12	May 21	May 26	June 11	Sept. 21	Early season.
Do.	O	1904	1906		S	2.3	3 to 4	Mar. 18	Mar. 18	May 23	June 8	May 28	June 14	Sept. 26	
Rupestris Martin.	O	1903	1905		S	3.8	2 to 5	Mar. 20	Mar. 20	May 22	June 5	June 13	June 20	Sept. 22	
Rupestris Melanica.	O	1907	1909		S	1.2	3 to 4	Mar. 17	Mar. 17	June 3	June 8	June 13	June 20	Sept. 25	
Rupestris Mission.	F	1907	1909		S	1.9	3 to 4	do.	Apr. 1	June 3	May 20	June 1	June 13	Sept. 25	
Rupestris St. George.	F	1903	1905		S	1.5	2 to 5	Mar. 18	Mar. 18	May 16	May 20	May 22	June 1	Sept. 18	
Do.	F	1903	1905		S	3.2	2 to 5	Mar. 22	Mar. 22	May 22	June 1	May 27	June 13	Sept. 21	
Rupestris X Perlandieri, No. 219A.	F	1907	1907		C, S	3.2	1 to 5	Mar. 22	Mar. 22	May 22	June 1	May 27	June 10	Sept. 21	
Salt Creek.	F	1903	1907		S	3.1	2 to 5	Mar. 19	Apr. 2	May 15	June 5	May 27	June 10	Sept. 21	
Sotom's Robusta.	F	1907	1907		C, S	4.1	2 to 4	Mar. 18	Apr. 2	June 15	June 10	June 7	June 18	Aug. 25	
Do.	O	1907	1907		S	4.8	5 to 5	Mar. 25	Apr. 1	June 16	June 22	June 30	July 5	Sept. 21	
Sotom's X Otello.	F	1907	1907		C, S	1.2	3 to 4	Mar. 20	Apr. 1	June 20	June 3	June 8	June 10	Sept. 25	
Sotom's X Otello, No. 1613.	O	1907	1909		C, S	3.2	2 to 4	Mar. 18	Mar. 18	May 22	June 7	May 30	June 11	Sept. 21	
Sotom's X Riparia, No. 1616.	O	1904	1906		C, S	3.3	2 to 4	Mar. 16	Mar. 16	May 22	June 7	May 28	June 8	Sept. 21	
Talbot's Carbone.	O	1904	1906		C, S	3.5	2 to 4	Mar. 16	Mar. 16	May 22	June 7	May 28	June 8	Sept. 21	
Valencia.	F	1907	1907		C, S	3.5	2 to 5	Mar. 19	Apr. 1	May 15	May 23	May 31	June 1	Aug. 25	
Luglienza No. 1.	F	1904	1906		S	1.5	3 to 5	Mar. 13	do.	May 18	May 20	May 25	May 30	Sept. 14	Do.
Adobe Giant.	F	1903	1905		S	1.7	3 to 6	Mar. 7	Apr. 3	May 11	do.	May 21	May 29	Aug. 25	Sept. 20
Deer Ridge.	F	1907	1907		C, S	1.5	4 to 6	Mar. 15	Mar. 15	May 23	May 19	May 21	May 23	Aug. 25	Sept. 19
Riparia X Rupestris, No. 3300.	F	1907	1907		C, S	2.1	4 to 6	Mar. 11	Mar. 25	May 10	May 19	May 21	May 23	Aug. 25	Sept. 19
Rupestris St. George.	F	1903	1905		C, S	2.1	4 to 6	Mar. 11	Mar. 25	May 10	May 19	May 21	May 23	Aug. 25	Sept. 19
Macrum.					S			Mar. 10	Apr. 5	May 20	May 25	May 25	June 5	Sept. 4	
Own roots.	GH	1907			S			do.	Mar. 30	May 25	June 2	June 1	June 7	Aug. 29	
Madreline Angevine.	GH	1905			S			do.	Mar. 30	May 25	June 2	June 1	June 7	Aug. 29	
Own roots.	O	1904	1907		S	1.8	2 to 5	Mar. 15	Mar. 15	May 19	May 30	May 24	June 11	Sept. 20	Oct. 2
Dog Ridge.	O	1903	1905		S	2.6	2 to 4	Mar. 15	Mar. 15	May 21	May 26	May 28	June 6	do.	Oct. 8
Lenoir.	O	1903	1905		S	5.1	2 to 4	Mar. 16	Mar. 16	May 23	May 31	June 1	June 15	do.	Oct. 2
Rupestris St. George.	O	1903	1905		S	5.1	2 to 4	Mar. 16	Mar. 16	May 23	May 31	June 1	June 15	Sept. 22	Sept. 25
Madreline Blanche.					C, S			Mar. 25	Mar. 29	May 30	June 6	June 3	June 11	Aug. 28	Sept. 19
Own roots.	GH	1905			C, S			Mar. 25	Mar. 29	May 30	June 6	June 3	June 11	Aug. 28	Sept. 19

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experimental vineyards in California—Continued.

Variety and stock (on own roots, if so stated)	Experiment vine-	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Mamelon:															
Own roots.....	GI	1907		<i>P. et.</i>	s	<i>Lbs.</i>	8	9	10	11	12	13	14	15	16
Dog Ridge.....	O	1904	1907	91	c, s	3.1	3 to 5	Mar. 15	Apr. 8	May 20	May 30	May 30	June 2	Sept. 22	Oct. 8
Lenoir.....	O	1903	1905	94	c, s	4.2	2 to 5	Mar. 16	Mar. 25	May 23	June 3	May 27	June 13	Sept. 19	Oct. 6
Rupestris St. George.....	O	1903	1905	90	s	4.7	2 to 4	Mar. 17	Mar. 17	May 21	June 8	May 30	June 13	Sept. 21	Oct. 8
Taylor Narbonne.....	O	1904	1906	87	s	2.3	3 to 5	Mar. 21	Mar. 21do.do.	May 29	June 16	Sept. 20	Oct. 4
Valencia.....	F	1907	1907	89	c, s	.9	Mar. 12	Mar. 12do.do.do.do.	Oct. 17
Manteno de Pillas:															
Own roots.....	GI	1904													
Adobe Giant.....	F	1904	1906	88	c, s	2.4	3 to 4	Mar. 11	Mar. 24	May 26	June 3	May 30	June 9	Sept. 20	Sept. 25
Aramon X Rupestris Ganzin, No. 1.....	F	1904	1906	86	c, s	2.6	3 to 6	Mar. 10	Mar. 25	May 13	May 19	May 27	May 27	Sept. 15	Sept. 30
Dog Ridge.....	F	1903	1905	86	c, s	2.5	3 to 5	Mar. 13	Mar. 22	May 12do.	May 23	May 30do.do.
Herbemont.....	F	1903	1905	85	c, s	1	3 to 5	Mar. 7	Mar. 15	May 13	May 20	May 21	May 21	Sept. 15do.
Lenoir.....	F	1903	1905	87	s	2.6	3 to 5do.	Mar. 27	May 15	May 24	May 24	May 24	Sept. 15do.
Do.....	F	1903	1905	89	s	2.6	3 to 4	Mar. 24	Mar. 27	May 20	June 1	May 20	May 24	Sept. 23	Oct. 9
Mourvedre X Rupestris, No. 1292.....	F	1904	1906	83	c, s	3.7	3 to 5	Mar. 12	Mar. 26	May 15	May 20	May 21	May 30	Sept. 10	Sept. 23
Riparia Cloire.....	F	1903	1905	83	c, s	2.8	3 to 5do.	Mar. 20	May 10	May 21	May 25	May 30	Sept. 14	Sept. 19
Riparia X Rupestris, No. 101.....	F	1904	1906	89	c, s	2.9	3 to 6do.	Mar. 30	May 12	May 22do.	May 28	Sept. 14	Sept. 20
Riparia X Rupestris, No. 5309.....	F	1904	1905	83	c, s	3.3	3 to 5	Mar. 1	Mar. 26	May 17	May 20do.	June 4	Sept. 8	Do.
Rupestris Marini.....	F	1903	1905	89	s	3.1	3 to 6	Mar. 13	Mar. 25	May 14	May 18	May 21	June 2	Sept. 10	Sept. 19
Rupestris St. George.....	F	1903	1905	89	c, s	3.1	2 to 4	Mar. 20	Mar. 25	May 21	June 2	May 27	June 15	Sept. 22	Sept. 30
Salt Creek.....	O	1903	1908	87	s	1.7	2 to 6	Mar. 6	Mar. 30	May 13	May 21	May 21	June 3	Sept. 25	Sept. 30
Solonis X Riparia, No. 1616.....	F	1903	1905	86	s	4	4 to 6	Mar. 12	Mar. 20	May 14	May 23	May 23	May 28	Sept. 13	Sept. 16
Taylor Narbonne.....	F	1904	1906	91	c, s	3.2	2 to 5	Mar. 6	Mar. 25	May 15do.do.do.do.do.
Marschano:															
Own roots.....	GI	1905													
Lenoir.....	O	1904		84	c, s	3.5	Mar. 25	Mar. 30	May 30	June 6	June 5	June 12	Sept. 23	Sept. 24
Maryville de Malaga:															
Lenoir.....	O	1904		85	s	2.1	3 to 5do.do.	May 20	June 8	May 27	June 24	Sept. 24	Oct. 20
Riparia X Rupestris, No. 3309.....	F	1907	1907	92	c, s	6.3	Mar. 13	Apr. 8	May 20	May 23do.do.	Sept. 25	Sept. 25
Solonis X Othello, No. 1613.....	F	1907	1907	87	s	2.1	3 to 5	Mar. 11do.	May 21	May 26	June 2do.	Sept. 20	Sept. 21

	GI	1904				Mar. 23	Apr. 10	May 29			May 29				Sept. 24
Marmola Tobend:															
Own roots.....															
Marmora:															
Lenoir.....	F	1905	85	S	1.5	2 to 6	Mar. 15	Apr. 3	May 18	May 20	May 24	May 24	May 24	May 30	Sept. 27
Riparia Gloire.....	F	1905	85	S	2.2	3 to 6	Mar. 8	Apr. 1	May 15	May 15	May 25	May 25	May 25	June 7	Sept. 19
Rupestris St. George.....	F	1905	86	C, S	2	3 to 7	Mar. 15	Mar. 28	May 18	May 25	May 25	May 25	May 25	June 7	Do.
Marsanne:															
Own roots.....	GI	1904		S			do.	Mar. 27	May 26	June 4	May 31	June 9	June 9	Aug. 28	Sept. 22
Marzemino:															
Own roots.....	GI	1904		S			Mar. 12	Apr. 3	do.	June 5	June 2	June 2	June 2	Sept. 20	Sept. 23
Do.....	O	1904		S		3 to 4	Mar. 23	Mar. 23	May 9	June 5	May 15	June 17	June 17	Sept. 23	Sept. 23
Mataro:															
Own roots.....	GI	1907		S			Mar. 20	Apr. 1	May 28	June 7	May 29	June 20	June 20	do.	Oct. 15
Aramon X Rupestris Ganzin, No. 1.....	O	1903	90	S	3.4	2 to 5	Mar. 18	Mar. 18	May 24	June 9	June 1	June 14	June 14	Sept. 27	Oct. 10
Dog Ridge.....	O	1905	92	S	2.4	2 to 5	do.	do.	May 24	June 4	May 28	June 10	June 10	Sept. 28	Oct. 11
Herbent.....	O	1903	86	S	1.2	3 to 5	do.	do.	May 21	June 6	May 29	June 12	June 12	Sept. 28	Oct. 8
Lenoir.....	O	1903	96	S	3.2	2 to 6	Mar. 22	Mar. 22	do.	June 5	do.	June 20	June 20	Sept. 28	Oct. 6
Mourvadre X Rupestris, No. 1202.....	O	1904	89	S	3.1	2 to 6	Mar. 17	Mar. 17	May 22	June 6	May 27	June 11	June 11	Sept. 27	Oct. 11
Rupestris Martin.....	O	1903	91	S	3.3	3 to 5	do.	do.	May 22	June 5	May 29	June 9	June 9	Sept. 24	Oct. 10
Rupestris St. George.....	O	1903	91	S	3.7	2 to 4	do.	do.	May 23	June 5	May 29	June 16	June 16	Sept. 7	Oct. 11
Salt Creek.....	O	1905	89	S	1.7	2 to 4	Mar. 20	Mar. 20	May 23	June 12	do.	June 16	June 16	Sept. 7	Oct. 11
Taylor Narbonne.....	O	1904	88	S	2.8	2 to 4	Mar. 24	Mar. 24	do.	June 8	do.	June 22	June 22	Sept. 27	Oct. 10
Melon:															
Dog Ridge.....	F	1905	90	S	3.6	4 to 6	Mar. 12	Mar. 30	May 15	May 25	May 27	June 10	June 10	Aug. 25	Sept. 20
Rupestris St. George.....	F	1905	88	S	2.1	4 to 5	Mar. 11	Mar. 28	May 12	May 17	May 20	May 27	May 27	do.	Sept. 19
Merlot:															
Own roots.....	GI	1904		S			Mar. 15	Apr. 1	May 26	May 30	June 5	June 5	June 5	Sept. 15	Sept. 15
Meslier:															
Own roots.....	GI	1905		S			Mar. 8	Apr. 3	May 20	do.	May 31	June 5	June 5	do.	Sept. 22
Aramon X Rupestris Ganzin, No. 1.....	O	1903	92	S	5	2 to 3	Mar. 19	Mar. 19	May 29	May 31	June 5	June 16	June 16	Sept. 25	Oct. 10
Lenoir.....	O	1905	89	S	4	2 to 5	Mar. 16	Mar. 16	May 26	do.	May 29	June 12	June 12	Sept. 20	Oct. 8
Rupestris St. George.....	O	1905	92	S	4.3	2 to 4	Mar. 21	Mar. 21	May 21	May 30	May 26	do.	do.	do.	Oct. 10
Meunier:															
Own roots.....	GI	1905		S			Mar. 15	Apr. 1	May 22	May 26	do.	June 4	June 4	Aug. 23	Oct. 23
Dog Ridge.....	O	1906	88	S	3.7	2 to 4	Mar. 20	Mar. 20	May 20	June 5	May 24	June 17	June 17	Sept. 20	Oct. 3
Lenoir.....	O	1905	85	S	3.4	3 to 5	Mar. 16	Mar. 16	May 20	May 31	May 26	June 13	June 13	Sept. 25	Oct. 18
Rupestris St. George.....	O	1905	90	C, S	4.3	2 to 5	Mar. 17	Mar. 17	May 22	do.	May 27	June 7	June 7	Sept. 15	Oct. 8
Meyer, No. 59:															
Dog Ridge.....	O	1906	90	S	2.5	2 to 5	Mar. 18	Mar. 18	May 26	June 8	June 1	June 16	June 16	Sept. 24	Oct. 4
Lenoir.....	O	1906	89	S	3.2	3 to 4	Mar. 20	Mar. 20	May 23	June 12	May 27	June 17	June 17	Sept. 25	Oct. 4
Meyer, No. 60:															
Dog Ridge.....	O	1906	93	S	3.1	3 to 5	Mar. 23	Mar. 23	May 26	June 7	June 2	June 14	June 14	Sept. 23	Oct. 1
Lenoir.....	O	1906	88	C, S	3.9	3 to 4	Mar. 20	Mar. 20	May 25	June 5	May 29	June 19	June 19	Sept. 25	Oct. 1
Meyer, No. 65:															
Lenoir.....	O	1906	93	S	4.3	3 to 5	Mar. 18	Mar. 18	May 21	June 8	May 25	June 20	June 20	Sept. 28	Oct. 20
Meyer, No. 91:															
Dog Ridge.....	O	1906	95	S	4.3	3 to 5	Mar. 17	Mar. 17	May 22	May 31	May 27	June 12	June 12	Sept. 27	Oct. 11
Lenoir.....	O	1906	91	S	3.3	3 to 4	Mar. 20	Mar. 20	May 20	June 4	May 23	June 17	June 17	Sept. 26	Oct. 11
Meyer, No. 95:															
Dog Ridge.....	O	1906	93	S	5.9	3 to 5	do.	do.	May 26	June 5	June 2	June 14	June 14	Sept. 29	Oct. 4
Lenoir.....	O	1906	93	S	5.3	3 to 5	Mar. 17	Mar. 17	May 22	do.	May 25	June 17	June 17	Sept. 30	Oct. 7

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting an resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year planted was	Year grafted.	Congeniality.	How pruned.	Weight of fruiting per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
Meyer, No. 103: Lenoir.....	O	1901	1906	P. cl. 88	s	2.4	8	9	10	11	12	13	14	15	16
Meyer, No. 107: Lenoir.....	O	1904	1906	94	c, s	4.4	2 to 4	Mar. 18	May 22	June 9	May 26	June 17
Meyer, No. 116: Dog Ridge.....	O	1904	1906	94	s	3.2	3 to 5	Mar. 19	May 26	June 4	June 1	June 8	Oct. 7	Oct. 10
Meyer, No. 51E: Dog Ridge.....	O	1904	1906	85	s	2.1	3 to 4	May 23	May 29	June 11	Sept. 29	Oct. 15
Millenium: Lenoir.....	O	1904	1906	89	s	3.3	May 24	Sept. 26	Oct. 10
Aramon X Rupestris Ganzin, No. 1: Lenoir.....	O	1904	1906	74	s	3.7	2 to 4	Mar. 16	May 21	June 2	May 26	June 12	Sept. 26	Oct. 9
Riparia X Rupestris, No. 101: Riparia X Rupestris, No. 336B.	O	1904	1907	60	s	1.2	3 to 5	Mar. 22	May 21	June 1	May 28	June 10	Sept. 25	Oct. 2
Solonis X Riparia, No. 1616: Mission: Own roots.....	O	1904	1907	68	s	1.3	3 to 4	May 20	June 1	May 25	June 13	Sept. 27	Oct. 6
Adobe Giant.....	G	1904	1906	90	c, s	4.3	Mar. 22	Mar. 27	May 26	June 4	May 30	June 9	Sept. 15	Sept. 28
Dog Ridge.....	F	1903	1905	92	c, s	4	3 to 4	Mar. 11	Apr. 1	May 1	May 25	May 8	May 30	Sept. 5	Sept. 21
Herbemont.....	F	1903	1905	81	s	3.8	2 to 4	Mar. 14	Mar. 30	May 15	June 8	May 28	June 7	Sept. 5	Sept. 20
Do.....	F	1903	1905	82	c, s	4.8	2 to 5	Mar. 22	Apr. 1	May 20	June 5	May 29	June 14	Sept. 28	Oct. 3
Do.....	F	1903	1905	81	s	4.1	2 to 5	Mar. 20	May 22	June 0	May 30	June 6	Sept. 3	Sept. 20
Do.....	F	1903	1905	84	s	1.3	3 to 6	Mar. 9	May 15	June 0	May 28	June 21	Sept. 23	Oct. 11
Do.....	F	1903	1905	83	s	4	2 to 4	Mar. 20	May 23	June 1	May 30	June 11	Sept. 18	Sept. 27
Riparia Gloire.....	O	1903	1905	80	c, s	3.3	2 to 5	Mar. 12	Mar. 25	May 20	June 7	May 24	June 28	Sept. 19	Sept. 24
Rupestris Mission.....	F	1903	1905	87	c, s	2.4	2 to 4	Mar. 14	Mar. 26	May 27	June 4	May 21	June 8	Sept. 1	Sept. 10
Do.....	F	1903	1905	86	s	2.8	2 to 4	Mar. 21	May 23	June 0	May 25	June 2	Sept. 1	Sept. 10
Rupestris St. George.....	F	1903	1905	85	c, s	5	3 to 5	Mar. 20	May 28	June 6	May 26	June 10	Sept. 10	Sept. 20
Do.....	F	1903	1905	83	c, s	5.1	2 to 5	Mar. 20	May 28	June 6	May 26	June 10	Sept. 8	Sept. 20
Salt Creek.....	F	1903	1905	83	c, s	2.3	2 to 5	Mar. 13	Apr. 1	May 23	June 10	May 29	June 20	Sept. 26	Oct. 10
Do.....	O	1903	1905	91	s	2.8	2 to 5	Mar. 22	May 21	June 10	May 26	June 20	Sept. 26	Oct. 10

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Muscad (S. P. I. No. 3063): Own roots.....	O	1904	4	P. et.	S	2	3 to 4	Mar. 20		May 12	June 12	May 17	June 16	Oct. 10	16
Muscad Albaridians: Own roots.....	G	1905			S			do.	Mar. 28	do.	May 29	May 18	June 4	Aug. 27	Sept. 25
Aramon X Rupestris Gamzin, No. 1. Lenoir.....	F	1904	1905	87	S	2.1	2 to 6	Mar. 22	Apr. 1	May 20	May 29	May 27	May 31	Aug. 20	Sept. 25
Mourvedre X Rupestris No. 1202. Riparia Gloire.....	F	1903	1905	87	S	1.7	2 to 6	Mar. 13	Mar. 30	May 12	May 23	May 23	May 27	Aug. 23	Sept. 15
Riparia X Rupestris, No. 101. Riparia X Rupestris, No. 3389. Riparia X Rupestris, No. 3389. Rupestris St. George.....	F	1903	1905	82	S	4.8	2 to 5	Mar. 14	do.	May 15	May 20	May 21	June 1	Aug. 18	Sept. 20
Riparia X Rupestris, No. 101. Riparia X Rupestris, No. 3389. Rupestris St. George.....	F	1904	1905	87	S	2.6	2 to 5	Mar. 20	Apr. 1	May 15	May 18	May 22	May 26	Aug. 20	Sept. 19
Riparia X Rupestris, No. 101. Rupestris St. George.....	F	1904	1905	91	S	2.7	2 to 5	Mar. 13	Mar. 30	May 16	May 23	May 24	do.	do.	Sept. 20
Rupestris St. George.....	F	1903	1905	92	S	5.7	2 to 5	Mar. 23	Apr. 1	May 12	May 23	May 21	May 28	Aug. 8	Do.
Solonis X Riparia, No. 1616. Taylor Narbonne.....	F	1904	1905	85	S	4.3	3 to 5	Mar. 24	do.	May 14	May 16	do.	May 29	Aug. 20	Sept. 2
Muscad Bonod: Berlandieri X Riparia, No. 420A. De Grasset.....	F	1904	1905	88	S	1.4	3 to 6	Mar. 21	Mar. 30	May 12	May 20	May 25	May 27	Aug. 18	Sept. 2
Lenoir.....	F	1909	1909	80	S	3	3 to 6	Mar. 12	do.	May 13	do.	May 24	May 26	Aug. 15	Sept. 20
Riparia X Rupestris, No. 101. Riparia X Rupestris, No. 3309. Rupestris St. George.....	F	1909	1907	80	S	1	3 to 5	Mar. 22	Apr. 8	May 18	May 22	May 29	do.	Sept. 15	Oct. 25
Solonis Robusta. Valencia.....	F	1907	1907	82	S	1.7	3 to 5	Mar. 26	do.	May 23	June 10	May 28	June 25	Sept. 17	Oct. 25
Muscad Capulines: Own roots.....	G	1905	1905	79	S	2.8	2 to 5	Mar. 18	Mar. 27	May 28	May 27	May 30	June 10	Sept. 10	Sept. 20
Deg Ridge.....	O	1904	1905	85	S	2.1	2 to 4	Mar. 26	do.	do.	June 11	do.	June 15	Sept. 20	Sept. 21
Rupestris St. George.....	O	1903	1905	85	S	3.4	2 to 5	Mar. 18	do.	May 21	May 31	May 26	June 14	Sept. 9	Sept. 23
Muscadeller: Own roots.....	G	1905	1905	89	S	2.4	3 to 5	Mar. 17	Mar. 28	do.	do.	June 8	Aug. 27	Sept. 20	Sept. 20
Deg Ridge.....	F	1903	1905	87	S	1.2	3 to 6	Mar. 8	Mar. 25	May 14	May 20	May 25	May 29	Aug. 15	Do.
Riparia Gloire.....	F	1903	1905	76	S	1.2	3 to 5	Mar. 12	Mar. 25	May 10	May 14	May 16	May 24	Aug. 24	Sept. 10
Rupestris St. George.....	F	1903	1905	89	S	2.3	3 to 5	Mar. 10	do.	May 16	May 20	May 24	May 29	Aug. 21	Sept. 19
Rupestris St. George.....	F	1903	1905	89	S	2.3	3 to 5	Mar. 13	do.	May 8	May 13	May 14	May 24	Aug. 20	Do.

Muscad Gros Noir Hatif: Own roots.....	G1	1905	90	S	2,9	3 to 4	Mar. 18	Mar. 27	May 14	May 27	May 20	June 1	Aug. 27	Sept. 23
Dog Ridge.....	O	1904	84	S	2,2	2 to 5	Mar. 17	Mar. 27	May 25	May 31	May 27	June 10	Sept. 15	Sept. 15
Lenoir.....	O	1903	84	S	2,9	3 to 6	Mar. 18	Mar. 27	May 22	June 1	May 28	June 12	do.	Oct. 6
Rupestris St. George.....	O	1903	86	S	2,9	3 to 6	Mar. 18	Mar. 27	May 22	June 1	May 28	June 14	Sept. 10	Oct. 6
Taylor Narbonne.....	O	1904	84	S	2,6	1 to 4	do.	Mar. 27	May 24	do.	May 29	do.	Sept. 15	Oct. 4
Muscad Hamburg: Own roots.....	G1	1905	86	C, S	3	2 to 4	Mar. 24	Mar. 28	May 26	May 27	May 30	June 3	Aug. 27	Sept. 23
Aramon X Rupestris Ganzin, No. 1. Dog Ridge.....	F	1907	86	S	3,9	2 to 5	Mar. 24	Apr. 1	May 16	May 23	May 27	June 3	Aug. 20	Sept. 23
Lenoir.....	O	1904	87	S	2	3 to 5	Mar. 24	Apr. 1	May 25	June 5	May 30	June 17	Sept. 25	Oct. 4
Monticola X Rupestris.....	F	1903	83	S	2,2	3 to 4	Mar. 26	Apr. 1	May 30	June 1	June 6	June 15	do.	Oct. 4
Riparia X Rupestris, No. 3389.....	F	1907	83	S	3,9	2 to 6	Mar. 23	Apr. 2	May 15	May 22	May 28	June 15	Aug. 21	Aug. 25
Rupestris St. George.....	F	1903	90	S	1,6	2 to 6	Mar. 17	Apr. 2	May 19	June 3	May 21	June 16	Aug. 15	Aug. 25
Rupestris X Berlandieri, No. 219A.....	F	1905	80	S	1,8	3 to 5	Mar. 20	Apr. 2	May 17	June 3	May 24	June 19	Sept. 15	Sept. 19
Taylor Narbonne.....	O	1907	84	S	1,8	3 to 5	Mar. 19	Apr. 2	May 21	June 22	May 29	June 17	Aug. 13	Oct. 4
Muscad Hamburg Noir d'Hongrie: Own roots.....	G1	1906	80	S	2,2	3 to 6	Mar. 19	Apr. 2	May 21	June 22	May 27	June 17	Sept. 25	Oct. 4
Muscad Noir d'Hongrie: Lenoir.....	F	1905	83	S	1,7	2 to 5	Mar. 14	Mar. 30	May 14	June 1	May 30	June 6	Aug. 27	Sept. 20
Do.....	F	1903	90	S	2,8	2 to 5	Mar. 22	Mar. 30	May 14	June 4	May 18	May 26	Aug. 24	Sept. 18
Riparia Gloire.....	O	1903	90	C, S	3,6	2 to 5	Mar. 10	Apr. 1	May 12	May 18	May 22	May 29	Sept. 25	Oct. 6
Rupestris St. George.....	F	1903	89	S	3,7	2 to 5	Mar. 14	Mar. 25	May 10	May 16	May 19	May 25	Aug. 22	Sept. 19
Do.....	O	1903	90	S	5,4	2 to 5	Mar. 16	Mar. 16	May 22	June 1	May 27	June 14	Sept. 15	Oct. 8
Muscad Noir Precoce: Aramon X Rupestris Ganzin, No. 2. Dog Ridge.....	F	1907	91	S	3,5	3 to 6	Mar. 21	Apr. 1	May 16	May 23	May 26	June 13	Aug. 25	Aug. 29
Lenoir.....	O	1904	81	S	9	2 to 4	Mar. 22	Apr. 1	May 23	May 31	May 29	June 13	Sept. 28	Oct. 8
Luifata.....	O	1903	88	S	2,9	2 to 4	Mar. 18	Apr. 1	May 26	June 3	June 1	June 10	do.	Do.
Monticola X Riparia, No. 18804.....	F	1907	92	C, S	5	3 to 5	Mar. 25	Apr. 1	May 16	May 22	May 29	June 13	Aug. 20	Aug. 25
Rupestris St. George.....	F	1907	92	C, S	6,3	3 to 5	Mar. 21	do.	do.	do.	do.	do.	do.	do.
Rupestris X Berlandieri, No. 219A.....	O	1903	91	S	4	3 to 4	Mar. 20	Apr. 1	May 24	May 30	do.	June 10	Sept. 20	Sept. 30
Solanis X Otthello.....	F	1907	88	S	2,8	3 to 6	do.	Apr. 1	May 15	May 22	May 27	June 10	Aug. 20	Aug. 24
Taylor Narbonne.....	F	1907	89	S	3,5	3 to 6	Mar. 25	Apr. 1	May 22	June 4	May 28	June 16	Sept. 25	Oct. 8
Muscad Rose: Dog Ridge.....	O	1904	85	S	2	2 to 5	Mar. 22	Mar. 22	May 23	June 4	June 28	June 16	Sept. 25	Oct. 8
Lenoir.....	F	1903	88	C, S	2,8	3 to 7	Mar. 7	Mar. 30	May 18	May 18	May 24	May 28	Aug. 26	Sept. 20
Riparia Gloire.....	F	1903	82	S	1	2 to 6	Mar. 6	Mar. 25	May 12	do.	May 21	May 25	Aug. 5	Sept. 10
Rupestris St. George.....	F	1903	88	C, S	3,9	2 to 6	Mar. 12	do.	May 15	May 16	May 31	May 31	Aug. 1	Sept. 19
Muscad Rouge de Madre: Lenoir.....	F	1903	86	C, S	3,3	3 to 6	do.	Mar. 28	May 12	May 15	May 18	May 27	Aug. 10	Do.
Riparia Gloire.....	F	1903	76	S	1	3 to 5	Mar. 16	Mar. 30	May 11	do.	May 19	May 24	Aug. 5	Sept. 19
Rupestris St. George.....	F	1903	86	C, S	3,3	2 to 6	Mar. 10	Apr. 1	May 10	May 21	May 24	May 26	Aug. 3	Do.
Muscad Talabot: Own roots.....	G1	1905	86	S	4,3	2 to 5	Mar. 12	Mar. 28	May 12	May 15	May 18	May 25	Aug. 7	Sept. 19
Lenoir.....	O	1903	86	S	3,3	3 to 6	do.	Apr. 3	May 26	June 5	June 1	June 18	Sept. 18	Sept. 25
Rupestris St. George.....	O	1903	86	S	4,5	3 to 6	Mar. 23	do.	May 25	June 5	May 27	June 14	do.	Do.
Napolcon: Own roots.....	G1	1904	86	S	4,5	3 to 6	do.	Apr. 1	May 24	June 6	May 29	June 13	Sept. 10	Sept. 22

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment yard.	Year planted was	Year grafted	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
I		3	4	5	6	7	8	9	10	11	12	13	14	15	16
Nasa Valentina: Own roots.....	Gi	1905			c, s	Lbs.		Mar. 8	Mar. 30	May 25	May 30	June 1	June 3	Sept. 21	Sept. 19
Riparistis St. George.....	F	1903	1903	83		1.8	2 to 5	Mar. 3	Mar. 15	May 10	May 15	May 20	May 28	Aug. 25	
Nebbiolo: Own roots.....	Gi	1905	1906		\$		2 to 4	Mar. 15	Apr. 3	May 22	June 1	May 27	June 5	Sept. 5	Sept. 24
Dog Ridge.....	O	1904	1906	83	\$	4.4	2 to 4	Mar. 20	Mar. 30	May 25	June 2	May 29	June 12	Sept. 25	Oct. 4
Lenoir.....	O	1903	1905	84	\$	3.4	2 to 4	Mar. 19	May 24	June 5	May 28	June 17	Oct. 8	Oct. 8
Riparistis St. George.....	O	1903	1905	84	\$	2	2 to 4	Mar. 24	May 20	June 3	May 26	June 15	Sept. 27	Sept. 30
Nebbiolo Bourgt: Dog Ridge.....	O	1904	1906	93	\$	4.1	3 to 5	Mar. 17	May 22	June 5	May 28	June 19	Sept. 23	Oct. 8
Lenoir.....	O	1903	1905	88	\$	3.8	2 to 5	Mar. 14	May 19	May 30	May 23	June 14	Sept. 25	Oct. 8
Riparistis St. George.....	O	1903	1905	88	\$	4.1	2 to 4	Mar. 18	May 24	May 31	May 30	June 10	Sept. 24	Oct. 8
Taylor Narbonne.....	O	1904	1907	81	\$	1.6	4 to 5	Mar. 30	May 19	May 28	May 24	June 9	Sept. 23	Oct. 2
Nebbiolo F. no.: Dog Ridge.....	O	1904	1906	89	\$	4	3 to 5	Mar. 18	do	do	May 25	June 15	do	Oct. 4
Lenoir.....	O	1903	1905	86	\$	2.8	3 to 5	Mar. 16	do	do	May 23	June 13	do	Sept. 28
Riparistis St. George.....	O	1903	1905	84	\$	2.8	2 to 4	Mar. 20	May 24	May 25	May 29	June 2	do	Sept. 30
Taylor Narbonne.....	O	1904	1906	78	\$	1.2	3 to 4	Mar. 21	May 19	May 30	May 24	June 10	do	do
Negrone di Gattinara: Own roots.....	Gi	1904			\$			Mar. 14	Apr. 1	May 22	do	May 27	June 5	Sept. 16	Sept. 22
Negro Amaro: Own roots.....	Gi	1904			\$			Mar. 20	Mar. 29	May 24	June 3	May 30	June 10	Aug. 27	Sept. 23
Nebrette di Castilla: Own roots.....	Gi	1904			c, s			Mar. 10	Mar. 30	May 22	May 28	May 27	June 3	Sept. 23	Sept. 25
Ocup di Boe: Own roots.....	Gi	1904			c, s			Mar. 17	Mar. 28	May 28	Sept. 15	Sept. 21
Obanex: Lenoir.....	O	1906	1909	93	\$	1.2		Mar. 18	June 5	June 14	June 9	June 20
Olivette Blanche: Adoba Blanc.....	F	1907	1907	98	c, s	7.7	2 to 5	Mar. 21	Apr. 2	May 19	May 23	May 29	June 20	Sept. 21	Oct. 10
Lenoir.....	O	1907	1907	88	\$	2.8	2 to 5	Mar. 20	May 20	June 8	May 26	June 20	Sept. 30	Oct. 10
Riparia X Grandes Feuilles.....	F	1907	1907	96	c, s	6.6	3 to 5	Mar. 12	Apr. 1	May 21	May 23	June 1	Aug. 25	Sept. 21
Riparia X Ruparistis, No. 101.....	F	1907	1907	97	c, s	6.2	3 to 5	Mar. 22	do	May 19	May 24	May 30	Aug. 26	Sept. 21

Olivetto Chaptal: Lenoir.....	1904 1907	83	s	2.1 8.1	2 to 5 3 to 5	Mar. 24 do.	Apr. 1	May 21 do.	June 10 22	May 26 29	June 17 20	Sept. 25 27	Oct. 25
Solanis Robusta.....	1907	81	s	2	3 to 6	Mar. 25	Apr. 5	May 23	June 10	May 27	June 25	Sept. 27	Oct. 20
Olivette Noir: Lenoir.....	1907	81	s	6.3	3 to 5	Mar. 23	do.	May 21	June 25	May 30	June 19	Sept. 15	Oct. 10
Mourvedre X Rupestris, No. 1202.....	1907	94	C, S	7.3	3 to 5	Mar. 23	do.	May 21	June 10	May 30	June 19	Sept. 17	Oct. 8
Riparia X Rupestris, No. 3300.....	1907	55	C, S	8.6	2 to 5	Mar. 25	do.	May 20	June 10	May 31	June 13	do.	Oct. 10
Solanis X Othello, No. 1613.....	1907	89	s	3.1	2 to 4	Mar. 19	do.	May 21	June 5	May 29	June 20	Sept. 27	Oct. 15
Olivette Rose: Lenoir.....	1907	87	s	2.2	3 to 5	Mar. 20	do.	May 19	June 5	May 24	June 17	Sept. 25	Oct. 15
Opimian: Lenoir.....	1907	87	s	2.2	3 to 5	Mar. 20	do.	May 19	June 5	May 24	June 17	Sept. 25	Oct. 15
Orleans: Own roots.....	1904	Gi	C, S	do.	do.	do.	Mar. 26	May 26	June 3	May 31	June 8	Aug. 29	Oct. 25
Paquadabito: Own roots.....	1904	Gi	s	5.6	1 to 4	Mar. 10	Mar. 27	May 25	June 8	do.	June 14	Sept. 20	Do.
Dog Ridge.....	1906	64	s	4.1	3 to 5	Mar. 16	do.	May 25	June 5	May 30	June 20	Sept. 28	Oct. 10
Lenoir.....	1905	94	S	5.8	3 to 5	Mar. 16	do.	May 25	June 5	June 1	June 19	Sept. 27	Oct. 8
Rupestris St. George.....	1905	83	s	2	2 to 4	Mar. 18	do.	May 19	June 1	do.	June 11	do.	Oct. 10
Taylor Narbonne.....	1906	83	s	2	2 to 4	Mar. 18	do.	May 19	June 1	May 25	June 13	do.	Oct. 10
Talarus: Own roots.....	1904	Gi	s	10.1	2 to 4	Mar. 26	Mar. 28	May 26	June 10	May 31	June 15	Sept. 20	Sept. 30
Panariti: Adobe Giant.....	1904	94	C, S	6.5	2 to 4	Mar. 7	do.	May 14	May 19	May 24	June 1	Aug. 3	Aug. 1
Aramon X Rupestris Ganzin, No. 1.....	1904	85	C, S	3.2	2 to 4	Mar. 10	Apr. 1	May 13	May 22	do.	June 27	July 28	Do.
Dog Ridge.....	1903	91	s	1.7	2 to 5	Mar. 7	Mar. 27	May 12	May 18	May 26	May 28	July 25	Do.
Herbmont.....	1903	76	s	1.7	2 to 5	Mar. 8	do.	May 12	May 18	May 29	do.	do.	Do.
Lenoir.....	1903	90	C, S	5.5	3 to 5	Mar. 8	Mar. 25	May 11	May 20	May 22	May 24	July 25	Aug. 8
Mourvedre X Rupestris, No. 1202.....	1904	94	S	5.5	2 to 5	Mar. 5	Apr. 1	May 10	May 18	May 25	May 25	do.	Aug. 3
Riparia Gloire.....	1903	94	C, S	1.2	2 to 4	do.	do.	May 12	May 20	May 25	May 25	do.	Aug. 1
Riparia X Rupestris, No. 101.....	1904	73	S	5.6	2 to 4	Mar. 8	Mar. 27	May 12	May 20	May 25	May 25	July 25	Aug. 1
Riparia X Rupestris, No. 3300.....	1904	94	C, S	1.9	3 to 6	Mar. 12	do.	May 13	May 16	May 25	May 27	Do.	Do.
Riparia X Rupestris, No. 3309.....	1903	86	S	6.1	3 to 6	Mar. 11	do.	May 12	May 18	May 23	May 30	Do.	Do.
Rupestris Marini.....	1903	96	C, S	5.1	2 to 5	Mar. 9	do.	May 12	May 18	May 23	May 27	do.	Do.
Rupestris St. George.....	1903	90	C, S	4.8	2 to 4	Mar. 7	do.	May 12	May 20	May 25	May 27	do.	Do.
Salt Creek.....	1906	50	C, S	3.4	2 to 4	Mar. 7	do.	May 12	May 20	May 25	May 27	do.	Do.
Solanis X Riparia, No. 1616.....	1904	92	C, S	3.4	2 to 4	do.	Mar. 28	May 13	May 20	May 25	May 28	July 28	Do.
Taylor Narbonne.....	1906	92	C, S	5.5	2 to 5	Mar. 14	Mar. 25	May 12	May 23	May 22	do.	Aug. 20	Sept. 21
Tare de Versalles: Adobe Giant.....	1904	93	C, S	4.1	2 to 5	Mar. 6	do.	May 11	May 16	May 22	May 27	Aug. 5	Sept. 20
Aramon X Rupestris Ganzin, No. 1.....	1906	91	C, S	7.3	2 to 5	Mar. 6	do.	May 11	May 16	May 24	May 27	Aug. 5	Sept. 20
Dog Ridge.....	1904	95	S	3.2	2 to 5	Mar. 7	Mar. 30	May 12	May 25	May 24	May 30	do.	Do.
Lenoir.....	1903	90	C, S	1.8	3 to 5	Mar. 4	Mar. 15	May 15	May 20	May 22	May 25	Aug. 25	Sept. 16
Mourvedre X Rupestris, No. 1202.....	1906	81	S	10.5	3 to 5	Mar. 5	do.	May 21	May 22	June 1	June 5	do.	Sept. 20
Pouray.....	1907	93	C, S	6.7	3 to 5	Mar. 21	do.	May 21	May 20	May 20	May 26	Aug. 5	Sept. 19
Riparia Gloire.....	1903	91	C, S	5.4	2 to 4	Mar. 20	Mar. 20	May 21	May 20	May 20	May 26	Aug. 5	Sept. 19
Riparia X Rupestris, No. 101.....	1905	84	S	4.4	3 to 5	Mar. 20	do.	May 21	May 15	May 24	May 25	Aug. 10	Sept. 20
Riparia X Rupestris, No. 3300.....	1904	94	C, S	5	2 to 5	Mar. 8	Mar. 28	May 12	May 15	May 27	May 30	Oct. 17	Sept. 19
Rupestris Merlot.....	1907	92	S	7.6	3 to 6	Mar. 20	Mar. 20	May 21	May 17	May 20	May 23	Aug. 10	Sept. 19
Rupestris St. George.....	1903	96	C, S	2.8	4 to 5	Mar. 6	Mar. 25	May 11	May 20	May 25	May 28	Aug. 25	Sept. 20
Solanis X Riparia, No. 1616.....	1904	84	S	2.8	4 to 5	Mar. 6	do.	May 11	May 20	May 25	May 28	Aug. 25	Sept. 20
Taylor Narbonne.....	1904	96	C, S	3 to 6	3 to 6	Mar. 5	do.	May 5	May 23	May 24	do.	Aug. 5	Do.

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California.—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blooming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea- son.	Late sea- son.	Early sea- son.	Late sea- son.	Early sea- son.	Late sea- son.	Early sea- son.	Late sea- son.
1	1	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Pedro Ximines: Own roots.....	Gi	1904	P. cl.	c, s	Lbs.	Mar. 12	Mar. 28	May 32	May 30	May 29	June 6	Aug. 29	Sept. 25
Adon 6000.....	F	1904	1906	84	c, s	210 4	Mar. 17	Mar. 26	May 13	May 20	May 24	May 26	Sept. 15	Sept. 21
Aron 6000.....	F	1904	1906	88	c, s	2.7	310 5	Mar. 5	Mar. 30	May 20	May 28	May 25	June 4	Sept. 8	Sept. 15
Dor Ridge.....	F	1903	1905	90	s	2.4	310 5	Mar. 13	Mar. 26	May 14	May 22	May 27	May 30	Sept. 20	Sept. 20
Herbmont.....	F	1903	1905	72	s	1.1	310 5	Mar. 25	May 15	May 20	May 24	June 1	Sept. 13	Do.
Lenoir.....	F	1903	1905	88	s	1.9	210 5	Mar. 27	May 22	May 26	Sept. 15	Sept. 19
Monvodie X Rupestris, No. 1202.....	F	1904	1906	94	c, s	3.7	310 6	Mar. 26	May 14	May 24	May 26	Sept. 10	Sept. 20
Riparia Gloire.....	F	1903	1905	89	c, s	3	210 5	Mar. 20	May 15	May 21	June 5	Sept. 19
Riparia X Rupestris, No. 101.....	F	1904	1906	94	c, s	3.8	410 7	Mar. 8	Mar. 26	May 14	May 20	May 22	May 26	Sept. 19	Sept. 20
Riparia X Rupestris, No. 3309.....	F	1904	1906	95	c, s	3.6	310 5	Mar. 12	Mar. 26	May 14	May 20	May 22	May 26	Sept. 20	Sept. 20
Rupestris Martin.....	F	1903	1905	93	c, s	3.5	210 5	Mar. 11	Mar. 25	May 11	May 25	Do.	Do.
Rupestris St. George.....	F	1903	1905	83	c, s	3.2	410 6	Mar. 28	May 15	May 23	May 25	Sept. 16	Sept. 16
Salt Creek.....	F	1903	1905	81	c, s	1.3	310 5	Mar. 30	May 12	May 16	May 21	May 30	Sept. 8	Sept. 19
Soleais X Riparia, No. 1616.....	F	1904	1906	92	c, s	2.9	310 5	Mar. 6	Mar. 20	May 15	May 23	May 28	May 28	Sept. 20	Sept. 20
Taylor Narbonne.....	F	1904	1906	90	c, s	2.2	310 5	Mar. 11	Mar. 25	May 14	May 25	Do.
Ferle Blanche: Lenoir.....	O	1903	1905	89	s	3.4	310 6	Mar. 25	May 14	May 25	May 27	Sept. 15	Do.
Rupestris St. George.....	O	1903	1905	91	s	4.4	210 4	Mar. 19	May 23	June 4	May 24	June 18	Sept. 23	Sept. 28
Lenoir.....	O	1904	1907	80	s	1.2	210 6	Mar. 21	May 18	June 3	May 27	June 15	Oct. 15
Ferruno: Oven roots.....	Gi	1904	s	Mar. 14	Mar. 29	May 24	June 4	May 29	June 10	Sept. 20	Sept. 23
Persian, No. 21: Oven roots.....	Gi	1904	c, s	Mar. 18	Mar. 30	May 30	Sept. 20
Persian, No. 25: Oven roots.....	Gi	1904	c, s	Mar. 16	Mar. 25	Sept. 24
Persian, No. 26: Oven roots.....	Gi	1904	c, s	Mar. 20	Mar. 28	June 3	June 5	June 11
Peru: Lenoir.....	O	1903	1909	92	s	3.2	310 4	Mar. 26	June 9	June 10	June 13	June 27	Sept. 25	Oct. 8

Petit Syrah: O. roots.	GI	1907 1904	1909	83	S	8	1, 2	Mar. 7	Apr. 1 Mar. 29	May 20 May 23	June 1	June 10	Sept. 20	Do.
Adobe Giant (Aestivalis X Monticola) X (Riparia X Rupestris, No. 364-5).	G	1904	1909	90	S	4, 2	Mar. 9	Apr. 1	May 25	June 1	June 10	Sept. 23	Do.	
Aramon X Rupestris Ganzin, No. 1.	G	1904	1909	85	S	3, 1	Mar. 18	Apr. 2	May 28	June 12	Sept. 24	Do.		
Do.	G	1904	1909	95	S	3, 5	Mar. 2	Mar. 30	May 20	June 12	Sept. 24	Oct. 4		
Aramon X Rupestris Ganzin, No. 9.	G	1904	1909	60	S	1, 5	Mar. 2	Mar. 2	June 15	June 15	Sept. 7	Oct. 8		
Australis	G	1904	1909	91	S	1, 1	Mar. 10	Apr. 1	May 27	June 18	July 2	Oct. 22		
Berlandieri, No. 1.	G	1904	1909	78	S	1, 5	Mar. 6	do.	June 3	June 3	Oct. 26	Do.		
Berlandieri, No. 2.	G	1904	1909	85	S	1	Mar. 10	Mar. 27	June 1	June 1	Sept. 28	Do.		
Berlandieri X Labott, No. 9	G	1904	1909	91	S	5	Mar. 5	Mar. 30	do.	do.	do.	Do.		
Berlandieri X Riparia, No. 34EM.	G	1904	1909	91	S	3, 5	Mar. 2	Apr. 1	May 28	May 28	Oct. 1	Do.		
Berlandieri X Riparia, No. 15-11.	G	1904	1909	82	S	1, 8	Mar. 4	Mar. 27	do.	do.	do.	Do.		
Berlandieri X Riparia, No. 420A.	G	1904	1909	72	S	5	Mar. 6	Mar. 28	do.	do.	do.	Do.		
Do.	G	1907	1909	76	S	1, 1	Mar. 21	Mar. 27	June 1	June 4	June 8	Sept. 20	Do.	
Bourisquon X Rupestris, No. 106-4.	G	1904	1909	88	S	3	Mar. 5	Apr. 1	May 27	June 3	June 13	Oct. 12		
(Bourisquon X Rupestris, No. 601) X Calicola, No. 12205.	G	1907	1909	86	S	1, 7	do.	Apr. 3	May 30	May 30	Sept. 26	Oct. 7		
Bourisquon X Rupestris, No. 3807.	G	1904	1909	97	S	3, 5	Mar. 7	Apr. 1	June 2	June 2	Sept. 20	Do.		
Cabernet X Berlandieri, No. 333.	G	1904	1909	90	S	4, 7	Mar. 8	Mar. 28	June 1	June 4	Sept. 26	Oct. 8		
Cabernet X Rupestris Ganzin, No. 33A.	G	1904	1909	90	S	2, 2	Mar. 6	Apr. 3	May 23	June 1	June 9	Oct. 10		
Captain	G	1908	1909	86	S	2	Mar. 3	Apr. 1	June 1	June 15	June 25	Sept. 26		
Chasselas X Berlandieri, No. 41B.	G	1904	1909	91	S	2	Mar. 2	Apr. 2	June 1	June 20	June 20	Oct. 7		
Constitution.	O	1907	1909	87	S	2, 3	Mar. 20	do.	do.	do.	do.	do.		
Do.	G	1904	1909	93	S	3	Mar. 6	Mar. 30	June 1	June 5	June 12	Sept. 26		
Dog Ridge.	G	1904	1909	88	S	2, 9	Mar. 19	do.	May 19	June 5	June 12	Sept. 27		
Do.	G	1903	1905	86	S	2, 8	Mar. 17	Mar. 10	May 25	June 4	June 17	Sept. 25		
Herbennont	G	1904	1909	93	S	2, 2	Mar. 3	Mar. 10	May 20	June 4	June 12	Sept. 20		
Hotopurp	G	1904	1909	80	S	1, 1	Mar. 4	Apr. 3	June 1	June 1	do.	do.		
July	G	1907	1909	70	S	4	Mar. 4	Apr. 1	do.	do.	do.	Do.		
Judge.	G	1907	1909	85	S	1, 5	Mar. 20	Apr. 1	May 23	June 5	June 11	Sept. 25		
Lenoir	G	1903	1905	89	S	2, 9	Mar. 6	Apr. 1	May 25	June 3	June 10	do.		
Monticola X Riparia, No. 554.	O	1904	1909	85	S	1, 5	Mar. 6	Apr. 1	June 2	June 6	June 17	Oct. 10		
Monticola X Riparia, No. 18908.	O	1907	1909	87	S	1, 2	Mar. 22	Mar. 10	June 2	do.	do.	do.		
Monticola X Rupestris.	G	1906	1909	85	S	1, 2	Mar. 10	Apr. 1	May 30	June 6	June 17	Oct. 10		
Motley.	G	1904	1909	86	S	3, 4	Mar. 8	do.	May 23	June 1	June 12	do.		
Mourvudre X Rupestris, No. 1202.	G	1904	1909	95	S	3, 2	Mar. 6	do.	do.	do.	do.	do.		
Do.	G	1904	1909	92	S	2	Mar. 21	do.	June 12	June 15	June 25	Oct. 7		
Mourvudre X Rupestris, No. 1203.	G	1905	1909	78	S	6	Mar. 4	do.	June 1	June 18	June 25	Sept. 5		
Pardes.	G	1907	1909	90	S	2, 2	Mar. 6	Mar. 20	May 25	May 30	June 9	Sept. 20		
Riparia du Colorado.	G	1904	1909	86	S	1	Mar. 10	do.	May 28	do.	do.	Sept. 26		
Riparia X (Cordifolia X Rupestris) No. 106-S.	G	1904	1909	89	S	3, 5	do.	Apr. 30	do.	do.	do.	do.		
Riparia X Rupestris, No. 101.	G	1904	1909	94	S	2, 5	Mar. 6	Apr. 2	May 30	June 7	June 20	Sept. 25		
Do.	O	1904	1906	85	S	2	Mar. 21	do.	May 20	June 1	June 25	Oct. 6		
Riparia X Rupestris, No. 101-H.	G	1904	1909	90	S	1, 7	Mar. 10	Mar. 30	June 1	June 16	June 26	Sept. 27		
Do.	O	1907	1909	72	S	5	Mar. 19	do.	June 2	June 10	June 25	Sept. 26		

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment yard.	Year planted was	Year grafted.	Congenality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Petit Syrah—Continued.															
Riparia × Rupesstris, No. 108-103.	G	1905	1909	81	8	1.2	2 to 3	Mar. 2	Apr. 3	June 5	June 2	June 6	June 15	Sept. 26	Oct. 8
Riparia × Rupesstris, No. 359b.	G	1901	1909	98	8	3.2	2 to 1	Mar. 9	Mar. 27	May 27	June 1	June 2	June 15	Sept. 25	Oct. 7
Do.	G	1907	1909	88	8	1.2	2 to 1	Mar. 19	Apr. 3	May 30	June 1	June 7	June 22	Sept. 26	Oct. 11
Riparia × Rupesstris, No. 3809.	G	1901	1909	93	8	3.7	2 to 5	Mar. 17	Apr. 17	May 21	June 1	June 9	June 22	Sept. 25	Oct. 8
Do.	G	1901	1906	91	8	2.7	2 to 5	Mar. 8	Mar. 27	May 28	June 1	June 9	June 26	Sept. 28	Oct. 8
Rupesstris des Causses.	G	1904	1909	89	8	2.7	2 to 5	Mar. 11	Mar. 27	June 1	June 3	May 29	June 20	Sept. 24	Oct. 13
Rupesstris Martin.	G	1903	1909	89	8	3.2	2 to 5	Mar. 18	Apr. 3	May 21	June 3	May 29	June 20	Sept. 24	Oct. 13
Do.	G	1903	1909	91	8	1.2	2 to 3	Mar. 6	Apr. 3	May 29	June 5	June 9	June 2	Sept. 25	Oct. 8
Do.	G	1904	1909	84	8	1.2	2 to 3	Mar. 20	Mar. 27	June 3	June 5	June 9	June 2	Sept. 25	Oct. 8
Rupesstris Mission.	G	1901	1909	86	8	1.2	2 to 3	Mar. 9	Mar. 27	May 27	June 1	June 9	June 2	Sept. 28	Oct. 8
Do.	G	1907	1909	88	8	1.2	2 to 3	Mar. 9	Apr. 1	May 29	June 1	June 9	June 2	Sept. 28	Oct. 8
Rupesstris St. George.	G	1903	1909	88	8	3.2	2 to 5	Mar. 1	Mar. 30	June 4	June 10	June 17	June 17	Sept. 26	Oct. 4
Do.	G	1903	1905	90	8	3.2	2 to 5	Mar. 29	Apr. 3	May 21	June 1	June 14	June 17	Sept. 27	Oct. 8
Rupesstris × Peckhamieri, No. 301A.	G	1901	1909	82	8	1.6	2 to 3	Mar. 8	Apr. 3	June 1	June 1	June 12	June 25	Sept. 25	Oct. 10
Rupesstris × Peckhamieri, No. 301B.	G	1901	1909	87	8	2.8	2 to 3	Mar. 6	Apr. 3	June 30	June 1	June 12	June 25	Sept. 25	Oct. 10
Rupesstris × Peckhamieri, No. 301D.	G	1901	1909	87	8	2	2 to 3	Mar. 6	Apr. 3	June 1	June 1	June 12	June 25	Sept. 25	Oct. 10
Rupesstris × Peckhamieri, No. 301E.	G	1905	1909	83	8	1.3	2 to 3	Mar. 4	Apr. 3	June 1	June 1	June 12	June 25	Sept. 26	Oct. 8
Rupesstris × Hybrid Azemar, No. 215.	G	1901	1909	81	8	1.6	2 to 3	Mar. 10	Mar. 18	May 26	June 1	June 12	June 20	Sept. 20	Oct. 8
Salt Creek.	G	1901	1909	82	8	1.4	2 to 5	Mar. 9	Apr. 1	May 27	June 1	June 12	June 6	Sept. 25	Oct. 11
Do.	G	1903	1905	88	8	1.5	2 to 5	Mar. 7	Apr. 5	May 22	June 6	June 20	June 20	Sept. 20	Oct. 8
Solonis Ordinaire.	G	1904	1909	70	8	1.7	3 to 4	Mar. 8	Apr. 1	May 28	June 10	June 15	June 20	Sept. 20	Oct. 7
Solonis Robusta.	G	1907	1909	91	8	1.7	3 to 4	Mar. 19	Apr. 1	May 28	June 9	June 9	June 20	Sept. 25	Oct. 7
Do.	G	1901	1909	91	8	1.6	3 to 4	Mar. 2	Apr. 1	May 28	June 3	June 7	June 15	Sept. 26	Oct. 8
Solonis × Othello.	G	1904	1909	77	8	1.2	2 to 1	Mar. 21	Mar. 30	May 27	June 10	June 15	June 15	Sept. 28	Oct. 7
Solonis × Othello, No. 1613.	G	1907	1909	82	8	1.2	2 to 1	Mar. 6	Mar. 30	May 27	June 10	June 10	June 17	Sept. 28	Oct. 9
Solonis × Riparia, No. 1616.	G	1904	1906	83	8	1.4	2 to 5	Mar. 17	Apr. 2	May 28	June 10	June 1	June 17	Sept. 20	Oct. 5
Do.	G	1904	1909	83	8	2.5	2 to 4	Mar. 1	Apr. 2	May 28	June 8	June 8	June 19	Sept. 23	Oct. 9
Taylor Narbonne.	G	1901	1909	83	8	1.9	2 to 4	Mar. 23	Apr. 2	May 25	June 8	May 30	June 19	Sept. 26	Oct. 8
Do.	G	1904	1906	88	8	1.9	2 to 4	Mar. 23	Apr. 2	May 25	June 8	May 30	June 19	Sept. 26	Oct. 8

TABLE VII.—Relative behavior and value for different nurseries of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California (Continued).

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Pis de Chevre des Alpes:															
Own roots.....	G	1904			\$	3.1	2 to 4	Mar. 15	Mar. 29	May 25	June 2	May 31	June 9	Sept. 24	Oct. 10
Dog Ridge.....	O	1904	1906	86	\$	2.9	3 to 5	Mar. 23	Mar. 29	May 19	May 31	May 23	June 12	Sept. 23	Oct. 9
Lenoir.....	O	1903	1905	90	\$	4.9	2 to 3	Mar. 21	do.	June 8	May 24	June 24	do.	Oct. 12
Rupestris St. George.....	O	1903	1905	92	\$	4.9	2 to 3	Mar. 22	May 24	May 31	May 3	June 12	Sept. 22	Oct. 12
Pis de Chevre Rouge:															
Lenoir.....	O	1904	1907	86	\$	3.2	3 to 5	Mar. 20	May 23	June 6	May 28	June 20	Sept. 20	Sept. 21
Pizzutella:															
Own roots.....	G	1904			c, s	4	3 to 5	Mar. 20	Mar. 28	May 30	June 1	June 5	Sept. 10	Sept. 23
Adobe Giant.....	F	1904	1906	89	c, s	4	3 to 5	Mar. 25	Apr. 3	May 20	May 30	May 27	June 7	do.	Sept. 20
Aramon X Rupestris Ganzin, No. 1.....	F	1904	1906	95	c, s	6.6	2 to 4	Mar. 12	Apr. 1	do.	May 29	May 26	June 3	Sept. 12	Do.
Dog Ridge.....	F	1903	1905	80	\$	1.1	2 to 5	Mar. 8	Mar. 30	May 15	May 22	May 27	May 28	Sept. 16	Do.
Herbemont.....	F	1903	1905	73	\$	1.1	3 to 5	Mar. 6	Mar. 25	do.	May 27	May 30	June 1	Sept. 9	Do.
Lenoir.....	F	1903	1905	94	\$	2	2 to 5	do.	Mar. 30	do.	May 20	May 26	May 27	Sept. 10	Sept. 15
Mourvedre X Rupestris, No. 1202.....	F	1904	1906	87	c, s	4.9	3 to 5	Mar. 14	Apr. 1	May 12	May 29	do.	June 4	Sept. 5	Sept. 19
Riparia X Rupestris, No. 101.....	F	1903	1905	96	c, s	7.3	2 to 5	Mar. 20	do.	May 18	May 28	May 23	June 3	Sept. 8	Sept. 20
Riparia X Rupestris, No. 3309.....	F	1904	1906	98	c, s	13.3	3 to 5	Mar. 13	Apr. 2	May 14	May 25	May 27	June 5	Sept. 10	Sept. 20
Rupestris Martin.....	F	1903	1905	95	c, s	5.1	3 to 5	Mar. 14	Apr. 6	May 16	May 25	May 25	June 8	Sept. 19	Do.
Rupestris St. George.....	F	1903	1905	82	c, s	6.5	2 to 5	Mar. 15	Apr. 1	do.	May 21	May 25	June 6	Sept. 10	Do.
Salt Creek.....	F	1903	1905	86	c, s	2.5	3 to 6	Mar. 14	Apr. 30	May 21	May 29	May 28	June 6	Sept. 5	Do.
Solonis X Riparia, No. 1616.....	F	1904	1906	90	c, s	5.6	2 to 5	Mar. 24	Apr. 1	May 20	May 29	May 23	June 6	Sept. 10	Do.
Taylor Narbonne.....	F	1904	1906	87	c, s	3.7	3 to 5	Mar. 14	Mar. 31	May 15	May 24	May 26	May 28	Sept. 8	Do.
Poulsard:															
Own roots.....	G	1904			\$			Mar. 17	Mar. 28	May 27	June 5	Sept. 15	Sept. 21
Precoce de Courtillet:															
Lenoir.....	O	1903	1905	79	\$	2.4	2 to 4	Mar. 18	May 22	May 30	May 26	June 14	Sept. 9	Sept. 25
Rupestris St. George.....	O	1903	1907	82	\$	3.4	2 to 4	Mar. 22	do.	May 31	May 29	June 10	Sept. 22	Do.
Prune de Cazouls:															
Aramon X Rupestris Ganzin, No. 2.....	F	1907	1909	88	\$			Mar. 26	May 21	Sept. 29	Oct. 5
Austrais.....	F	1907	1909	90	\$			Mar. 31	May 25	Sept. 5	Oct. 5
Berlandieri X Riparia, No. 420A.....	F	1907	1909	87	\$			Mar. 23	May 21	Sept. 15	Oct. 5

O	1904	91	S	2 S	2 to 5	Mar. 22	May 23	June 8	May 30	June 20	Sept. 23	Oct. 10
O	1907	91	S	Mar. 22	May 23	June 8	May 30	June 20	Sept. 23	Oct. 10
F	1907	90	S	Apr. 3	May 23	June 8	May 30	June 20	Sept. 23	Oct. 10
F	1907	90	S	Mar. 20	May 18	June 8	May 30	June 20	Sept. 23	Oct. 10
F	1907	90	S	Mar. 20	May 18	June 8	May 30	June 20	Sept. 23	Oct. 10
F	1907	87	S	Mar. 26	May 23	June 8	May 30	June 20	Sept. 23	Oct. 10
F	1907	87	S	Apr. 5	May 23	June 8	May 30	June 20	Sept. 23	Oct. 10
F	1907	89	S	Mar. 22	May 23	June 8	May 30	June 20	Sept. 23	Oct. 10
Purple Damascus:													
Gi	1905	C, S	Mar. 26	May 30	June 6	June 13	Sept. 10	Sept. 24
F	1904	86	S	Mar. 12	May 11	June 23	May 23	Sept. 8	Sept. 21
F	1904	85	S	Mar. 27	May 11	June 23	May 23	Sept. 11	Sept. 20
F	1904	92	S	Apr. 1	May 17	June 23	May 26	Sept. 8	Do.
F	1903	92	S	Apr. 7	May 16	June 23	May 30	Sept. 22	Do.
F	1904	95	S	Mar. 7	May 15	June 23	May 28	Sept. 8	Do.
F	1903	89	S	Mar. 7	May 15	June 23	May 28	Sept. 19	Do.
F	1904	88	S	Mar. 30	May 8	June 24	May 20	Sept. 5	Do.
F	1904	93	S	Apr. 3	May 24	June 24	May 26	Sept. 12	Do.
F	1904	95	S	Mar. 12	May 16	June 24	May 29	Sept. 10	Do.
F	1903	87	S	Mar. 15	May 16	June 24	May 29	Sept. 10	Do.
F	1904	92	S	Mar. 14	May 14	June 21	May 29	Sept. 12	Do.
F	1904	92	S	Mar. 14	May 14	June 21	May 29	Sept. 12	Do.
Quangine:													
Gi	1904	S	Mar. 19	May 26	June 6	June 30	Sept. 19	Sept. 25
O	1901	82	S	Apr. 1	May 26	June 6	June 30	Sept. 19	Sept. 25
O	1903	90	S	May 23	June 8	June 2	Sept. 29	Oct. 6
O	1903	91	S	May 23	June 8	June 2	Sept. 28	Oct. 8
O	1904	89	S	Mar. 20	May 21	June 5	May 27	Oct. 13
O	1904	89	S	Mar. 19	May 23	June 5	May 29	Do.
Razaki Zolot:													
Gi	1905	89	C, S	Mar. 26	May 26	June 6	June 26	Sept. 15	Sept. 23
F	1903	90	C, S	Mar. 21	May 10	June 8	May 19	Sept. 23	Oct. 1
O	1904	90	C, S	Mar. 21	May 10	June 8	May 19	Sept. 23	Oct. 1
Refresco:													
Gi	1904	S	Mar. 20	May 29	June 6	June 6
O	1901	91	C, S	Mar. 24	May 23	June 8	June 8	Sept. 21	Oct. 8
O	1903	90	C, S	Mar. 16	May 23	June 8	May 29	Sept. 21	Oct. 15
O	1903	92	S	Mar. 19	May 25	June 8	June 2	Sept. 23	Oct. 8
O	1904	88	S	Mar. 22	May 27	June 31	June 2
Ribier:													
O	1903	92	S	Mar. 15	May 23	June 1	May 26	Sept. 28	Sept. 23
O	1903	93	S	Mar. 23	May 22	June 3	May 28	Sept. 29	Sept. 23
Robin Noir:													
Gi	1904	S	Mar. 16	May 26	June 30	May 31
Rodites:													
O	1904	1907	S	Mar. 21	May 25	June 8	May 30
Rese d'Italie:													
O	1903	84	S	Mar. 17	May 22	June 5	June 28
O	1904	83	S	Mar. 24	June 1	June 3	May 28
O	1903	90	C, S	Mar. 16	May 23	June 31	June 6	Sept. 26	Sept. 30
O	1904	92	S	Mar. 21	June 1	May 31	June 6	Sept. 29	Oct. 8
O	1904	84	S	Mar. 16	June 2	June 3	June 7	Sept. 15	Oct. 10
O	1904	84	S	Mar. 16	June 2	June 3	June 7	Sept. 15	Oct. 10

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Rothschilder:															
Own roots.	GI	1904		P.ct.	C, S	Lbs.		Mar. 15	Mar. 28	May 24	June 1	May 29	June 7	Aug. 24	Oct. 20
Roussseau:															
Own roots.	GI	1905			C, S			Mar. 18	Mar. 30	May 25	June 6	do.	June 12	Aug. 26	Sept. 23
Dog Ridge.	O	1904	1906	91	S	4.2	3 to 5	Mar. 16	do.	do.	Mar. 30	do.	June 15	Sept. 25	Oct. 10
Léonor.	O	1903	1905	87	S	5.2	2 to 4	do.	do.	May 27	May 31	June 2	June 16	Sept. 20	Oct. 9
Riparia X Rupestris, No. 3306.	O	1904	1905	91	C, S	4.7	1 to 3	Mar. 20	Mar. 19	May 15	June 5	June 5	June 18	Sept. 23	Oct. 5
Riparia X Rupestris de Jaeger.	F	1907	1907	91	C, S	3.8		Mar. 19	Mar. 16	May 22	June 5	May 27	June 17	Aug. 25	Oct. 17
Rupestris St. George.	O	1903	1905	90	S	2.4	2 to 4	Mar. 16	Mar. 19	May 16	June 5	May 27	June 17	Sept. 15	Sept. 26
Solonis X Riparia, No. 1016.	F	1907	1907	84	S	2.2		Mar. 19	Mar. 21	May 21	June 1	May 24	June 10	Oct. 17	Sept. 26
Taylor Narbonne.	O	1904	1906	80	S	1.5	2 to 4	Mar. 21	Mar. 26	May 19	June 1	May 24	June 10	Sept. 25	Do.
Royal Ascot:															
Own roots.	GI	1905			S			Mar. 15	Mar. 26	May 20	do.	May 26	June 6	Aug. 27	Do.
Saint Laurent:															
Own roots.	GI	1905			S			Mar. 18	Apr. 1	May 26	May 30	May 30	June 5	Aug. 29	Sept. 23
Adobe Giant.	F	1904	1906	83	S	2.4	3 to 5	Mar. 14	do.	May 15	May 18	May 23	May 28	Aug. 10	Sept. 6
Aramon X Rupestris Ganzin No. 1.	F	1904	1906	90	C, S	3.7	2 to 6	Mar. 14	Apr. 2	May 16	May 19	May 22	May 30	Aug. 5	Sept. 10
Dog Ridge.	F	1903	1905	81	S	2.6	2 to 5	Mar. 15	Apr. 1	May 11	May 22	May 24	May 31	Aug. 1	Sept. 6
Herbmont.	F	1903	1905	74	S	3.7	2 to 5	Mar. 14	do.	May 10	May 25	May 25	June 1	do.	Sept. 10
Léonor.	F	1903	1905	67	S	3.8	2 to 6	Mar. 12	Mar. 30	May 12	May 20	May 20	May 27	Aug. 24	Sept. 11
Mourvedre X Rupestris, No. 1202.	F	1904	1906	89	C, S	3.3	2 to 6	Mar. 15	do.	May 15	do.	May 25	May 27	Aug. 8	Sept. 8
Riparia Cloire.	F	1903	1905	82	S	2.1	2 to 5	Mar. 18	Apr. 1	May 14	do.	May 23	May 26	Aug. 3	Do.
Riparia X Rupestris, No. 101.	F	1904	1906	86	S	2.1	2 to 5	Mar. 14	Mar. 30	May 7	do.	May 23	May 27	Aug. 10	Sept. 10
Riparia X Rupestris, No. 3309.	F	1904	1906	83	S	1.5	2 to 5	Mar. 13	Apr. 5	May 14	May 17	do.	May 29	do.	Sept. 8
Rupestris Martin.	F	1903	1905	69	S	1.5	2 to 5	Mar. 24	Apr. 1	do.	do.	do.	May 31	Aug. 5	Do.
Rupestris St. George.	F	1903	1905	91	C, S	2.4	2 to 6	Mar. 15	Mar. 27	May 13	May 20	do.	May 30	Aug. 1	Do.
Salt Creek.	F	1903	1905	75	S	1.3	2 to 6	do.	do.	May 15	May 17	do.	May 29	Sept. 9	Do.
Solonis X Riparia, No. 1016.	F	1904	1906	72	S	1.1	2 to 6	Mar. 6	Mar. 19	May 14	May 17	May 22	May 29	Sept. 9	Do.
Taylor Narbonne.	F	1904	1906	50	S	1.3		Mar. 14	Mar. 31	May 16	do.	May 21	May 26	Aug. 24	Do.
Saint Macaire:															
Own roots.	GI	1904	1909	94	S	3.7	3 to 5	Mar. 12	Mar. 31	May 24	June 2	May 28	June 7	Sept. 18	Sept. 25
Aramon X Rupestris Ganzin, No. 1.	O	1904	1906	91	S	3.2	2 to 4	Mar. 19	do.	May 25	June 5	May 30	June 20	Sept. 28	Oct. 8
Dog Ridge.	O	1904	1906	91	S			Mar. 19	do.	May 25	June 5	May 30	June 20	Sept. 28	Oct. 8

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	Low pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Sanmillon—Continued.															
Lenoir.....	0	1903	1905	P. et.	C, S	2 to 6	8	9	10	11	12	13	14	15	16
Monticola X Riparia, No. 18848.....	0	1907	1909	85	S	3 to 8	2 to 4	Mar. 23	May 20	June 5	May 26	June 18	Sept. 23	Oct. 8
Meurcedra X Rupestris, No. 1247.....	0	1904	1905	85	S	3 to 4	2 to 4	Mar. 16	May 20	June 3	May 29	June 17	Sept. 25	Oct. 6
Riparia X Rupestris, No. 101.....	0	1904	1905	89	S	3 to 4	3 to 4	Mar. 21	May 23	June 2	May 29	June 18	Sept. 24	Oct. 11
Riparia X Rupestris, No. 33405.....	0	1907	1909	91	C, S	1 to 4	3 to 4	Mar. 20	June 2	June 8	June 6do.	Sept. 20	Do.
Riparia X Rupestris, No. 33309.....	0	1904	1905	89	C, S	2 to 4	3 to 4	May 23	June 2	May 29do.	Oct. 5
Rupestris Marlin.....	0	1903	1905	88	C, S	4 to 7	2 to 4	Mar. 20	do.	June 4	May 28	June 20	Sept. 24	Oct. 5
Rupestris St. George.....	0	1903	1905	86	S	1 to 9	3 to 4	Mar. 16	May 26	June 1	May 29	June 11	Sept. 22	Oct. 8
Salt Creek.....	0	1903	1905	89	S	2 to 1	2 to 4	Mar. 19	May 23	June 10	May 29	June 24	Sept. 20	Oct. 11
Solanis X Othello, No. 1613.....	0	1907	1909	88	S	1 to 1	2 to 4	Mar. 20	May 25	June 3	June 8	June 11	Sept. 24	Oct. 4
Solanis X Riparia, No. 1616.....	0	1904	1905	88	C, S	2 to 7	2 to 4	Mar. 17	do.	June 5	May 30	June 21	Sept. 24	Oct. 8
Taylor Narbonne.....	0	1904	1905	90	C, S	2 to 9	2 to 4	Mar. 20	May 20do.	June 2	June 18do.	Oct. 4
Sanmillon Blanc:															
Own roots.....	0	1905	C, S	Mar. 19	Mar. 31	May 28	May 29	June 1	June 4	Aug. 29	Sept. 25
Sarnie:															
Own roots.....	0	1904	S	Mar. 10	Mar. 29	May 27	May 30	June 6	Sept. 20	Sept. 22
Dog Ridge.....	0	1904	1907	91	C, S	4 to 3	3 to 6	Mar. 18	May 25	June 1	May 30	June 21	Sept. 23	Oct. 10
Lenoir.....	0	1903	1905	89	S	3 to 7	3 to 5	Mar. 17	May 22	June 5	May 26	June 18	Sept. 20	Oct. 6
Rupestris St. George.....	0	1903	1905	91	S	5 to 5	3 to 5	Mar. 20	May 19	June 3do.	June 14	Sept. 15	Do.
Taylor Narbonne.....	0	1904	1905	86	S	3 to 6	2 to 4	Mar. 19	May 22	June 2	May 27do.	Oct. 10
Servan Rose:															
Lenoir.....	0	1904	1907	86	S	2 to 5	2 to 5	Mar. 18do.	June 7	May 28	June 20	Sept. 28	Oct. 13
Sticien:															
Own roots.....	0	1905	S	Mar. 26	Apr. 1	May 28	May 29	June 4	June 28	Aug. 28	Sept. 27
Aramon X Rupestris Ganzin, No. 1.....	0	1904	88	S	6 to 3	3 to 6	Mar. 18	June 1	June 5	June 5	June 10	Sept. 29	Oct. 15
Lenoir.....	0	1904	1907	87	S	2 to 9	3 to 6	Mar. 16	May 21	June 4	May 27	June 20	Sept. 9	Oct. 6
Slankamenka:															
Own roots.....	0	1904	S	Mar. 14	Apr. 3	May 29	May 30	June 5	Aug. 23	Sept. 22
Souvenir du Congrès:															
Dog Ridge.....	0	1904	1906	85	S	2 to 2	3 to 5	Mar. 20	May 12	June 1	June 3	June 17	Sept. 21	Oct. 1
Lenoir.....	0	1903	1905	93	S	6 to 2	3 to 4	Mar. 20	May 26	June 3	May 30do.	Sept. 23	Oct. 15
Rupestris St. George.....	0	1903	1905	89	S	3 to 7	2 to 4	Mar. 19do.do.do.	June 16	Sept. 20	Sept. 22

Spurbonna:	Gi	1904		S		Mar. 17	Apr. 1	May 27		June 2		June 15	Aug. 29	Sept. 24
Own roots.....	O	1904	3, 5	c, s	2 to 4	Mar. 21	May 19	June 2	June 5	June 8	June 12	June 19	Oct. 12	Sept. 24
Sucre de Marselle:	O	1904	1, 6	s	3 to 5	Mar. 16	May 22	June 3	June 5	June 8	June 12	June 19	Sept. 20	Oct. 8
Dog Ridge.....	O	1903	4, 4	s	2 to 5	Mar. 24	May 25	June 5	June 8	June 12	June 16	June 19	Sept. 15	Oct. 12
Lenoir.....	O	1903	1, 3	s	2 to 5	Mar. 18	May 19	June 2	June 5	June 8	June 12	June 19	Sept. 21	Oct. 12
Rupestris St. George.	O	1904	8	s	2 to 5	do.	May 6	June 8	June 12	June 16	June 19	June 25	do.	do.
Taylor Narbonne.....	O	1904				do.	May 6	June 8	June 12	June 16	June 19	June 25	do.	do.
Sutliffa:														
Own roots.....	Gi	1904		c, s		Mar. 14	May 29	June 6	June 13	June 2	June 6	June 13	Sept. 5	Sept. 5
Own roots.....	Gi	1904		c, s		Mar. 12	May 28	June 3	June 10	June 3	June 10	June 16	Sept. 16	Sept. 16
Do.....	O	1904	3, 2	s	2 to 4	Mar. 24	May 21	June 11	June 18	June 31	June 16	June 16	Sept. 24	Sept. 24
Aramon X Rupestris Ganzin, No. 1.....	O	1907	5, 3	s	3 to 5	Mar. 26	May 24	June 14	June 14	June 14	June 28	June 28	Oct. 3	Oct. 8
Do.....	F	1907	5, 3	c, s	3 to 6	Mar. 10	May 17	June 4	June 4	June 4	June 4	June 4	Aug. 30	Sept. 21
Aramon X Rupestris Ganzin, No. 2.....	F	1909	4, 5	s	4	Mar. 22	May 28	June 17	June 17	June 17	June 17	June 17	Oct. 16	Oct. 16
Australis.....	F	1909	4	s	2 to 6	Mar. 20	May 24	June 10	June 10	June 10	June 22	June 22	Sept. 16	Oct. 16
Dog Ridge.....	O	1904	2, 5	c, s	3 to 5	do.	May 25	June 25	June 25	June 25	June 25	June 25	Sept. 25	Sept. 25
Hooprup.....	F	1909	8, 2	s	3 to 5	Mar. 14	May 23	June 15	June 15	June 15	June 30	June 30	Oct. 16	Oct. 16
Lenoir.....	O	1904	2, 8	c, s	3 to 6	Mar. 8	May 23	June 15	June 15	June 15	June 30	June 30	Sept. 24	Sept. 30
Monticola X Riparia, No. 554.....	F	1907	5, 5	c, s	3 to 6	Mar. 12	May 18	June 23	June 23	June 23	June 5	June 5	Aug. 25	Oct. 17
Monticola X Riparia, No. 18804.....	F	1907	11	s	3 to 6	Mar. 16	May 22	June 5	June 5	June 5	June 5	June 5	Sept. 21	Sept. 21
Monticola X Riparia, No. 3306.....	F	1909	3	s	3 to 6	Mar. 17	May 25	June 26	June 26	June 26	June 26	June 26	Sept. 5	Sept. 5
Monticola X Riparia, No. 18808.....	F	1907	3, 5	s	3 to 6	Mar. 20	May 26	June 26	June 26	June 26	June 26	June 26	Aug. 26	Aug. 26
Monticola X Riparia, No. 18815.....	F	1907	3, 5	s	3 to 6	Mar. 20	May 26	June 26	June 26	June 26	June 26	June 26	Sept. 21	Sept. 21
Monticola X Rupestris.....	F	1909	3, 5	s	3 to 6	Mar. 12	May 21	June 26	June 26	June 26	June 26	June 26	Aug. 27	Aug. 27
Pomroy.....	F	1907	3, 8	c, s	3 to 6	Mar. 19	May 18	June 23	June 23	June 23	June 23	June 23	Aug. 27	Aug. 27
Ramsey.....	F	1907	3, 9	c, s	3 to 6	do.	do.	do.	do.	do.	do.	do.	Aug. 25	Aug. 25
Riparia X (Cordifolia Rupestris, No. 100-8).....	F	1909	3, 5	s	3 to 5	Mar. 7	May 22	June 22	June 22	June 22	June 22	June 22	Oct. 16	Oct. 16
Riparia X Rupestris, No. 101.....	O	1907	4, 8	s	3 to 5	Mar. 22	May 23	June 23	June 23	June 13	June 28	June 25	Sept. 24	Oct. 1
Riparia X Rupestris, No. 101-14.....	F	1909	8, 6	s	3 to 5	Mar. 8	May 20	June 21	June 21	June 13	June 28	June 25	Oct. 16	Oct. 16
Riparia X Rupestris, No. 3306.....	F	1909	2, 6	s	3 to 5	Mar. 20	May 20	June 21	June 21	June 21	June 21	June 21	do.	do.
Riparia X Rupestris, No. 3309.....	O	1904	4, 1	c, s	3 to 5	Mar. 24	May 27	June 27	June 27	June 12	June 2	June 14	Sept. 23	Oct. 5
Riparia X Rupestris de Jaeger, No. 201.....	F	1907	2, 1	c, s	3 to 6	Mar. 18	May 22	June 22	June 22	June 12	June 2	June 14	Sept. 23	Oct. 5
Rupestris des Caussees.....	F	1907	2	c, s	3 to 5	Mar. 19	May 18	June 23	June 23	June 12	June 2	June 14	Sept. 23	Oct. 5
Rupestris Marlin.....	F	1909	2, 5	s	3 to 5	Mar. 31	May 16	June 24	June 24	June 24	June 4	June 4	do.	do.
Rupestris Mission.....	F	1907	4, 6	s	3 to 5	Mar. 20	May 21	June 21	June 21	June 21	June 4	June 4	Oct. 16	Oct. 16
Rupestris X Berlandieri, No. 219A.....	F	1907	4, 3	c, s	3 to 5	Mar. 12	May 24	June 24	June 24	June 24	June 4	June 4	do.	do.
Salt Creek.....	F	1907	2, 3	c, s	3 to 5	Mar. 20	May 18	June 23	June 23	June 23	June 5	June 5	Sept. 5	Sept. 5
Solonis Robusta.....	F	1907	4, 3	c, s	3 to 6	Mar. 27	May 19	June 23	June 23	do.	do.	do.	Aug. 25	Aug. 25
Solonis X Otello, No. 1613.....	F	1907	5, 5	s	3 to 6	Mar. 22	May 23	June 23	June 23	do.	do.	do.	do.	do.
Solonis X Riparia, No. 1616.....	O	1904	4, 8	c, s	2 to 6	Mar. 16	May 24	June 24	June 24	June 10	May 29	June 25	Sept. 23	Do.
Yermorel.....	F	1907	4, 3	c, s	3 to 6	Mar. 18	May 22	June 22	June 22	June 10	May 29	June 25	Sept. 23	Sept. 28
Viala.....	F	1907	4	c, s	3 to 6	Mar. 19	May 17	June 23	June 23	May 22	June 6	June 6	Aug. 27	Sept. 21
Sutliffa Rosea:														
Lenoir.....	O	1904	2, 1	s	2 to 1	Mar. 20	May 24	June 24	June 24	June 8	May 28	June 20	Sept. 26	Oct. 6

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Sylvaner:															
Own roots.....	Cl	1904													
Aramon X Rupestris Ganzin, No. 1.....	O	1904	1905		c,s		2 to 4	Mar. 17	Mar. 26	May 26	May 28	May 31	June 1	Aug. 28	Sept. 23
Dog Ridge.....	O	1903	1905	95	c,s	6.4	2 to 5	Mar. 19	May 19	May 19	June 8	May 24	June 20	Sept. 20	Oct. 11
Herbement.....	O	1903	1905	92	c,s	3.8	2 to 5	Mar. 15	May 20	May 20	June 3	do.	June 15	Sept. 22	Oct. 8
Lenoir.....	O	1903	1905	93	c,s	4.5	2 to 4	Mar. 22	do.	do.	do.	do.	June 13	Sept. 20	Oct. 9
Riparia X Rupestris, No. 101.....	O	1904	1906	90	c,s	5.9	2 to 4	Mar. 18	May 19	do.	do.	May 24	June 16	Sept. 21	Oct. 6
Riparia X Rupestris, No. 3300.....	O	1904	1906	89	c,s	2.2	1 to 4	Mar. 21	do.	do.	do.	do.	June 24	Sept. 22	Oct. 9
Rupestris Martin.....	O	1903	1905	86	c,s	3.6	2 to 4	Mar. 17	May 23	do.	do.	do.	June 24	Sept. 20	Oct. 8
Rupestris St. George.....	O	1903	1905	86	c,s	3.1	2 to 5	Mar. 20	May 18	June 1	June 2	May 28	June 19	do.	do.
Sylvaner.....															
Rupestris St. George.....	O	1903	1905	86	c,s	3.5	2 to 4	Mar. 21	May 18	June 1	June 1	May 23	June 14	do.	do.
Syrian:															
Own roots.....	Cl	1904			s										
Tadone:															
Dog Ridge.....	O	1904	1906	88	c,s	1.9	3 to 5	Mar. 18	May 24	June 1	June 1	May 29	June 17	Sept. 30	Oct. 4
Lenoir.....	O	1903	1905	90	s	2.2	2 to 6	Mar. 15	May 26	June 2	June 3	May 30	June 18	Sept. 28	Oct. 10
Rupestris St. George.....	O	1903	1905	88	s	2.8	2 to 4	Mar. 16	May 24	June 3	June 3	May 29	June 18	do.	do.
Taylor Narbonne.....	O	1904	1906	84	s	1.6	2 to 6	Mar. 19	May 26	June 2	June 2	June 2	June 16	Sept. 27	Oct. 4
Tannat:															
Own roots.....	Cl	1904			c,s										
Dog Ridge.....	O	1904	1906	93	s	3.5	3 to 5	Mar. 26	May 25	June 7	June 7	May 30	June 9	Aug. 29	Sept. 23
Lenoir.....	O	1903	1905	89	s	2.1	2 to 5	Mar. 20	May 23	June 5	June 5	May 28	June 25	Sept. 20	Oct. 6
Rupestris St. George.....	O	1903	1905	89	s	4.1	3 to 6	do.	do.	do.	do.	do.	June 25	Sept. 27	Oct. 6
Taylor Narbonne.....	O	1904	1907	74	s	1.5	2 to 4	Mar. 24	May 24	June 6	June 6	May 29	June 20	Sept. 24	Oct. 8
Teneron:															
Own roots.....	Cl	1904			s										
Lenoir.....	O	1904	1907	89	s	2.6	2 to 6	Mar. 24	May 22	June 8	June 8	May 27	June 9	Sept. 15	Sept. 24
Riparia X Rupestris, No. 3300.....	O	1904		94	c,s	7	2 to 4	Mar. 20	May 12	June 4	June 4	May 16	May 18	Oct. 1	Oct. 8
Terret Monstre:															
Adobe Giant.....	F	1907	1907	95	c,s	3.3	2 to 4	do.	May 15	May 23	May 23	May 30	do.	Sept. 16	Sept. 21
Canada.....	F	1907	1907	97	c,s	5.1	3 to 4	Mar. 14	May 15	do.	do.	May 29	do.	do.	do.
Lenoir.....	O	1904	1907	89	s	2.4	2 to 5	Mar. 25	May 24	June 8	June 8	May 29	June 23	June 23	Oct. 7
Riparia X Rupestris, No. 101.....	F	1907	1907	95	c,s	5.5	3 to 5	Mar. 23	May 20	May 21	May 21	May 31	do.	Sept. 15	Sept. 21

Rupestris des Caussettes.	1907	97	c, s	4.7	3 to 5	Mar. 20	do.	May 16	May 23	May 30	Sept. 16	Do.
Solonis X Riparis, No. 1616.	1907	94	c, s	5.1	3 to 5	Mar. 17	do.	May 17	do.	May 28	do.	Oct. 17
Valencia.	1907	98	s	7.8	3 to 4	Mar. 17	do.	May 17	May 22	May 30	do.	do.
Tinta Amarella:												
Own roots.	1904	95	s	6.6	2 to 4	Mar. 19	Mar. 30	May 28	June 3	June 2	Sept. 18	Sept. 20
Dog Ridge.	1904	96	c, s	4	2 to 4	Mar. 19	do.	May 27	do.	June 3	Sept. 25	Oct. 6
Lenoir.	1903	90	c, s	4	2 to 4	Mar. 17	do.	May 20	do.	May 29	do.	Do.
Rupestris St. George.	1905	94	s	4.7	2 to 4	Mar. 17	do.	May 24	June 5	May 29	Sept. 26	Do.
Taylor Narbonne.	1906	88	s	2.3	2 to 4	Mar. 21	do.	May 25	do.	May 30	June 25	Oct. 4
Tinta Cao:												
Own roots.	1904	94	s	7.7	2 to 4	Mar. 15	Apr. 5	May 24	May 30	May 29	Aug. 29	Sept. 22
Dog Ridge.	1904	94	c, s	4.9	2 to 4	Mar. 21	do.	do.	June 5	do.	do.	Oct. 8
Lenoir.	1903	95	s	5.1	2 to 5	Mar. 17	do.	May 19	June 8	May 25	Sept. 25	Do.
Rupestris St. George.	1905	94	s	5.1	2 to 5	Mar. 20	do.	do.	June 1	May 25	Sept. 26	Oct. 6
Taylor Narbonne.	1906	90	s	3.6	1 to 4	Mar. 19	do.	May 23	May 31	May 29	Sept. 30	Oct. 4
Tinta de Madeira:												
Own roots.	1904	92	s	3.8	2 to 4	Mar. 19	Mar. 29	May 24	May 30	do.	June 7	Sept. 20
Dog Ridge.	1907	92	c, s	6.3	2 to 4	Mar. 18	do.	May 25	June 1	June 5	Aug. 28	Oct. 2
Lenoir.	1903	94	s	6	2 to 4	Mar. 20	do.	May 26	June 1	June 6	Sept. 24	Oct. 4
Rupestris St. George.	1905	96	c, s	2	2 to 4	Mar. 20	do.	May 27	May 31	May 27	Sept. 10	Sept. 23
Taylor Narbonne.	1907	87	s	2	2 to 4	Mar. 18	do.	May 26	June 3	June 2	Sept. 10	Sept. 23
Torok Goher Noir:												
Own roots.	1904	90	s	2.7	2 to 5	Mar. 12	Mar. 28	do.	May 31	May 31	Sept. 18	Sept. 25
Traminer:												
Own roots.	1907	88	s	2.6	2 to 4	Mar. 16	Apr. 1	do.	June 3	June 2	Sept. 23	Oct. 5
Aramon X Rupestris Ganzin, No. 1.	1906	88	c, s	1.7	2 to 5	Mar. 20	do.	May 22	June 3	June 2	Sept. 25	Oct. 10
Dog Ridge.	1903	82	s	3.1	2 to 5	Mar. 18	do.	May 20	June 6	May 26	do.	do.
Herbmont.	1905	90	s	1.5	2 to 5	Mar. 20	do.	May 25	June 2	June 20	Sept. 20	Oct. 5
Lenoir.	1903	84	s	1.3	2 to 5	Mar. 18	do.	May 26	June 3	do.	Sept. 23	Oct. 6
Mourvedre X Rupestris, No. 1202.	1906	85	s	2.7	2 to 5	Mar. 17	do.	May 22	June 2	do.	Sept. 26	Do.
Riparia X Rupestris, No. 101.	1904	89	c, s	2.2	2 to 5	Mar. 20	do.	May 23	June 1	May 28	Sept. 24	Do.
Riparia X Rupestris, No. 3309.	1905	85	s	3.5	2 to 4	Mar. 17	do.	May 20	June 4	May 28	Sept. 25	Oct. 6
Rupestris Martin.	1903	92	c, s	1.9	2 to 5	Mar. 20	do.	May 24	June 1	May 27	Sept. 12	Sept. 30
Rupestris St. George.	1905	88	s	2.3	2 to 4	Mar. 18	do.	May 22	June 5	May 27	Sept. 26	Oct. 6
Self Creek.	1903	88	s	3.4	2 to 5	May 17	do.	May 26	June 10	June 3	Sept. 21	Oct. 5
Solonis X Riparia, No. 1616.	1906	92	s	3.3	3 to 5	Mar. 15	Apr. 1	May 26	May 28	June 2	Sept. 23	Oct. 5
Taylor Narbonne.	1906	92	s	2.1	2 to 4	Mar. 18	do.	May 27	June 1	June 3	Sept. 23	Oct. 5
Triplarin:												
Own roots.	1904	91	c, s	2.6	2 to 6	Mar. 20	do.	May 23	June 10	May 29	Sept. 20	Oct. 1
Lenoir.	1905	90	c, s	2.9	3 to 5	Mar. 16	do.	do.	June 5	June 1	Sept. 9	Oct. 8
Rupestris St. George.	1905	84	s	2.6	2 to 6	Mar. 20	do.	May 22	June 2	May 28	Sept. 24	Oct. 8
Trifere du Japon:												
Lenoir.	1907	92	s	2.9	3 to 5	Mar. 16	do.	do.	June 5	do.	Sept. 27	Oct. 15
Triomphe:												
Lenoir.	1907	92	s	2.6	2 to 6	Mar. 20	do.	May 23	June 10	May 29	Sept. 29	Oct. 10
Triplot:												
Own roots.	1904	90	s	2.1	2 to 4	Mar. 22	do.	do.	June 5	do.	Sept. 27	Oct. 15
Trojka:												
Own roots.	1904	90	s	2.1	2 to 4	Mar. 20	Apr. 10	May 31	June 12	June 6	Sept. 24	Oct. 25
Own roots.	1904	90	s	2.6	2 to 6	Mar. 26	Mar. 29	May 24	June 3	May 29	June 10	Aug. 20

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment vineyard.	Year stock was planned.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
Trousseau:															
Owens roots.....	Cl	1904	1906		c, s	Lvs.	1 to 5	Mar. 20	Mar. 26	May 26	May 30	June 31	June 7	Aug. 21	Sept. 20
Dog Ridge.....	O	1901	1905	63	s	4.6	1 to 5	Mar. 22	Mar. 26	May 26	June 3	June 2	June 5	Sept. 25	Oct. 28
Herbimont.....	O	1903	1905	63	c, s	3.1	1 to 4	Mar. 20	Mar. 26	May 26	June 1	June 1	June 13	Sept. 24	Oct. 11
Riparia St. George.....	O	1903	1905	63	c, s	4.8	1 to 4	Mar. 16	Mar. 21	May 21	June 1	May 30	June 14	Sept. 25	Oct. 25
Taylor Narbonne.....	O	1904	1906	94	s	6.5	2 to 4	Mar. 21	Mar. 27	May 27	June 3	June 3	June 14	Sept. 23	Oct. 25
Ullidale:															
Blanche d'Ambree.....	Cl	1905			c, s			Mar. 20	Mar. 31	May 28	May 30	June 1	June 1	Sept. 15	Sept. 25
Valdevenas:															
Dog Ridge.....	O	1904	1906	92	s	4	2 to 4	Mar. 16	Mar. 18	May 26	June 3	June 2	June 15	Sept. 27	Oct. 7
Lehigh.....	O	1904	1906	90	s	3	3 to 4	Mar. 18	Mar. 18	May 27	June 1	June 1	June 14	Sept. 28	Oct. 7
Valdevenas:															
Araucan X Ruprestis Ganzi, No. 1.....	Cl	1901	1905	65	s	4.7	2 to 5	Mar. 10	Apr. 1	May 26	May 30	May 30	June 5	Aug. 29	Sept. 23
Dog Ridge.....	O	1903	1905	84	c, s	3.9	1 to 5	Mar. 20	Mar. 20	May 21	June 4	May 27	June 15	Sept. 24	Oct. 11
Herbimont.....	O	1903	1905	65	c, s	5.7	1 to 4	Mar. 25	Mar. 25	May 24	June 5	May 29	June 20	Sept. 20	Oct. 8
Lehigh.....	O	1903	1905	65	s	4.4	2 to 4	Mar. 15	Mar. 15	May 26	June 3	June 3	June 17	Sept. 20	Oct. 11
Menyevre X Ruprestis, No. 1202.....	O	1904	1906	91	c, s	3.9	2 to 5	Mar. 18	Mar. 18	May 26	June 2	May 30	June 13	Sept. 25	Oct. 9
Riparia X Ruprestis, No. 101.....	O	1904	1906	95	s	4.2	2 to 5	Mar. 20	Mar. 20	May 27	June 2	May 30	June 15	Sept. 24	Oct. 9
Riparia X Ruprestis, No. 3360.....	O	1904	1906	91	s	6.3	1 to 4	Mar. 18	Mar. 18	May 27	June 3	May 31	June 18	Sept. 23	Oct. 6
Ruprestis Martin.....	O	1903	1905	96	s	5.6	2 to 4	Mar. 20	Mar. 20	May 22	June 3	May 25	June 17	Sept. 25	Oct. 8
Ruprestis St. George.....	O	1903	1905	92	c, s	2.8	2 to 5	Mar. 20	Mar. 20	May 22	June 3	May 27	June 16	Sept. 22	Oct. 11
Salt Creek.....	O	1903	1905	91	s	3.4	2 to 5	Mar. 18	Mar. 18	May 25	June 10	May 29	June 4	Sept. 24	Oct. 10
Solonis X Riparia, No. 1616.....	O	1904	1906	92	c, s	5	2 to 5	Mar. 23	Mar. 23	May 25	June 8	May 29	June 23	Sept. 23	Oct. 10
Taylor Narbonne.....	O	1904	1906	92	c, s			Mar. 13	Mar. 28	May 22	May 28	May 27	June 3	Aug. 28	Sept. 20
Valdevenas:															
Dog Ridge.....	Cl	1904	1906	92	c, s	3	3 to 5	Mar. 20	Mar. 20	May 24	June 1	May 29	June 16	Sept. 24	Oct. 9
Herbimont.....	O	1903	1905	88	c, s	2.7	3 to 5	Mar. 24	Mar. 24	May 20	June 5	May 26	June 17	Sept. 23	Oct. 8
Lehigh.....	O	1903	1905	90	s	3.1	2 to 4	Mar. 20	Mar. 20	May 20	June 3	May 25	June 14	Sept. 24	Oct. 8
Riparia X Ruprestis, No. 3306.....	O	1907	1909	79	s	9		do	do	June 1	do	June 5	June 14	Sept. 24	Oct. 8
Ruprestis Martin.....	O	1903	1905	84	s	1.8	2 to 4	Mar. 24	Mar. 24	May 19	June 5	May 25	June 20	Sept. 26	Oct. 6
Ruprestis St. George.....	O	1903	1905	85	s	2.1	2 to 5	Mar. 24	Mar. 24	May 20	June 31	May 27	June 10	Sept. 25	Do.
Salt Creek.....	O	1903	1905	88	s	2.4	2 to 5	Mar. 20	Mar. 20	May 19	June 1	May 25	June 16	Sept. 20	Oct. 11

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Experiment yard.	Year stock was planted.	Year grafted.	Congeniality.	How pruned.	Weight of pruning per vine.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-selling date.		Fruit-ripening date.	
								Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.	Early sea-son.	Late sea-son.
White Hanapeol:															
Adobe Chant.	F	1904	1906	88	s	1.5	3 to 5	Mar. 24	Apr. 2	May 20	May 28	May 29	June 4	Sept. 5	Sept. 12
Aramon X Rupestris Ganzin, No. 2.	F	1907	1909	88	s	1.5	2 to 5	Mar. 20	Apr. 2	May 15	May 31	May 25	June 7	Sept. 16	Sept. 20
Australis.	F	1907	1909	90	s	3	2 to 5	Mar. 30	Apr. 1	May 22	May 30	May 26	June 8	Sept. 10	Sept. 20
Berlandieri X Riparia, No. 420A.	F	1907	1909	88	s	3	2 to 5	Apr. 1	Apr. 1	May 20	May 30	May 26	June 8	Sept. 10	Sept. 20
Dee Ridge.	F	1904	1906	91	s	3.8	2 to 5	Apr. 13	Apr. 6	May 18	May 30	May 29	June 8	Sept. 5	Sept. 20
Honolulu.	F	1907	1909	87	s	2.2	3 to 5	Mar. 20	Apr. 5	May 21	May 27	May 29	June 6	Sept. 12	Sept. 19
Monticola X Rupestris.	F	1907	1909	89	s	4	3 to 5	Mar. 27	Apr. 3	May 15	May 25	May 26	May 30	Sept. 4	Sept. 20
Neurvedere X Rupestris, No. 1202.	F	1903	1905	89	s	3.5	3 to 5	Mar. 14	Apr. 5	May 20	May 25	May 30	June 7	Sept. 8	Sept. 20
Riparia X (Cordifolia X Rupestris, No. 106-8).	F	1907	1909	90	s	3.5	2 to 5	Mar. 24	Apr. 2	do.	do.	May 25	June 7	Sept. 8	Sept. 20
Riparia X Rupestris, No. 101.	F	1904	1906	86	s	1.3	2 to 5	Mar. 12	Apr. 2	May 15	May 31	May 25	June 7	Sept. 16	Sept. 20
Riparia X Rupestris, No. 336b.	F	1907	1909	88	s	3	2 to 5	Mar. 30	Apr. 1	May 22	May 30	May 26	June 8	Sept. 10	Sept. 20
Riparia X Rupestris, No. 336c.	F	1904	1906	92	s	4.1	2 to 5	Mar. 14	Apr. 1	May 20	May 30	May 26	June 8	Sept. 10	Sept. 20
Riparia X Martin.	F	1907	1909	80	s	2.9	3 to 6	Mar. 23	Apr. 5	May 24	May 27	May 29	June 6	Sept. 12	Sept. 19
Rupestris St. George.	F	1903	1905	82	s	2	3 to 6	Mar. 15	Apr. 5	do.	do.	May 29	June 6	Sept. 12	Sept. 19
Rupestris X Berlandieri, No. 219A.	F	1907	1909	84	s	2.6	3 to 6	Mar. 31	Apr. 3	May 15	May 25	May 26	May 30	Sept. 4	Sept. 20
Taylor Narbonne.	F	1904	1906	89	s	2	3 to 6	Mar. 8	Apr. 3	May 14	May 25	May 26	May 30	Sept. 4	Sept. 20
Villa.	F	1907	1909	87	s	2	3 to 6	Mar. 24	Apr. 3	May 15	May 25	May 26	May 30	Sept. 4	Sept. 20
White Tokay:															
Owntoots.	Cl	1904			s			Mar. 12	Mar. 28	May 22	May 30	May 27	June 5	Aug. 18	Sept. 24
Wilmot, Hamburg.	O	1904			s	2.7	3 to 4	Mar. 18	Mar. 28	May 16	May 31	May 21	June 5	Sept. 23	Sept. 25
Do.	Cl	1905			c, s			Mar. 14	Mar. 28	May 27	May 28	June 2	do.	Aug. 28	Sept. 25
Wilmot, No. 16:															
Owntoots.	Cl	1905			c, s			Mar. 20	Mar. 31	May 15	May 26	May 26	June 12	Sept. 25	Oct. 10
Dee Ridge.	O	1904	1906	89	s	2.4	3 to 6	Mar. 21	Mar. 21	May 20	June 2	May 29	June 16	Sept. 28	Oct. 15
Lenoir.	O	1903	1905	91	s	2.8	2 to 5	Mar. 20	Mar. 20	May 25	May 31	May 29	June 16	Sept. 25	Oct. 30
Rupestris St. George.	O	1903	1905	89	s	2.2	3 to 6	Mar. 21	Mar. 21	May 22	do.	May 28	June 14	Sept. 25	Sept. 30
Zanfo.	Cl	1904			s			Mar. 22	Mar. 27	do.	May 22	do.	June 7	Sept. 5	Sept. 24

Zinzillasa: Ow'n roots. Zinfandel:	GI	1904	S	Mar. 20	Apr. 1	May 24	June 3	May 30	June 9	Aug. 29	Sept. 25
Ow'n roots.	GI	1907	1906	S	92	4.1	Mar. 30	June 27	June 1	June 2	June 14	Sept. 15	Sept. 23
Aramon X Rupestris Ganzin, No. 1.	O	1907	1909	S	83	1	May 30	June 4	June 9	June 15	Sept. 20	Oct. 11
Berlandieri X Riparia, No. 420A	O	1903	1905	S	93	3	May 23	June 5	May 29	June 17	Sept. 25
Dog Ridge	O	1903	1905	S	86	2.5	Mar. 22	May 22	June 5	May 28	June 15	do.	Oct. 8
Herbmont.	O	1903	1905	S	91	2.9	Mar. 24	do.	do.	do.	do.	Oct. 11
Lenoir	O	1907	1909	S	85	1.9	Mar. 17	do.	May 27	June 20	Sept. 20	Oct. 8
Monticola X Riparia, No. 18815	O	1904	1906	S	80	2.9	Mar. 18	June 1	June 3	June 6	June 23	do.	Oct. 10
Mourvedre X Rupestris, No. 1202	O	1904	1906	S	85	2.5	May 27	June 20	Sept. 20	Oct. 8
Riparia X Rupestris, No. 101	O	1907	1909	S	82	1.3	June 4	June 21	do.	Oct. 10
Riparia X Rupestris, No. 101-14	O	1907	1909	S	86	1.8	May 28	June 23	Sept. 25	Oct. 11
Riparia X Rupestris, No. 3506	O	1901	1906	S	89	1.8	Mar. 18	May 31	June 1	May 27	June 20	Sept. 20	Oct. 11
Riparia X Rupestris, No. 3509	O	1903	1905	S	87	3.5	Mar. 20	May 24	June 2	June 4	June 16	Sept. 20	Oct. 5
Rupestris Martin	O	1907	1909	S	80	1.4	June 4	June 21	Sept. 20	Oct. 11
Rupestris Metallica	O	1907	1909	S	70	June 5	June 19	Sept. 25	Oct. 13
Rupestris Mission	O	1907	1909	S	87	June 4	June 21	Sept. 20	Oct. 11
Rupestris St. George	O	1903	1905	S	92	3.1	Mar. 18	May 22	June 1	May 27	June 13	Sept. 28	Oct. 10
Salt Creek	O	1903	1905	S	86	1.5	Mar. 22	May 23	June 10	May 29	June 20	Sept. 20	Oct. 11
Solom's Obusta	O	1907	1909	S	79	1.6	Mar. 18	May 30	June 1	June 6	June 15	do.	do.
Solom's X Ribolla, No. 1613	O	1907	1909	S	68	1	Mar. 20	June 1	June 5	June 5	June 20	Sept. 25
Solom's X Riparia, No. 1616	O	1904	1906	S	72	1.1	May 29	June 26	Sept. 29	Oct. 10
Taylor Narbonne	O	1904	1906	S	82	2.1	May 25	June 22	Sept. 25	Oct. 8

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tartaric (grams per 100 c.c.).	Cluster.			Berry.			Use.
	1909							Size.	Shape.	Compact or loose.	Size.	Shape.	Color.	
	1909	1910	1911	1912	1913									
I	17	18	19	20	21	52	23	25	26	27	28	29	30	
Ach-I-Soum:						<i>P. cent.</i>								
Dog Ridge.....								t	t	t	t	w	s, st	
Lenoir.....								t	t	t	t	w	s, st	
Actoni-Maceroni:								t	t	t	t	w	s, st	
Monicola X Rupestris.....	2	10			3	21.8	0.5780	t	t	t	t	w	s, t	
Mourvedre X Rupestris, No. 1202.....				3	3	25	.4240	t	t	t	t	w	s, t	
Mourvedre X Rupestris, No. 1202.....								t	t	t	t	w	s, t	
Solonis X Riparia, No. 1416.....			6		3	24.5	.4200	t	t	t	t	w	s, t	
Actoni-Y:								t	t	t	t	w	s, t	
Lenoir:	6	3	7			24.8	.1750	t	t	t	t	w	s, t	
Monicola X Rupestris.....				2		20	.0050	t	t	t	t	w	s, t	
Molly.....								t	t	t	t	w	s, t	
Admirable.....	25		21	7		22.5	.7150	cy	m	m	m	w	t, w	
Riparia X Rupestris, No. 3309.....								t	t	t	t	w	t, w	
Affendhaler.....			10	11½	8	19		t	c	c	c	b	s, t	
Own roots.....								t	c	c	c	w	s, st, t	
Agodia.....								t	c	c	c	w	s, t	
Riparia X Rupestris, No. 3309.....			8	3		21.8	.5600	t	c	c	c	w	s, t	
Ahaneur hon Ahmed:								t	c	c	c	w	s, t	
Ajaki Odia.....	8					28.5	.6562	t	m	m	m	w	t	
Mourvedre X Rupestris, No. 1202.....			6					t	m	m	m	w	t	
Riparia X Rupestris, No. 101.....			3	6		24.5	.5475	t	m	m	m	w	t	
Rupestris Metallica.....								t	m	m	m	w	t	
Ajmi:								t	s	s	s	w	t	
Lenoir.....								t	s	s	s	w	t	
Aleatico:								t	s	s	s	w	t	
Own roots.....	5	3	3	25		29	.7994	t	c	c	c	b	t, w	
Dog Ridge.....	25	10	5	2	5	27	.8235	t	c	c	c	b	t, w	
Lenoir.....	25	10	14		20	23.5		t	c	c	c	b	t, w	
Rupestris St. George.....	40	12	15	1	27½	23.5	.8156	t	c	c	c	b	t, w	
Taylor Narbonne.....	3					23.2	.8500	t	c	c	c	b	t, w	

Alexandria:		6	2	8	25.2	.6600	m	t	l	l	0	w	F, S, St, t, w, j
Own roots.....		8	5	11	26.5	.5900	m	t	l	l	0	w	F, S, St, t, w, j
Aramon X Rupestris Ganzin, No. 1.....		8	4	4	27.5	.5300	m	t	l	l	0	w	F, S, St, t, w, j
Aramon X Rupestris Ganzin, No. 2.....		8	5 1/2	9	28.5	.5250	m	t	l	l	0	w	F, S, St, t, w, j
Australis.....		11 1/2	9	9	29	.5250	m	t	l	l	0	w	F, S, St, t, w, j
Berlandieri X Riparia, No. 420A.....		10	15	3	26.5	.6284	m	t	l	l	0	w	F, S, St, t, w, j
Dog Ridge.....		2	8	5 1/2	25.5	.6284	m	t	l	l	0	w	F, S, St, t, w, j
Herbemont.....		8	8	0	27	.6056	m	t	l	l	0	w	F, S, St, t, w, j
Lenoir.....		1	8	4	28	.6150	m	t	l	l	0	w	F, S, St, t, w, j
Monticola X Riparia, No. 18808.....		10	12	15	28	.5325	m	t	l	l	0	w	F, S, St, t, w, j
Mourvedre X Rupestris, No. 1202.....		15	3	12	26.5	.6550	m	t	l	l	0	w	F, S, St, t, w, j
Riparia X Cordifolia X Rupestris, No. 106-S.....		10	3	12	27	.6018	m	t	l	l	0	w	F, S, St, t, w, j
Riparia X Rupestris, No. 101.....		10	3	7	27	.5350	m	t	l	l	0	w	F, S, St, t, w, j
Riparia X Rupestris, No. 101-14.....		12	3	18	28.5	.4800	m	t	l	l	0	w	F, S, St, t, w, j
Riparia X Rupestris, No. 3306.....		12	5	5	27	.4575	m	t	l	l	0	w	F, S, St, t, w, j
Riparia X Rupestris, No. 3309.....		2	4	5	27.5	.6490	m	t	l	l	0	w	F, S, St, t, w, j
Rupestris Marlbin.....		2	1 1/2	5 1/2	25.5	.4987	m	t	l	l	0	w	F, S, St, t, w, j
Rupestris X Berlandieri, No. 219A.....		10	1	2	26.7	.5738	m	t	l	l	0	w	F, S, St, t, w, j
Salt Creek.....		10	1	3	19	.8700	m	t	l	l	0	w	F, S, St, t, w, j
Solonis Robasia.....		4	5	10	28	.5181	m	t	l	l	0	w	F, S, St, t, w, j
Solonis X Othello, No. 1613.....		4	2	3	28	.5181	m	t	l	l	0	w	F, S, St, t, w, j
Solonis X Riparia, No. 1616.....		10	3	2	28	.6425	m	t	l	l	0	w	F, S, St, t, w, j
Taylor Narbonne.....		10	3	2	28	.6425	m	t	l	l	0	w	F, S, St, t, w, j
Viola.....		10	3	2	28	.6425	m	t	l	l	0	w	F, S, St, t, w, j
Alicante.....		25	10	10	24.4	.7068	m	t	l	l	0	b	t, w
Own roots.....		30	12	25	24	.7020	m	t	l	l	0	b	t, w
Lenoir.....		25	10	10	24.4	.7068	m	t	l	l	0	b	t, w
Riparia X Rupestris, No. 3306.....		30	12	25	24	.7020	m	t	l	l	0	b	t, w
Rupestris St. George.....		30	12	25	24	.7020	m	t	l	l	0	b	t, w
Rupestris St. George.....		30	12	25	24	.7020	m	t	l	l	0	b	t, w
Alicante (Black Alicante):		25	15	17	24.8	.6680	m	cy	c	m	r	b	w
Own roots.....		40	30	5	21.5	1.3200	m	cy	c	m	r	b	w
Alicante Bouschet:		40	30	5	24.9	.9707	m	cy	c	m	r	b	w
(Aestivatus X Monticola) X (Riparia X Rupestris, No. 554-5).....		40	30	5	24.9	.9707	m	cy	c	m	r	b	w
Aramon X Rupestris Ganzin, No. 1.....		40	8	3	22	1.0612	m	cy	c	m	r	b	w
Aramon X Rupestris Ganzin, No. 2.....		40	8	3	21.5	.9075	m	cy	c	m	r	b	w
Berlandieri X Riparia, No. 420A.....		40	8	6	21	1.1672	m	cy	c	m	r	b	w
Constantia.....		40	8	10	20	.8300	m	cy	c	m	r	b	w
Dog Ridge.....		20	6	7	20.6	1.0008	m	cy	c	m	r	b	w
Herbemont.....		50	13	8	21.5	1.0270	m	cy	c	m	r	b	w
Lenoir.....		50	13	8	21.5	1.0270	m	cy	c	m	r	b	w
Monticola X Riparia, No. 18808.....		50	13	8	21.5	1.0270	m	cy	c	m	r	b	w
Monticola X Riparia, No. 18815.....		50	13	8	21.5	1.0270	m	cy	c	m	r	b	w
Mourvedre X Rupestris, No. 1202.....		50	13	8	21.5	1.0270	m	cy	c	m	r	b	w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling's scale.	Acid, as tartaric (c.c.) per 100	Cluster.			Berry.		Use.	
	1909	1910	1911	1912	1913			Shape.	Compact or loose.	Size.	Shape.	Color.		
	17	18	19	20	21									
1						22	23	24	25	26	27	28	29	30
Alcanta Bouschet—Continued.														
Riparia × Rupestris, No. 101.	60	10	25	2	P. cent. 21.5	1.0000	m	cv	c	m	r	b	w
Riparia × Rupestris, No. 101-14.	4	m	cv	c	m	r	b	w
Riparia × Rupestris, No. 3306.	11	3	5	21	1.3237	m	cv	c	m	r	b	w
Riparia × Rupestris, No. 3309.	60	10	6	3	23.8	.9650	m	cv	c	m	r	b	w
Rupestris Martin.	60	10	8	15	23	.8750	m	cv	c	m	r	b	w
Rupestris Metallica.	3	m	cv	c	m	r	b	w
Rupestris Mission.	60	15	8	6	22.4	.9073	m	cv	c	m	r	b	w
Rupestris St. George.	60	6	8	4	22.5	.7612	m	cv	c	m	r	b	w
Salt Creek.	60	6	5	m	cv	c	m	r	b	w
Solomis Robusta.	4	m	cv	c	m	r	b	w
Solomis × Obello, No. 1613.	4	5	23	1.0350	m	cv	c	m	r	b	w
Solomis Riparia, No. 1616.	40	20	7	238810	m	cv	c	m	r	b	w
Taylor Narbonne.	50	8	9	3	7	22	1.8950	m	cv	c	m	r	b	w
Almeria:														
Own roots.														
Herbement.	25	12	23	.7537	l	t	c	l	o	w	s, st, t
Lenoir.	20	15	20.8	.6100	l	t	c	l	o	w	s, st, t
Riparia × Rupestris, No. 101.	8	10	6	3	23.5	.6500	l	t	c	l	o	w	s, st, t
Riparia × Rupestris, No. 3309.	10	15	11	15	8	22.6	.7650	l	t	c	l	o	w	s, st, t
Solomis × Riparia, No. 1616.	30	30	20	23.5	.6700	l	t	c	l	o	w	s, st, t
Aneb el Cadi:														
Adobe Giant.	10	25.5	.7837	l	cy	m	m-1	ob	b	s, st, t
Lenoir.	10	10	9	20	.7533	l	cy	m	m-1	ob	b	s, st, t
Monticola × Rupestris.	l	cy	m	m-1	ob	b	s, st, t
Mourvedre × Rupestris, No. 1202.	8	23	.5700	l	cy	m	m-1	ob	b	s, st, t
Riparia Grand Glabre.	12	25	.6150	l	cy	m	m-1	ob	b	s, st, t
Angelina:														
Adobe Giant.	l	t	c	l	ob	r	s, st, t
Lenoir.	4	2	8	3	21.2	.5825	v-1	t	c	l	ob	r	s, st, t
Mourvedre × Rupestris, No. 1202.	25	.3105	v-1	t	c	l	ob	r	s, st, t

Armon:	60	20	19	4	8	20.7	.9300	t	c	l	o	b	W, S, t
Aramon × Rupestris Ganzin, No. 1.....	3	15	43	1.0008	t	c	l	o	b	W, S, t
Australis.....	5	2	22.6	t	c	l	o	b	W, S, t
Barnes.....	1	6000	t	c	l	o	b	W, S, t
Cornucopia.....	8	23.5	t	c	l	o	b	W, S, t
De Grasset.....	40	12	20	7	6	19.8	t	c	l	o	b	W, S, t
Dog Ridge.....	50	20	12	3	12	20	t	c	l	o	b	W, S, t
Herbement.....	20	20	5625	t	c	l	o	b	W, S, t
Hotrump.....	60	22	12	18	t	c	l	o	b	W, S, t
Lenoir.....	t	c	l	o	b	W, S, t
Monticola × Rupestris.....	40	8	15	2	3	19.5	t	c	l	o	b	W, S, t
Mourvedre × Rupestris, No. 1202.....	3	36	22	t	c	l	o	b	W, S, t
Ponroy.....	t	c	l	o	b	W, S, t
Ramsey.....	t	c	l	o	b	W, S, t
Riparia × Rupestris, No. 101.....	t	c	l	o	b	W, S, t
Do.....	60	12	16	8	24	21.7	t	c	l	o	b	W, S, t
Riparia × Rupestris, No. 3309.....	50	10	2	20.7	t	c	l	o	b	W, S, t
Riparia × Rupestris de Jaeger.....	2	19	t	c	l	o	b	W, S, t
Rupestris des Causseuses.....	t	c	l	o	b	W, S, t
Rupestris des Semis, No. 81-2.....	15	22	t	c	l	o	b	W, S, t
Rupestris Martin.....	4	t	c	l	o	b	W, S, t
Do.....	50	3	1	23.4	t	c	l	o	b	W, S, t
Rupestris St. George.....	55	10	11	4	16	24.5	t	c	l	o	b	W, S, t
Rupestris × Berlandieri, No. 219 A.....	20	20	21.5	t	c	l	o	b	W, S, t
Salt Creek.....	5	20	t	c	l	o	b	W, S, t
Do.....	60	10	17	2	5	24	t	c	l	o	b	W, S, t
Solonis × Othello.....	3	6	26	21.8	t	c	l	o	b	W, S, t
Solonis × Riparia, No. 1616.....	40	25	9	1	20.8	t	c	l	o	b	W, S, t
Do.....	55	20	15	22.9	t	c	l	o	b	W, S, t
Taylor Narbonne.....	4	21	t	c	l	o	b	W, S, t
Do.....	40	4	19.4	t	c	l	o	b	W, S, t
Valencia.....	13	20.8	t	c	l	o	b	W, S, t
Vermorel.....	15	25	20.5	t	c	l	o	b	W, S, t
Viola.....	3	3	23	t	c	l	o	b	W, S, t
Askari.....	4	8	22.2	t	c	l	o	b	W, S, t
Own roots.....	23.6	cy	l	m-l	o	b	S, t
Aspiran Noir.....	20	25	30	25	20.5	t	c	m	o	b	S, t, w
Own roots.....	t	c	l	l	b	S, t
Augulato.....	t	l	l	l	b	S, t
Adobe Giant.....	3	1½	5	4	20.8	t	l	l	l	b	S, t
Lenoir.....	t	l	l	l	b	S, t
Baba: Own roots.....	25	12	t	l	l	l	b	S, t
Dog Ridge.....	20	8	25	18	23.6	t	l	l	l	b	S, t
Lenoir.....	40	15	10	22.4	t	l	l	l	b	S, t
Rupestris St. George.....	30	10	30	25	t	l	l	l	b	S, t
Taylor Narbonne.....	50	8	8	21	t	l	l	l	b	S, t
.....	25	t	l	l	l	b	S, t
.....	5475	t	l	l	l	b	S, t

Barducci:	Adobe Giant.....	10	8	1½	22.5	.6502	l	cy	s	s	r	w	w
	Lenoir.....	6	3	15	19.6	.4350	l	cy	s	s	r	w	w
Bastardo:	Own roots.....	8	6	2	28	.7375	l	cy	c	c	r	b	b
	Dog Ridge.....	50	4	4½	25.9	.6750	l	cy	c	c	r	b	b
	Lenoir.....	60	11	7	24.3	.5208	l	cy	c	c	r	b	b
	Rupestris St. George.....	8	4	½	23.8	.9412	l	cy	c	c	r	b	b
	Taylor Narbonne.....	20	10	8	23.6	.5418	m	cy	l	l	r	b	b
Beclan:	Own roots.....	45	6	15	19.7	.8225	m	cy	l	l	r	b	b
	Dog Ridge.....	50	7½	20	19.7	.8225	m	cy	l	l	r	b	b
	Lenoir.....	50	8	6	23.3	.6693	m	cy	l	l	r	b	b
	Rupestris St. George.....	40	6	2	21	.6250	m	cy	l	l	r	b	b
	Taylor Narbonne.....	10	8	25	21.4	.5235	m	t	c	c	r	b	b
Bellino:	Own roots.....	6	10	2	21.3	.5775	m	t	c	c	r	b	b
	Adobe Giant.....	12	11	10	21.4	.5003	m	t	c	c	r	b	b
	Aramont X Rupestris Ganzin, No. 1.....	6	1	1	23	.6937	m	t	c	c	r	b	b
	Dog Ridge.....	6	1	7	21.9	.4931	m	t	c	c	r	b	b
	Do.....	12	1	3	25	.5025	m	t	c	c	r	b	b
	Do.....	5	8	11	23.4	.5700	m	t	c	c	r	b	b
	Mourvedre X Rupestris, No. 1202.....	10	14	20	24	.6675	m	t	c	c	r	b	b
	Ponroy.....	8	12	2	21.2	.4637	m	t	c	c	r	b	b
	Riparia X Rupestris, No. 101.....	12	15	4	22.6	.5600	m	t	c	c	r	b	b
	Riparia X Rupestris, No. 3309.....	8	11½	3	22.4	.6408	m	t	c	c	r	b	b
	Rupestris des Caussettes.....	12	4	23	21.9	.7937	m	t	c	c	r	b	b
	Rupestris St. George.....	8	12	8	21.9	.7937	m	t	c	c	r	b	b
	Do.....	12	4	1½	24.2	.5475	m	t	c	c	r	b	b
	Solon X Riparia, No. 1616.....	6	2	3	23	.6970	m	t	c	c	r	b	b
	Taylor Narbonne.....	12	2	14	24	.6600	m	t	c	c	r	b	b
	Do.....	8	2	2	21	.6675	m	t	c	c	r	b	b
	Vermorel.....	20	18	3	22.3	.4275	m	t	c	c	r	b	b
Bengi:	Own roots.....	2	18	3	26	.5575	m	cy	c	c	r	w	w
	Bermestia Blanca.....	20	18	3	26	.5575	m	cy	c	c	r	w	w
	Rupestris St. George.....	2	18	3	26	.5575	m	cy	c	c	r	w	w
Besson:	Own roots.....	20	18	3	26	.5575	m	cy	c	c	r	w	w
Bholtri:	Own roots.....	20	12	3	23	.5831	m	r	l	l	ob	w	w
Bicane:	Adobe Giant.....	20	12	3	23.9	.5875	m	r	l	l	ob	w	w
	Australs.....	20	11½	9	23.6	.6425	m	r	l	l	ob	w	w
	Barras.....	20	2	13	23.6	.6425	m	r	l	l	ob	w	w
	Berlandieri X Riparia, No. 420B.....	20	2	13	23.6	.6425	m	r	l	l	ob	w	w
	Constanita.....	20	2	13	23.6	.6425	m	r	l	l	ob	w	w
	De Grasset.....	20	5	9	22.6	.7875	m	r	l	l	ob	w	w
	Dog Ridge.....	20	5	9	22.6	.7875	m	r	l	l	ob	w	w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Varyety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).						Sugar, Balling's scale.	Acid, as tartaric (c.c.) per 100	Cluster.			Berry.			Use.
	1								Shape.	Compact or loose.	Size.	Shape.	Color.		
	1909	1910	1911	1912	1913	1913									
Burger—Continued.	17	18	19	20	21	21	22	23	24	25	26	27	28	29	30
Aramon X Rupestris Ganzin, No. 2.			4		3	3	17	0.9525	1	t	c	1	I	w	w
Berlandieri X Riparia, No. 420A.			2	4	3	3	21.5	.7350	1	t	c	1	I	w	w
Constandia.			3		12	12	21	.8025	1	t	c	1	I	w	w
Columbard X Rupestris No. 2302.			14	6	4	4	20.5	.9650	1	t	c	1	I	w	w
Dog Ridge.	60	5	18	3	20	17.9	1.0175	1.0175	1	t	c	1	I	w	w
Herbmont.	30	15	4	4	4	19	1.7300	1.7300	1	t	c	1	I	w	w
Lenoir.	20		10	3		24	1.0200	1.0200	1	t	c	1	I	w	w
Monticola X Riparia, No. 18898.			10	5	5	22.2	1.7744	1.7744	1	t	c	1	I	w	w
Mourvedre X Rupestris, No. 1202.	40	10	10	5	5	19	1.0050	1.0050	1	t	c	1	I	w	w
Riparia X Rupestris, No. 101.	40	10	12	1		23	.8025	.8025	1	t	c	1	I	w	w
Riparia X Rupestris, No. 101-14.			15	1	15	20	1.1742	1.1742	1	t	c	1	I	w	w
Riparia X Rupestris, No. 3306.	60	15				18.7	.7062	.7062	1	t	c	1	I	w	w
Riparia X Rupestris, No. 3369.	50	10	12		10½	17.9	1.0294	1.0294	1	t	c	1	I	w	w
Rupestris Madrilca.					4				1	t	c	1	I	w	w
Rupestris Mission.									1	t	c	1	I	w	w
Rupestris St. George.	30	5	10	4		21.1	.7462	.7462	1	t	c	1	I	w	w
Salt Creek.	30	10	8	2	6	31.5	1.0108	1.0108	1	t	c	1	I	w	w
Solonis Kobusca.			23	2		21.5	.9057	.9057	1	t	c	1	I	w	w
Solonis X Onitela, No. 1613.			13	10	2	22.5	1.5719	1.5719	1	t	c	1	I	w	w
Solonis X Riparia, No. 1616.	15	15	4			20.5	.8596	.8596	1	t	c	1	I	w	w
Taylor Narbonne.	50	15	19	6	8	16.7	.9337	.9337	1	t	c	1	I	w	w
Cabonnet Sauvignon.									1	t	c	1	I	w	w
Own roots.									m	t	m	m	I	b	w
Aramon X Rupestris Ganzin, No. 1.	15	6	5	1½	1	24.5	.7950	.7950	m	t	m	m	I	b	w
Dog Ridge.	30	2				21.3	.7825	.7825	m	t	m	m	I	b	w
Herbmont.	30	11	17	17	3	23.6	.8525	.8525	m	t	m	m	I	b	w
Lenoir.	25	8	4	2	4	27.00	.7700	.7700	m	t	m	m	I	b	w
Mourvedre X Rupestris, No. 1202.	10	4				23.4	.7925	.7925	m	t	m	m	I	b	w
Riparia X Rupestris, No. 101.	20	3	5			23.2	.6562	.6562	m	t	m	m	I	b	w
Riparia X Rupestris, No. 3309.	20	2	6		2	24.3	.7458	.7458	m	t	m	m	I	b	w
						25.4	.7100	.7100	m	t	m	m	I	b	w

Rupestris Martin.....	25	2	6	1	3	22.2	9475	m	t	m	m	r	b	w
Rupestris St. George.....	25	3	6	1	5	23.5	.7258	m	t	m	m	r	b	w
Salt Creek.....	20	5	9	1	5	25.1	.8775	m	t	m	m	r	b	w
Solonis X Riparia, No. 1616.....	20	12	6	2	2	23.6	.7862	m	t	m	m	r	b	w
Taylor Narbonne.....	25	4	9	3	3	20.6	.9225	m	t	m	m	r	b	w
Calabrian:														
Own roots.....		3	2	23.6	.5862	l	cy	l	l	r	w	t, s
Australis.....		5	3	24.6	.6175	l	cy	l	l	r	w	t, s
Lenoir.....		2	8	1	3	23.8	.5556	l	cy	l	l	r	w	t, s
Do.....		8	1	4	3	25.4	.6650	l	cy	l	l	r	w	t, s
Monticola X Rupestris.....		12	7	6 ³	10	25.5	.5175	l	cy	l	l	r	w	t, s
Riparia Gloire.....		12	7	3	4	26	.5393	l	cy	l	l	r	w	t, s
Rupestris des Caussettes.....		10	9	15	24	23.6	.5531	l	cy	l	l	r	w	t, s
Rupestris St. George.....		10	9	15	24	23.6	.5531	l	cy	l	l	r	w	t, s
Calmette:														
Adobe Glant.....		5	3	24.2	.7225	m	t	l	l	r	b	w
Aramon X Rupestris Ganzin, No. 1.....		2	4	24	.7425	l	t	l	l	r	b	w
Do.....		40	40	9	2	21.6	.7775	m	t	l	l	r	b	w
Berlandier X Riparia, No. 420B.....		5	6	25.2	.6625	m	t	l	l	r	b	w
Constantia.....		10	8	21.3	.9900	m	t	l	l	r	b	w
Herbmont.....		20	8	6	21	22.5	.8131	m	t	l	l	r	b	w
Lenoir.....		4	4	13	7	28	.6367	m	t	l	l	r	b	w
Monticola X Riparia, No. 18904.....		4	4	13	7	26	.6700	m	t	l	l	r	b	w
Monticola X Riparia, No. 18815.....		2	4	22.6	.6725	m	t	l	l	r	b	w
Monticola X Rupestris.....		2	4	22.6	.6725	m	t	l	l	r	b	w
Ponroy.....		m	t	l	l	r	b	w
Ramsay.....		m	t	l	l	r	b	w
Riparia X Rupestris, No. 101.....		m	t	l	l	r	b	w
Do.....		40	30	13	16	25	.6000	m	t	l	l	r	b	w
Riparia X Rupestris, No. 3209.....		m	t	l	l	r	b	w
Do.....		50	20	14	28	20.1	.8430	m	t	l	l	r	b	w
Riparia X Rupestris de Jaeger.....		m	t	l	l	r	b	w
Rupestris X Berlandier, No. 219A.....		m	t	l	l	r	b	w
Rupestris X Berlandier, No. 301-37-152.....		6	6	24.7	.7408	m	t	l	l	r	b	w
Salt Creek.....		m	t	l	l	r	b	w
Solonis Robusta.....		m	t	l	l	r	b	w
Solonis Othello.....		10	4	23	.7800	m	t	l	l	r	b	w
Solonis X Riparia, No. 1616.....		30	15	3	22	23.5	.6320	m	t	l	l	r	b	w
Do.....		30	15	3	3	21.1	.6350	m	t	l	l	r	b	w
Taylor Narbonne.....		m	t	l	l	r	b	w
Vermorel.....		m	t	l	l	r	b	w
Carlignane:														
Own roots.....		20	5	22.7	.8475	m	t	l	l	r	b	w
Aramon X Rupestris Ganzin, No. 1.....		60	40	15	9	22.7	.8475	m	t	l	l	r	b	w
Do.....		60	40	15	9	22.7	.8475	m	t	l	l	r	b	w
Dog Ridge.....		60	12	20	16	21.7	1.0575	m	t	l	l	r	b	w
Herbmont.....		50	10	14	5	18	.9015	m	t	l	l	r	b	w
Lenoir.....		55	20	16	12	20.5	.9619	m	t	l	l	r	b	w
Mourvedre X Rupestris, No. 1202.....		60	15	15	8	21	1.2924	m	t	l	l	r	b	w
Riparia X Rupestris, No. 101.....		60	10	11	5	22.4	.8006	m	t	l	l	r	b	w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—(Continued).

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tartaric (grams per 100 c. c.).	Cluster.			Berry.			Use.	
	1909	1910	1911	1912	1913			Shape.	Compact or loose.	Size.	Shape.	Color.	Size.		Shape.
Chasselas Rose de Falloux:															
Own roots.....	2					<i>P</i>	0.3475	m	cy	c	m	f	w, f		
Aramon X Rupestris Ganzin, No. 1.....	25	4	4	1½	7	21.6	6115	m	cy	c	m	f	w, f		
Do.....		3	3½	16	4	20.5	3967	m	cy	c	m	f	w, f		
Australis.....	20					21.9	4850	m	cy	c	m	f	w, f		
Dog Ridge.....	16					20.8	5643	m	cy	c	m	f	w, f		
Lenoir.....	20					20.2	5458	m	cy	c	m	f	w, f		
Monroville X Rupestris, No. 1202.....	20		1		5	19.8	5550	m	cy	c	m	f	w, f		
Riparia X Rupestris, No. 101.....	20					20	4200	m	cy	c	m	f	w, f		
Do.....	20		4		6	20.8	1510	s	cy	c	m	f	w, f		
Riparia X Rupestris, No. 3309.....	30							m	cy	c	m	f	w, f		
Do.....	12		12		4	20.6	6243	s	cy	c	m	f	w, f		
Do.....	20	4	8	3	10	24.8	4131	s	cy	c	m	f	w, f		
Do.....	20		11		4	21.2	5700	s	cy	c	m	f	w, f		
Solonis X Riparia, No. 1046.....	20							m	cy	c	m	f	w, f		
Taylor Narbonne.....	15					20.8	5412	s	cy	c	m	f	w, f		
Do.....			5	2	2½	17	5400	s	cy	c	m	f	w, f		
Do.....				4	8	21.8	5225	s	cy	c	m	f	w, f		
Do.....				1	6			m	cy	c	m	f	w, f		
Chasselas Rose Royal:															
Aramon X Rupestris Ganzin, No. 1.....	10		3			20.3	4870	s	cy	c	m	f	w, w		
Dog Ridge.....	25					20.6	4125	s	cy	c	m	f	w, w		
Lenoir.....	25		4		3	22.3	4350	s	cy	c	m	f	w, w		
Rupestris St. George.....	11		6½		2	23	5275	s	cy	c	m	f	w, w		
Taylor Narbonne.....	3					21.2	4725	s	cy	c	m	f	w, w		
Chasselas Rouge:															
Own roots.....	4		5	5	18	21.9	4346	s	cy	c	m	f	w, w		
Lenoir.....	20		5		8	20.8	4675	s	cy	c	m	f	w, w		
Rupestris St. George.....	15		8	½	3	22	5146	s	cy	c	m	f	w, w		
Chasselas St. Bernard:															
Canada.....								m	cy	f	m	f	w, w		
Lenoir.....								m	cy	f	m	f	w, w		
Riparia X Rupestris, No. 101.....	6					17.8	3600	m	cy	f	m	f	w, w		

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Ballin g scale.	Acid, as tartaric c.c.) (grams per 100	Cluster.			Berry.			Use.
	1909	1910	1911	1912	1913			Size,	Shape	Compact or Loose	Size,	Shape.	Color.	
Corinthine Rose:														
Adobe Giant.....						23	0.8400	c	cy	s	I	I	I	I, F, W
Canada.....				1½				c	cy	s	I	I	I	I, F, W
Constantia.....					3	26.5	.9400	c	cy	s	I	I	I	I, F, W
Lenoir.....		2				28	.6525	c	cy	s	I	I	I	I, F, W
Monicoia X Rupestris.....		2				26	.5475	c	cy	s	I	I	I	I, F, W
Mourvedre X Rupestris, No. 1202.....								c	cy	s	I	I	I	I, F, W
Riparia Grand Glabre.....								c	cy	s	I	I	I	I, F, W
Riparia X Rupestris, No. 101.....								c	cy	s	I	I	I	I, F, W
Vernorel.....								c	cy	s	I	I	I	I, F, W
Conistano:														
Australis.....		15	41			20.6	1.1058	c	†	s	I	I	I	I, W
Barnes.....								c	†	s	I	I	I	I, W
Lenoir.....	20	4	10	8	28	18.4	.8890	c	†	s	I	I	I	I, W
Monicoia X Rupestris.....		6	37	25		21.1	.6750	c	†	s	I	I	I	I, W
Rupestris Martin.....		4				19.8	1.0870	c	†	s	I	I	I	I, W
Rupestris St. George.....		8	10			19.6	1.2050	c	†	s	I	I	I	I, W
Comnelton:														
Own roots.....		8	6	12		20	.5750	†	†	1	o	o	o	I, S
Aramon X Rupestris Ganzin, No. 1.....	40					20.2	.6525	†	†	1	o	o	o	I, S
Lenoir.....			8			21	.9575	†	†	1	o	o	o	I, S
Riparia Glorio.....						20	.9575	†	†	1	o	o	o	I, S
Rupescis X Berlandieri, No. 301A.....			8			21	.7725	†	†	1	o	o	o	I, S
St. Hubista.....			8					†	†	1	o	o	o	I, S
Grables Burgandy.....								†	†	1	o	o	o	I, S
Grables:														
Own roots.....	45			4	3	22	.9186	†	cy	m	I	I	I	W
Aramon X Rupestris Ganzin, No. 1.....	30					21.3	.8017	†	cy	m	I	I	I	W
Dog Ridge.....	40	12		2		21.3	.8419	†	cy	m	I	I	I	W
Lenoir.....	40	10	24½			21.2	.8880	†	cy	m	I	I	I	W
Riparia X Rupestris, No. 101.....	50	8	10	12	16	21.2	.8275	†	cy	m	I	I	I	W
Riparia X Rupestris, No. 3309.....	60	5	13	5	5	22.4	.7665	†	cy	m	I	I	I	W
Rupestris St. George.....	30	8	17	4	31	22.6	.8055	†	cy	m	I	I	I	W
Solonis X Riparia, No. 1616.....	50	20		3		19.5	.8150	†	cy	m	I	I	I	W
Taylor Narbonne.....	30	8		3	3	21.8	.6844	†	cy	m	I	I	I	W

	5	4	10	8	12	20	.8355	m	t	l	s	o	b	w
Croetto:														
Own roots.....														
Crujidero:	20		25	5	7	23.9	.5898	m	t	l	l	o	w	t
Own roots.....														
Danague:				21		22	.8025	m	t	l	l	r	b	t, s
Own roots.....	5		14		1	24.1	.5100	m	t	l	l	r	b	t, s
Dog Ridge.....	35			6		25.4	.5220	m	t	l	l	r	b	t, s
Lenoir.....	40		44		4	23.9	.5494	m	t	l	l	r	b	t, s
Rupestris St. George.....						22.6	.6150	m	t	l	l	r	b	t, s
Taylor Narbonne.....									cy	l	l	ob	w	t, s, r
Dattier de Beyrouth.....										c	l	ob	w	t
Lenoir.....				4						c	l	ob	w	t
Deacon Superb:						27	.9300	l	cy	c	l	ob	w	t
Own roots.....										c	l	ob	w	t
Diamant:				6		22	.6225	m	t	c	l	r	w	t, s
Own roots.....										c	l	r	w	t, s
Dodrelabi:						21.8	.4200	m	r	l	l	r	b	t, s
Own roots.....	2	2		3		16.7	.6990	m	r	l	l	r	b	t, s
Aramon X Rupestris Ganzin, No. 1.....			11	6	2	18.8	.7375	m	r	l	l	r	b	t, s
Dog Ridge.....	60	10				18.9	.6540	m	r	l	l	r	b	t, s
Herbement.....		8	12		4	18.8	.7375	m	r	l	l	r	b	t, s
Lenoir.....	40	10	12	14	9	18	.8437	m	r	l	l	r	b	t, s
Mourvedre X Rupestris, No. 1202.....			3			19.6	.5700	m	r	l	l	r	b	t, s
Rupestris St. George.....	30	10	28		12	20	.7600	m	r	l	l	r	b	t, s
Solanis X Riparia, No. 1616.....			8	3	5	20.2	.6545	m	r	l	l	r	b	t, s
Taylor Narbonne.....	50	3	9							l	l	ob	b	t, s
Drnekusi:						22.3	.5661	m	t	l	l	ob	b	t, s
Own roots.....			25	25	6					l	l	ob	b	t, s
Dronkane:						22.8	.8348	v, l	l	l	l	ob	w	t, s
Lenoir.....	10	2	13			27	.7500	v, l	l	l	l	ob	w	t, s
Rupestris des Causseilles.....				8						l	l	ob	w	t, s
Due de Macenta:										l	l	ob	w	t, s
Rupestris St. George.....	15	20	3		3	23.5	.5231	l	l	l	l	ob	w	t, s
Due de Malakoff:										l	l	ob	w	t, s
Own roots.....			4	7	8	24.8	.7106	l	t	m	l	r	w	t, s
Emethian:										m	l	r	w	t, s
Dog Ridge.....										l	l	o	w	t, s, st
Lenoir.....			5			23	.5737	m	t	l	l	o	w	t, s, st
Emporor.....										l	l	o	w	t, s, st
Lenoir.....										l	l	o	w	t, s, st
Rupestris St. George.....										l	l	o	w	t, s, st
Erz Kouml:										l	l	o	w	t, s, st
Lenoir.....										l	l	o	w	t, s, st
Rupestris St. George.....										l	l	o	w	t, s, st
Etraire de l'Adhui:										l	l	o	w	t, s, st
Own roots.....										l	l	o	w	t, s, st
Fajoumi Jaume:										l	l	o	w	t, s, st
Lenoir.....	6		5			21.2	.8005	l	cy	c	m	ob	w	t, s
Rupestris St. George.....			2	14		20	.6445	l	cy	l	l	ob	w	t, s
Faphly:										l	l	ob	w	t, s
Lenoir.....	25	10	13		15	19.7	.7369	v, l	cy	l	l	ob	r	t, s, st
Riparia X Rupestris, No. 3309.....			8			27	.6307	v, l	cy	l	l	ob	r	t, s, st
Rupestris St. George.....			24			26.6	.7500	v, l	cy	l	l	ob	r	t, s, st

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tartaric (c.c.) per 100	Cluster.			Berry.		Use.	
								Size.	Shape.	Compact or loose.	Size.	Shape.		Color.
	1909	1910	1911	1912	1913									
1	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Fehér Gohér Noir: Own roots.....	15	12	10	35	23	0.7406	1	cy	c	1	o	b	t, s, w
Fehér Som: Lenoir.....	10	5	6	4	12	24.5	.8775	1	cy	c	1	o	b	t, s, w
Fehér Szagos: Rupestris St. George.....						29	.6150	1	cy	c	1	o	b	t, s, w
Adobe Giant: Aramon X Rupestris Ganzin, No. 1.....	3	35	10	29	19	27	.5205	1	t	1	1	o	w	t, s, r
Dog Ridge.....	40	30	8	1	20	22	.5745	1	t	1	1	o	w	t, s, r
Herbemont.....	30	25	3	20	9	24.5	.6230	1	t	1	1	o	w	t, s, r
Lenoir.....	10	4	4	3	25.2	.6076	1	t	1	1	o	w	t, s, r
Mourvedre X Rupestris, No. 1202: Riparia Gloire.....	20	10	2	3	23.1	.5300	1	t	1	1	o	w	t, s, r
Riparia X Rupestris, No. 101: Riparia X Rupestris, No. 3309.....	40	20	4	9	4	23.8	.5320	1	t	1	1	o	w	t, s, r
Rupestris Martin.....	20	30	8	20	5	23.6	.5260	1	t	1	1	o	w	t, s, r
Rupestris St. George.....	40	40	5	15	23.5	.5043	1	t	1	1	o	w	t, s, r
Salt Creek.....	50	20	30	31	20.5	.5700	1	t	1	1	o	w	t, s, r
Solomis X Riparia, No. 1616: Taylor Narbonne.....	6	40	10	25	6	26	.4575	1	t	1	1	o	w	t, s, r
Ferrara: Herbemont.....	15	12	2	6	0	21.6	.5535	1	t	1	1	o	w	t, s, r
Lenoir.....	40	3	20	28	9	24.1	.5081	1	t	1	1	o	w	t, s, r
Fintondo: Own roots.....	10	15	6	28	3	27.4	.4890	1	t	1	1	o	w	t, s, r
Dog Ridge.....	25	15	23.6	.5812	v1	1	1	1	ob	b	t, s, st
Lenoir.....	12	12	19	.7575	v1	1	1	1	ob	b	t, s, st
Rupestris St. George.....	30	5	12	6	23.5	.7950	m	cy	c	m	ob	b	t, s, w
Taylor Narbonne.....	25	18	8	4	27	21.9	.7695	m	cy	c	m	ob	b	t, s, w
	30	5	14	1½	22.1	.7860	m	cy	c	m	ob	b	t, s, w
	35	8	9	10	22.4	.8175	m	cy	c	m	ob	b	t, s, w
						22.4	.7175	m	cy	c	m	ob	b	t, s, w

Flame Tokay:	1	2	3	25	21.4	.5212	t	c	l	ob	r	t, s
Own roots.....					25							
Canada.....					24	.4800	t					
Dog Ridge.....	30	12	18		15	.6600	t					
Lenoir.....	8		14		21.6	.5250	t					
Monticola X Riparia, No. 554			22		27	.7575	t					
Monticola X Riparia, No. 18304			18				t					
Monticola X Riparia, No. 18308			18		26	.5850	t					
Monticola X Riparia, No. 18185			11		24	.4875	t					
Riparia a Grandes Feuilles			16		22.1	.5950	t					
Riparia X Rupestris, No. 101	30	25	15		21.9	.6225	t					
Riparia X Rupestris, No. 3369	30	15	15				t					
Do.....					21	.4800	t					
Rupestris Martin.....			8		24	.5325	t					
Salt Creek.....			4		20	.3175	t					
Taylor Narbonne.....			13				t					
Folle Blanche:												
Rupestris St. George.....	20	15		3	21.7	.8175	l					
Foster:												
Own roots.....	25		4		26	.4612	t					
Lenoir.....	6	10	4	7	23.5	.5119	t					
Riparia Gloire.....	20	7	3	8	25.4	.5535	t					
Rupestris St. George.....	20	12	8	15	25.6	.5920	t					
Frankenthal Precoce:												
Herbemont.....												
Lenoir.....	15	2	4		24.1	.7350	l					
Monticola X Riparia, No. 18808		3	3	15	23	.6425	l					
Monticola X Riparia, No. 18815		4	4	12	24.5	.6600	l					
Rupestris St. George.....	20	1	6		25.4	.6487	l					
Solonis X Othello.....			4	8	28	.7125	l					
Solonis X Riparia, No. 1616			4	2	20.5	.6900	l					
Fredonction:												
Dog Ridge.....												
Lenoir.....	8		7		21.8	.8025	l					
Rupestris St. George.....												
Fresa de Monferat:												
Own roots.....				1	26	.8925	l					
Gamay de Bourgogne:												
Own roots.....				1	23	.8025	l					
Dog Ridge.....	25		9	4	25.9	.9525	l					
Lenoir.....	20		6		24.8	1.0762	l					
Rupestris St. George.....	15	3		2	25.7	.9712	l					
Gamay Teinturier:												
Own roots.....				1	28	.6581	l					
Dog Ridge.....	50	3	9	15	26.6	.7912	l					
Lenoir.....	30	8	12	1	24.7	.8500	l					
Rupestris St. George.....	50	8	12	1	26.9	.9225	l					
Taylor Narbonne.....	40	2			28.1	.7420	l					
Genyüz Tramminer:												
Own roots.....				1	24.9	.5625	l					

Riparis × Rupestris, No. 101.	40	5	8	1	18	20.6	.7185	m-l	t	l	l	r	w
Riparis × Rupestris, No. 3306.		10	3		5	19.5	.7725	m-l	t	l	l	r	w
Riparis × Rupestris, No. 3309.					5	21.3	.7325	m-l	t	l	l	r	w
Rupestris Martin.									t	l	l	r	w
Rupestris Mission.					4	22	.8100	m-l	t	l	l	r	w
Rupestris St. George.	30	4	4	4	2	19	.7435	m-l	t	l	l	r	w
Salt Creek.		12			4	20.4	.6470	m-l	t	l	l	r	w
Solonis Robusta.			4			21.5	.8212	m-l	t	l	l	r	w
Solonis × Othello, No. 1613.	25	10	3		5	21	.8137	m-l	t	l	l	r	w
Solonis × Riparis, No. 1616.	50	10	3			17.8	.7575	m-l	t	l	l	r	w
Taylor Narbonne.		10	11	4	4	19.6	.7290	m-l	t	l	l	r	w
Grenache:									t	l	l	r	w
Own roots.	10	30	20	32	30	24	.7245	m-l	t	c	m	ob	l, w
Dog Ridge.	40	12	15	2		22.5	.8794	m-l	t	c	m	ob	l, w
Lenoir.	50	15	18	11	36	22.8	.7800	m-l	t	c	m	ob	l, w
Rupestris St. George.	40	3	4	3	23	24	.9075	m-l	t	c	m	ob	l, w
Taylor Narbonne.	50	5	11		2	24.8	.7170	m-l	t	c	m	ob	l, w
Gros Blanc de Lausanne:									t	c	m	ob	l, w
Lenoir.	20	8	11		17	22.2	.5250	m	t	c	m	ob	w
Solonis × Othello.		2				23.3	.4950	m	t	c	m	ob	w
Gros Guillaume:					1½	28	.5250	m	r	c	l	r	l, s
Lenoir.						25.6	.5091	m	cy	l	m	r	w
Gros Manzenc:	½		3	8		21.5	.8250	m	t	c	l	ob	w
Own roots.									t	c	l	ob	s, t, st
Own roots.									t	c	l	ob	w
Gros Verdor:									t	c	l	ob	w
Own roots.	35	20		35½	12	21	.9000	m	cy	c	l	r	w
Dog Ridge.	20	3				23.5	.8118	m	cy	c	l	r	w
Do.	40	15	5		31	20.7	1.0075	m	cy	c	l	r	w
Lenoir.	12	12	12	4	2	24.1	.7181	m	cy	c	l	r	w
Do.	25	15		20	15	23.8	.8081	m	cy	c	l	r	w
Rupestris St. George.	20	10	14	6	38	22.5	.8445	m	cy	c	l	r	w
Do.	20	5	9			22	.7108	m	cy	c	l	r	w
Taylor Narbonne.									t	l	m	r	w
Hebron:	3	5	8	2		22.3	.5895	m-l	t	l	m	r	w
Lenoir.						21	.7875	m-l	t	l	m	r	w
Rupestris St. George.					1½	20	.7300	m-l	t	l	m	r	w
Solonis × Othello, No. 1613.									t	l	m	r	w
Huasco:									t	l	m	r	w
Own roots.	15	6	2	4		26	.6196	m	t	l	l	o	w
Aramon × Rupestris Ganzin, No. 1.		10	5	3	2	26	.3265	m	t	l	l	o	w
Do.			10			22	.6450	m	t	l	l	o	w
Aramon × Rupestris Ganzin, No. 2.				12		28	.5550	m	t	l	l	o	w
Australis.			4			25	.9850	m	t	l	l	o	w
Berlandieri × Riparis, No. 420A.			3	5		29.5	.4625	m	t	l	l	o	w
Lenoir.	10	8	3	3½		26.5	.6277	m	t	l	l	o	w
Monticola × Riparis, No. 18808.			4			30	.6150	m	t	l	l	o	w
Mourvedre × Rupestris, No. 1202.	15	6	3	19	3	26	.5415	m	t	l	l	o	w
Riparia Gloire.			3	12	1	26	.5975	m	t	l	l	o	w
Riparia × (Cordifolia × Rupestris, No. 106-8).			6			25.5	.5025	m	t	l	l	o	w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling °s	Acid, as tartaric (c.c.) per 100	Cluster.			Berry.			Usc.				
	1909	1910	1911	1912	1913			21	22	23	24	25	26		27	28	29	30
Huasco—Continued.																		
Riparia X Rupestris, No. 101	4		1	2				0.5717	m	t	l	l	0	w	s, t, r			
Do.		2					29.9	.6225	m	t	l	l	0	w	s, t, r			
Riparia X Rupestris, No. 3309.	15	11	3	2			26.3	.5262	m	t	l	l	0	w	s, t, r			
Rupestris St. George.			5	8	4		24.5	.6000	m	t	l	l	0	w	s, t, r			
Solonis Robusta.			5	5			27	.5500	m	t	l	l	0	w	s, t, r			
Solonis X Riparia, No. 1616.			5				29	.4950	m	t	l	l	0	w	s, t, r			
Taylor Narbonne.		5					28	.6750	m	t	l	l	0	w	s, t, r			
Viala.				8			27.2	.6100	m	t	l	l	0	w	s, t, r			
Hunisa:							27		m	t	l	l	0	w	s, t, r			
Adobe Giant.			2				28	.5175	l	cy	l	l	0	b	s, t, st			
Australis.			12	35			20	.7725	l	cy	l	l	0	b	s, t, st			
Berlandieri X Riparia, No. 420A				15			21	.5325	l	cy	l	l	0	b	s, t, st			
Constantia.			12	22			21		l	cy	l	l	0	b	s, t, st			
Dog Ridge.									l	cy	l	l	0	b	s, t, st			
Riparia X Rupestris, No. 3309.									l	cy	l	l	0	b	s, t, st			
Solonis Robusta.									l	cy	l	l	0	b	s, t, st			
Solonis X Riparia, No. 1616.			14	12			21	.5775	l	cy	l	l	0	b	s, t, st			
Hunisa (S. P. I. No. 6124):									l	cy	l	l	0	b	s, t, st			
Own roots:									l	cy	l	l	0	b	s, t, st			
Hunisa (S. P. I. No. 8583):							18.2	1.0125	l	cy	l	l	0	b	s, t, st			
Own roots:	10								l	cy	l	l	0	b	s, t, st			
Hutab:									m	l	l	m	0	w	t			
Own roots:							22	.5850	m	l	l	m	0	w	t			
Hycales:									l	t	l	m	r	w	s, t, w			
Lenoir.	15	10	13			9	22	.6850	l	t	l	m	r	w	s, t, w			
Rupestris Mission.									l	t	l	m	r	w	s, t, w			
Rupestris St. George.		2	19				23	.4815	l	t	l	m	r	w	s, t, w			
Solonis X Othello, No. 1613.			9	12			25.2	.4762	l	t	l	m	r	w	s, t, w			
Imperial:									l	t	l	m	r	w	s, t, w			
Own roots:				8		5	20.5	.6442	m	t	c	m-l	r	b	t, s, w			

Inzolia Bianca:	8	5	5	2	23.1	.6267	m-1	t	c	m	ob	w	t, w
Lenoir.....	1½	10	10	23	.6930	m-1	t	c	m	ob	w	t, w
Rupestris St. George.....	2	4	4	15	25.2	.6562	s	cy	c	s	r	w	w
Johannisberger:	11½	5	1	2	26.5	.6555	m	cy	c	s	r	r	t, w
Ow'n roots.....	12	2	28	6	27.7	.6400	m	cy	c	s	r	r	t, w
Lenoir.....	16	6	3	20	28.8	.7095	m	cy	c	s	r	r	t, w
Riparia Gloire.....	30	17.6	.6712	m	t	c	m-1	ob	w	t, s
Rupestris St. George.....	2	35	8	35	22.7	.5850	l	cy	c	s	r	b	w
Keechumsh-Aly-Blanc:	40	30	10	30	23.4	.6644	l	cy	c	s	r	b	w
Ow'n roots.....	15	30	4	2	23.7	.4950	l	cy	c	s	r	b	w
Lenoir.....	8	15	5	18	23.9	.5365	l	cy	c	s	r	b	w
Riparia Gloire.....	20	40	10	36	24.8	.4305	l	cy	c	s	r	b	w
Rupestris St. George.....	2	2½	24.1	.7425	l	t	l	m	o	w	t, f, w
Keechumsh-Aly-Blanc:	20	5	11	4	24	.7652	l	t	l	m	o	w	t, f, w
Ow'n roots.....	20	3	5	35	21.4	.5796	l	t	c	l	r	b	s, t, w
Lenoir.....	4	2	1	½	24.9	.6262	s	cy	c	s	r	w	w
Riparia × Rupestris, No. 3309.....	3	1	10	4	25.3	.6510	m	cy	l	l	r	b	w
Herbement.....	15	1½	10	23	.8717	m	cy	l	l	r	b	w
Lenoir.....	15	8	9½	4	22.4	.7765	m	cy	l	l	r	b	w
Klemburger:	45	7	12	1	21.2	.9210	m	cy	l	l	r	b	w
Ow'n roots.....	15	2	24.3	.4395	m	cy	l	l	r	b	w
Köbner:	3	6	22.1	.5550	m	t	l	l	ob	r	s, t
Riparia × Rupestris, No. 3309.....	50	4	21.2	.5250	l	cy	l	m	ob	w	w
Herbement.....	15	7	20.3	.7050	l	cy	l	m	ob	w	w
Lenoir.....	15	8	1½	23.3	.7037	m-1	l	l	l	ob	w	w, s, t
Klemburger:	30	10	8	6	26	.6944	m-1	l	l	l	ob	w	w, s, t
Ow'n roots.....	30	40	8	30	24.3	.7350	m-1	l	l	l	ob	w	w, s, t
Köbner:	30	40	8	18	22.3	.5130	m	cy	c	s	r	w	w
Riparia Gloire.....	40	30	5	3	27.5	.8475	s	cy	c	s	r	w	w
Rupestris St. George.....	10	6	0	22.3	.6094	s	cy	c	s	r	w	w
Léani Zolo:	42	35	14	47	22.6	.6187	s	cy	c	s	r	w	w
Lig'nian Blanc:	42	35	14	47	22.6	.6388	s	cy	c	s	r	w	w
Ow'n roots.....	40	30	5	43	27.5	.8475	s	cy	c	s	r	w	w
Aramon × Rupestris Ganzin, No. 1.....	10	6	0	22.6	.6187	s	cy	c	s	r	w	w
Dog Ridge.....	42	35	14	47	22.6	.6388	s	cy	c	s	r	w	w
Do.....	42	35	14	47	22.6	.6388	s	cy	c	s	r	w	w

Key:

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tartaric c. c. per 100	Cluster.			Berry.			Use.
	1909	1910	1911	1912	1913			Size.	Shape.	Compact or loose.	Size.	Shape.	Color.	
Lignan Blanc—Continued.														
Hotspur.....	25	5	3	11	5	24.1	.6175	S	CY	C	I	W	W	W
Lenor.....	28	10	2	2	4	23.2	.6660	S	CY	C	I	W	W	W
Do.....	28	10	2	2	4	21.3	.6940	S	CY	C	I	W	W	W
Mourvedre X Rupestris, No. 1202.....	25	30	3	41	8	22.3	.6315	S	CY	C	I	W	W	W
Do.....	8	35	8	40	11	24.8	.6615	S	CY	C	I	W	W	W
Riparia Gloire.....	30	30	20	6	0	22.9	.6276	S	CY	C	I	W	W	W
Riparia X Rupestris, No. 101.....	20	50	25	13	4	23	.6135	S	CY	C	I	W	W	W
Riparia X Rupestris, No. 3309.....	40	10	10	10	10 ¹	22.8	.6350	S	CY	C	I	W	W	W
Rupestris St. George.....	40	30	5	18	26	22.4	.6185	S	CY	C	I	W	W	W
Do.....	10	20	4	12	0	1.0462	1.0462	S	CY	C	I	W	W	W
Solonis X Othello.....	10	20	15	10	0	24	.5174	S	CY	C	I	W	W	W
Solonis X Riparia, No. 1616.....	6	40	10	23	0	22.4	.5681	S	CY	C	I	W	W	W
Taylor Narbonne.....	50	5	3	24.2	.6150	S	CY	C	I	W	W	W
Do.....	10	10	15	21	8	23.6	.5090	I	I	I	I	W	W, S, t	W, S, t
Lislan:														
Own roots.....	10	10	15	21	8	23.6	.5090	I	I	I	I	W	W, S, t	W, S, t
(Aestivalls X Monticola) X (Riparia X Rupestris, No. 554-9).....	40	40	26	6	1	23	.6300	I	I	C	I	W	W, S, t	W, S, t
Aramon X Rupestris Ganzin, No. 1.....	40	40	26	6	1	23.3	.5806	I	I	C	I	W	W, S, t	W, S, t
Aramon X Rupestris Ganzin, No. 2.....	40	40	26	6	1	23.3	.5806	I	I	C	I	W	W, S, t	W, S, t
Berklandier X Riparia, No. 4203.....	5	5	5	5	2	23.7	.6187	I	I	C	I	W	W, S, t	W, S, t
Columband X Riparia, No. 2502.....	5	5	5	5	2	23.7	.6187	I	I	C	I	W	W, S, t	W, S, t
Constantia.....	4	4	4	4	3	22	.5025	I	I	C	I	W	W, S, t	W, S, t
Cornucopia.....	4	4	4	4	2	23.8	.6375	I	I	C	I	W	W, S, t	W, S, t
Do.....	4	4	4	4	2	23.3	.3475	I	I	C	I	W	W, S, t	W, S, t
De Grasset.....	4	4	4	4	2	23.3	.3475	I	I	C	I	W	W, S, t	W, S, t
Dog Ridge.....	30	10	17	6	8	25.6	.4950	I	I	C	I	W	W, S, t	W, S, t
Herbement.....	15	12	2	4	12	23.3	.6375	I	I	C	I	W	W, S, t	W, S, t
Lenor.....	6	5	9	2	4	22.3	.5344	I	I	C	I	W	W, S, t	W, S, t
Monticola X Riparia, No. 18804.....	6	5	9	2	4	23.1	.5450	I	I	C	I	W	W, S, t	W, S, t
Monticola X Riparia, No. 18808.....	6	5	9	2	4	23.1	.5450	I	I	C	I	W	W, S, t	W, S, t
Mourvedre X Rupestris, No. 1202.....	30	10	13	2	15	23.7	.6070	I	I	C	I	W	W, S, t	W, S, t
Do.....	30	10	13	2	15	24.5	.6070	I	I	C	I	W	W, S, t	W, S, t
Do.....	30	10	13	2	15	22.9	.6530	I	I	C	I	W	W, S, t	W, S, t

Var. Name	25	10	16	2	2	22.3	6300	l	l	c	m	r	w	s	t
Riparia X Rupestris, No. 101						21	6300	l	l	c	m	r	w	s	t
Do						24	6300	l	l	c	m	r	w	s	t
Riparia X Rupestris, No. 101-14						24	6487	l	l	c	m	r	w	s	t
Riparia X Rupestris, No. 3306					5	24	6750	l	l	c	m	r	w	s	t
Riparia X Rupestris, No. 3309				19		23	6255	l	l	c	m	r	w	s	t
Do						23	6660	l	l	c	m	r	w	s	t
Rupestris Martin	60	15	9		5	23.3		l	l	c	m	r	w	s	t
Rupestris Metallien	30	10	14		8	24	6287	l	l	c	m	r	w	s	t
Rupestris Mission					3	25	6112	l	l	c	m	r	w	s	t
Rupestris St. George					2	25	5475	l	l	c	m	r	w	s	t
Do	6	3	2		1	23.7	5169	l	l	c	m	r	w	s	t
Rupestris X Berlandieri, No. 219A				1 1/2		23	5250	l	l	c	m	r	w	s	t
Salt Creek	15	8	5		10	25	6075	l	l	c	m	r	w	s	t
Solanis Robusta		6	4			24.5	4575	l	l	c	m	r	w	s	t
Do			5		4	23.3	6800	l	l	c	m	r	w	s	t
Solanis X Otello			1			23	4537	l	l	c	m	r	w	s	t
Solanis X Otello, No. 1013			1 1/2		2	24	6450	l	l	c	m	r	w	s	t
Solanis X Riparia, No. 1016	15	50	10		3	24	6469	l	l	c	m	r	w	s	t
Taylor Narbonne	30	10	25		3	24.6	5815	l	l	c	m	r	w	s	t
Valencia		3	3		8	21.2	4825	l	l	c	m	r	w	s	t
Lughecca Neri								l	l	c	m	r	w	s	t
Adobe Giant	30	10	4	26		22.4	5031	l	l	c	m	r	w	s	t
Dog Ridge	45	25		6	8	24	5694	l	l	c	m	r	w	s	t
Riparia X Rupestris, No. 3309				2		24	6825	l	l	c	m	r	w	s	t
Rupestris St. George	20	25	5	43	6	24	4740	l	l	c	m	r	w	s	t
Macmer								l	l	c	m	r	w	s	t
Own roots	1 1/2			4	1	25	6917	s		c	m	r	w	s	t
Madeleine Angevine								l	l	c	m	r	w	s	t
Own roots		3			2	22.6	4950	l	l	c	m	r	w	s	t
Dog Ridge	15	2	12		1	24.5	5392	l	l	c	m	r	w	s	t
Lenoir	15	2	1		2 1/2	22.3	5120	l	l	c	m	r	w	s	t
Rupestris St. George	6	2	2		3	21.7	5756	l	l	c	m	r	w	s	t
Madeleine Blanche								l	l	c	m	r	w	s	t
Own roots						25.9	6975	l	l	c	m	r	w	s	t
Madeleine d'Ambre								l	l	c	m	r	w	s	t
Dog Ridge	6	2	16		1/2	23.5	7800	l	l	c	m	r	w	s	t
Lenoir	3	2	3		5	20.9	7294	l	l	c	m	r	w	s	t
Rupestris St. George					3	21.3	8057	l	l	c	m	r	w	s	t
Madeleine Rose								l	l	c	m	r	w	s	t
Own roots						22	7500	l	l	c	m	r	w	s	t
Riparia Gloire			2		2	25.5	5587	l	l	c	m	r	w	s	t
Madeleine Royale								l	l	c	m	r	w	s	t
Own roots	25	4			4	31	7500	l	l	c	m	r	w	s	t
Dog Ridge	10	2			3	27.4	5725	l	l	c	m	r	w	s	t
Lenoir			3		3	25.7	6600	l	l	c	m	r	w	s	t
Mahaia								l	l	c	m	r	w	s	t
Own roots	20	25	2	7	9	23.9	5205	l	l	c	m	r	w	s	t
Adobe Giant	60	70	15	42	30	22.6	5550	l	l	c	m	r	w	s	t
Aramon X Rupestris Ganzin, No. 1	50	40	3	43	8	22.7	6045	l	l	c	m	r	w	s	t
Canada								l	l	c	m	r	w	s	t
Constantia						23	5775	l	l	c	m	r	w	s	t

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, scale, <i>P. cent.</i>	Acid, as tartaric (C.C.) per 100	Size.	Cluster.		Berry.			Use.
	1909	1910	1911	1912	1913				Shape.	Compact or loose.	Size.	Shape.	Color.	
Malaga—Continued.														
Dog Ridge.....	60	40	10	20	32	23.1	0.6150	l	t	m	l	ob	w	s, t, r
Herbmont.....	2	20	2	4	24.8	0.6933	l	t	m	l	ob	w	s, t, r
Lenoir.....	30	20	20	4	2	22.8	1.392	l	t	m	l	ob	w	s, t, r
Mourvedro X Rupestris, No. 1202.....	45	30	15	45	12	22.9	5.650	l	t	m	l	ob	w	s, t, r
Riparia Gloire.....	50	25	15	44	1.8	22.7	6.189	l	t	m	l	ob	w	s, t, r
Riparia X Rupestris, No. 101.....	40	70	20	43	10	23.6	6.120	l	t	m	l	ob	w	s, t, r
Riparia X Rupestris, No. 3309.....	40	30	30	22	1.8	23.7	5.995	l	t	m	l	ob	w	s, t, r
Rupestris Martin.....	11	7	11	11	22.5	1.790	l	t	m	l	ob	w	s, t, r
Rupestris St. George.....	40	30	15	30	20	22.9	5.715	l	t	m	l	ob	w	s, t, r
Salt Creek.....	30	15	5	31	8	22.8	6.165	l	t	m	l	ob	w	s, t, r
Solonis X Riparia, No. 1616.....	50	70	35	22	12	22.7	6.006	l	t	m	l	ob	w	s, t, r
Taylor Narbonne.....	60	60	15	33	4	23.9	5.600	l	t	m	l	ob	w	s, t, r
Malaga Blanc:														
Lenoir.....	15	18	3	26.9	5.125	m-l	l	l	m	ob	w	w
Riparia Gloire.....	4	15	1	7	4	25.3	5.790	m-l	l	l	m	ob	w	w
Rupestris St. George.....	15	25	4	30	26	26.5	5.720	m-l	cy	c	s	r	d	w
Malbeck:
Own roots.....	2	2	22	3	24	8.025	s	t	c	l	ob	b	s, t, w
Malvasia:
Own roots.....	2	6	2	27	26	5.556	s	t	c	l	ob	b	s, t, w
Lenoir.....	s	t	c	l	ob	b	s, t, w
Rupestris St. George.....	3	24	9.375	s	t	c	l	ob	b	s, t, w
Malvasia de Broglio:
Own roots.....	4	15	23.1	6.375	l	t	l	m	r	w	w
Malvasia Rosaria:
Own roots.....	15	8	2	12	24.5	7.162	l	t	c	l	ob	b	w
Dog Ridge.....	40	50	5	40	0	23.4	7.125	l	t	c	l	ob	b	w
Herbmont.....	21	8.025	l	t	c	l	ob	b	w
Lenoir.....	25	15	4 ¹	6	24.3	6.354	l	t	c	l	ob	b	w
Riparia Gloire.....	30	40	5	20	26	22.4	6.995	l	t	c	l	ob	b	w
Rupestris St. George.....	25	25	10	10	0	24.1	6.669	l	t	c	l	ob	b	w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tartaric (c.c.) per 100	Cluster.			Berry.			Use.
	1909							Size.	Shape.	Compact or loose.	Size.	Shape.	Color.	
	1910	1911	1912	1913	1913									
I	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Malaria—Continued.						<i>P. cent.</i>								
Mourvedre × Rupestris, No. 1202.....	50	5	6	4	21.1	0.8587	m	cy	c	m	r	b	w
Rupestris Martin.....	40	8	2	16	21.4	.8212	m	cy	c	m	r	b	w
Rupestris St. George.....	50	12	15	10	21.7	.7200	m	cy	c	m	r	b	w
Salt Creek.....	45	12	4	4	22	.8719	m	cy	c	m	r	b	w
Taylor Narbonne.....	30	10	8	10	4	22.6	.7350	m	cy	c	m	r	b	w
Melon:														
Dog Ridge.....	20	20	10	15	23.1	.5131	m	r	c	l	ob	w	s, t
Rupestris St. George.....	15	15	15	8	23.5	.5096	m	r	c	l	ob	w	s, t
Merlot: Own roots.....	2	2	3	24.7	.5950	m	cy	m	s	r	b	w
Meslier: Own roots.....	3	1½	1	24.7	.9800	s	cy	c	s	r	w	w
Aramon × Rupestris Ganzin, No. 1.....	6	5	22.4	1.0650	s	cy	c	s	r	w	w
Lenoir.....	8	3	4	20	21.7	1.0918	s	cy	c	s	r	w	w
Rupestris St. George.....	15	1	5½	1	6	22.8	.9224	s	cy	c	s	r	w	w
Meunier: Own roots.....	4	2	2	28.4	.6975	s	cy	c	s	r	b	w
Dog Ridge.....	25	1	16	25.1	.7050	s	cy	c	s	r	b	w
Lenoir.....	20	3	5	9	21.5	.8344	s	cy	c	s	r	b	w
Rupestris St. George.....	40	3	29	3	7	28.4	.8160	s	cy	c	s	r	b	w
Meyer, No. 59: Dog Ridge.....	12	5	25.8	.5250	l	t, cy	c	m	ob	w	s, t, w
Lenoir.....	6	22	.5825	l	t, cy	c	m	ob	w	s, t, w
Meyer, No. 60: Dog Ridge.....	30	3½	25.1	.6112	l	t, cy	c	m	ob	w	s, t, w
Lenoir.....	1	4	22	.7349	l	t, cy	c	m	ob	w	s, t, w
Meyer, No. 65: Lenoir.....	20	8	13	19	20.6	.6885	l	t	m	m	r	r	w
Meyer, No. 94: Dog Ridge.....	20	21.2	.2400	m-1	cy	m	l	ob	w	s, t
Lenoir.....	2	11	20.2	.3316	m-1	cy	m	l	ob	w	s, t

Year	Region	Area	Yield	Quality	Notes
10	Meyer, No. 95:	2	21.1	.6150	S, t
20	Dog Ridge	3	19	.5875	S, t
	Lenoir	13			W
	Meyer, No. 103:		23	.6000	W
	Lenoir	5			b
	Meyer, No. 107:		19	.8637	W
	Lenoir	9			b
	Meyer, No. 116:		23.2	.6300	W
	Dog Ridge	4	24.1	.5925	W
	Lenoir				W
	Meyer, No. 515:		20.1	.7425	W
	Dog Ridge	5½	20.6	.7275	W
	Lenoir				W
	Millennium:				S, t, w
	Aramon X Rupestris Ganzin, No. 1:		23.7	.6885	S, t, w
	Lenoir	11	4	.7237	S, t, w
	Riparia X Rupestris, No. 101:	11	6	.5950	S, t, w
	Riparia X Rupestris, No. 3309:	2	21.9	.7125	S, t, w
	Solonis X Riparia, No. 1615:	1	22.3	.6825	S, t, w
	Mission:				S, t, w
	Own roots:		20.2	.5885	W, S, t
	Adobe Giant:	10	15	.5010	W, S, t
	Dog Ridge:	4	26	.4889	W, S, t
	Do:	5	15	.7500	W, S, t
	Herbemont:	20	21.2	.4837	W, S, t
	Do:	4	29.4	.7295	W, S, t
	Lenoir:	20	12	.5070	W, S, t
	Do:	30	3	.6125	W, S, t
	Riparia Gloire:	15	10	.4871	W, S, t
	Rupestris Mission:	10	2	.4605	W, S, t
	Do:	5	10	.7395	W, S, t
	Rupestris St. George:	30	18	.5450	W, S, t
	Do:	40	12	.5910	W, S, t
	Salt Creek:	20	15	.5685	W, S, t
	Do:	30	16	.6412	W, S, t
	Molineria Cordo:				S, t
	Lenoir:	3	23	.5175	W
	Mondeuse:				W
	Own roots:		25.4	.6170	W
	Aramon X Rupestris Ganzin, No. 1:	4	6	.7781	W
	Dog Ridge:	15	4	.9430	W
	Herbemont:	8	3	.6895	W
	Do:	20	7	.8930	W
	Lenoir:	20	10	.8125	W
	Mourvedre X Rupestris, No. 1202:	10	5	.8750	W
	Riparia X Rupestris, No. 101:	40	12	.7200	W
	Riparia X Rupestris, No. 3309:	40	6	.8220	W
	Rupestris Martin:	50	5	.7600	W
	Rupestris St. George:	5	15	.7700	W
	Salt Creek:	10	6	.8231	W
	Solonis X Riparia, No. 1616:	25	12		W
	Taylor Narbonne:	25	5		W
	Do:	10	13		W

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Baillings scale, <i>P. cent.</i>	Acid, as tartaric c. c.). (grams per 100)	Cluster.			Berry.			Use.
								Size.	Shape.	Compact or loose.	Size.	Shape.	Color.	
	1909	1910	1911	1912	1913									
I	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Monaca Nero: Own roots.....	5	10	4	18	23.8	0.5700	l	t	c	m	r	b	w, t
Mourvada: Own roots.....	3	3	2	8	6	20.2	.8235	m-1	t	m	m	r	b	w
Aramon X Rupestris Ganzin, No. 1.....	40	10	19	12	12	19.6	1.0540	m-1	t	m	m	r	b	w
Dog Ridge.....	23.1	1.0125	m-1	t	m	m	r	b	w
Herbemont.....	20	.8450	m-1	t	m	m	r	b	w
Lenoir.....	24	.8250	m-1	t	m	m	r	b	w
Riparia X Rupestris, No. 101.....	60	8	15	4	31	22.5	.9320	m-1	t	m	m	r	b	w
Riparia X Rupestris, No. 3309.....	60	5	12	3	15	22.2	.9825	m-1	t	m	m	r	b	w
Solonis X Riparia, No. 1616.....	40	25	11	3	3	21.7	.9855	m-1	t	m	m	r	b	w
Taylor Narbonne.....	60	10	11	12	22.2	.9622	m-1	t	m	m	r	b	w
Mourisco Bisanca: Own roots.....	10	6	6	18	23	.5600	l	t	l	m	ob	w	t, s, w
Dog Ridge.....	50	10	21	.4450	l	t	l	m	ob	w	t, s, w
Lenoir.....	60	10	40	19.4	.6725	l	t	l	m	ob	w	t, s, w
Rupestris St. George.....	25	12	18	30	24.4	.5750	l	t	l	m	ob	w	t, s, w
Taylor Narbonne.....	40	2	6	23.4	.5300	l	t	l	m	ob	w	t, s, w
Mourisco Preto: Own roots.....	12	10	2	3	25.7	.5662	m	t	c	s	o	b	w
Dog Ridge.....	50	6	10	8	24.8	.7637	m	t	c	s	o	b	w
Lenoir.....	40	8	2	21	23.3	.8401	m	t	c	s	o	b	w
Rupestris St. George.....	6	24.7	.7516	m	t	c	s	o	b	w
Muscadelle du Bordelais: Adobe Giant.....	10	4	2	9	4	25.4	.6615	m-1	l	m	m	r	w	w
Aramon X Rupestris Ganzin, No. 1.....	12	8	2	2	2	27.3	.5420	m-1	l	m	m	r	w	w
Lenoir.....	3	5	2	3	3	26.5	.6445	m-1	l	m	m	r	w	w
Mourvedre X Rupestris, No. 1202.....	15	5	5	8	12	28.9	.5885	m-1	l	m	m	r	w	w
Riparia Gloire.....	1	4	5	8	4	28.8	.6355	m-1	l	m	m	r	w	w
Riparia X Rupestris, No. 101.....	20	4	7	8	16	28.2	.5730	m-1	l	m	m	r	w	w
Riparia X Rupestris, No. 3309.....	20	2	8	19	15	27.7	.5750	m-1	l	m	m	r	w	w
Rupestris St. George.....	25	6	10	8	4	28.2	.6245	m-1	l	m	m	r	w	w
Solonis X Riparia, No. 1616.....	10	0	3	30.5	.5772	m-1	l	m	m	r	w	w
Taylor Narbonne.....	10	5	8	32.1	.6450	m-1	l	m	m	r	w	w

	15	1	1	5	2	16.2	.6000	m-l	t, cy	m	ob	b	s, t
Muscad (S. P. I. No. 3063):													
Own roots.....													
Muscad Albardiens:													
Own roots.....													
Aramon X Rupestris Ganzin, No. 1.....	4	8	1	5	2	23.2	.7569	m-l	1	m	ob	b	s, t
Lenoir.....	8	3	3	1 ¹ / ₂	2	27	.5225	m-l	1	m	ob	b	s, t
Mourvedre X Rupestris, No. 1202.....	12	10	3	18	20	24.9	.5125	m-l	1	m	ob	b	s, t
Riparia Gloire.....	25	6	2	32	1	26.6	.4940	m-l	1	m	ob	b	s, t
Riparia X Rupestris, No. 101.....	8	12	1	21	6	26.2	.5580	m-l	1	m	ob	b	s, t
Riparia X Rupestris, No. 3309.....	30	0	10	13	8	26.4	.5060	m-l	1	m	ob	b	s, t
Rupestris St. George.....	15	5	8	8	6	28	.4462	m-l	1	m	ob	b	s, t
Solonis X Riparia, No. 1616.....	5	0	1	2	27.1	.4375	m-l	1	m	ob	b	s, t
Taylor Narbonne.....	6	4	5	18	24.7	.4312	m-l	1	m	ob	b	s, t
Muscad Bonod:													
Berlandieri X Riparia, No. 420A.....			13	23	.7500	m-l	t	1	o	w	s, t
De Grasset.....			3	30	.4650	m	t	1	o	w	s, t
Lenoir.....	10	2	9 ¹ / ₂	2	22.3	.6094	m-l	t	1	o	w	s, t
Riparia X Rupestris, No. 101.....			14	28	.5750	m-l	t	1	o	w	s, t
Riparia X Rupestris, No. 3309.....			31	.5475	m-l	t	1	o	w	s, t
Rupestris St. George.....			25	.6750	m-l	t	1	o	w	s, t
Solonis Robusta.....			8	27	.8925	m-l	t	1	o	w	s, t
Valencia.....			3	28	.5175	m-l	t	1	o	w	s, t
Muscad Capusines:													
Own roots.....													
Dog Ridge.....	15					27.2	.8250	1	t	1	o	b	s, t
Lenoir.....	40		6			25.3	.7175	1	t	1	o	b	s, t
Rupestris St. George.....	40		11		4	24.3	.7575	1	t	1	o	b	s, t
Muscadall:													
Own roots.....													
Dog Ridge.....	8	5	1	12	6	25.4	.6525	m	cy	c	r	w	s, w
Lenoir.....	4	4	15	9	2	28.3	.6555	m	cy	c	r	w	s, w
Riparia Gloire.....	1	4	1	1	2	25.2	.7080	m	cy	c	r	w	s, w
Rupestris St. George.....	4	3	1	1	25.5	.6200	m	cy	c	r	w	s, w
Muscad Gros Noir Hadri:	20	3	1	6	10	30.1	.5790	m	cy	c	r	w	s, w
Muscad Hamburg:													
Own roots.....													
Dog Ridge.....	3	2	3	4	23.7	.6350	m-l	1	1	o	b	s, t, w
Lenoir.....	25	5	5	1	4	24.4	.7540	m-l	1	1	o	b	s, t, w
Rupestris St. George.....	12	5	7	23.3	.8075	m-l	1	1	o	b	s, t, w
Taylor Narbonne.....	30	5	2	24.4	.8525	m-l	1	1	o	b	s, t, w
Muscad Hamburg:													
Own roots.....													
Aramon X Rupestris Ganzin, No. 1.....			3	26.7	.5550	m	1	1	o	b	s, t, w
Dog Ridge.....	30	1 ¹ / ₂	4	27	.5775	m	1	1	o	b	s, t, w
Lenoir.....	40	2	10	3	2	24.5	.7080	m	1	1	o	b	s, t, w
Monticola X Rupestris.....			4	23	.6285	m	1	1	o	b	s, t, w
Riparia X Rupestris, No. 3309.....			4	28.2	.6075	m	1	1	o	b	s, t, w
Rupestris St. George.....			2	24.5	.6050	m	1	1	o	b	s, t, w
Rupestris X Berlandieri, No. 219A.....	25	12	13	22.7	.6150	m	1	1	o	b	s, t, w
Taylor Narbonne.....	15	3	2 ¹ / ₂	26.5	.5925	m	1	1	o	b	s, t, w
Muscad Hamburg Noir d'Hongrie:			5	23.8	.6025	m	1	1	o	b	s, t, w
Own roots.....			2	23.8	.6918	m	1	1	o	b	s, t, w

	1	5	10	3	2	22	.4575	22	24.6	.4706	m	m	t	m	l	l	r	r	w	w	S ₁ , t, w	S ₂ , t, w	
Nasa Vaientiana:																							
Own roots.....	10	10	10	3	2	22	.4706	22	24.6	.4706	m	m	t	m	l	l	r	r	w	w	S ₁ , t, w	S ₂ , t, w	
Rupestris St. George.....																							
Nebbiolo:																							
Own roots.....	5	5	5	3½	4	22.7	.7202	22.7	24.4	.7202	m	c	cy	m	m	m	r	r	b	b	w	w	
Dog Ridge.....	40	15	8	8	3½	24.4	.6133	24.4	22.8	.6133	m	c	cy	m	m	m	r	r	b	b	w	w	
Lenoir.....	35	8	12	2	3½	22.8	.7050	22.8	25.3	.7050	m	c	cy	m	m	m	r	r	b	b	w	w	
Rupestris St. George.....	5	4	4	2	6½	25.3	.5400	25.3	24.3	.5400	m	c	cy	m	m	m	r	r	b	b	w	w	
Nebbiolo Bourgt:																							
Dog Ridge.....	10	1½	1½	1	1	24.3	1.0012	24.3	23.5	1.0012	m	c	t	m	m	m	r	r	b	b	w	w	
Lenoir.....	20	5	5	1½	9	23.5	.9632	23.5	23.8	.9632	m	c	t	m	m	m	r	r	b	b	w	w	
Rupestris St. George.....	5	2	2	2	9	23.8	.8666	23.8	24.3	.8666	m	c	t	m	m	m	r	r	b	b	w	w	
Taylor Narbonne.....	5	2	2	2	9	25	.8587	25	23.7	.8587	m	c	t	m	m	m	r	r	b	b	w	w	
Nebbiolo Fino:																							
Dog Ridge.....	12	1½	2½	2½	4	23.7	1.0125	23.7	23.9	1.0125	m	c	t	m	m	m	r	r	b	b	w	w	
Lenoir.....	15	2	2	6	4	23.9	.7987	23.9	24.3	.7987	m	c	t	m	m	m	r	r	b	b	w	w	
Rupestris St. George.....	10	1	1	6	4	24.3	.9262	24.3	26.3	.9262	m	c	t	m	m	m	r	r	b	b	w	w	
Taylor Narbonne.....																							
Negrara di Gattinara:																							
Own roots.....	2	3	3	2	2	26.3	.5550	26.3	27.5	.5550	m	c	lt	m	m	m	r	r	b	b	w	w	
Negro Amaro.....																							
Own roots.....	8	8	3	19	1	27.5	.9950	27.5	23.8	.9950	m	c	t	m	s	s	r	r	b	b	w	w	
Neiretía di Castilla.....																							
Own roots.....	3	2	2	3	2	23.8	.5450	23.8	25.5	.5450	m	m	t	m	m	m	r	r	b	b	w	w	
Oera di Boe.....																							
Own roots.....	1	2	2	15	7	25.5	.7313	25.5	17	.7313	s	c	cy	m	m	m	ob	ob	b	b	w	w	
Ohanez.....																							
Lenoir.....																							
Olivette Blanche:																							
Adops Ghant.....																							
Lenoir.....	12	4	8	32	4	17	.8475	17	19.7	.8475	l	m	t	m	l	l	ob	ob	w	w	S ₁ , t	S ₂ , t	
Riparia à Grandes Feuilles.....																							
Riparia X Rupestris, No. 101.....	3	3	21	21	4	20.1	.7125	20.1	23.6	.7125	l	m	t	m	l	l	ob	ob	w	w	S ₁ , t	S ₂ , t	
Lenoir.....	8	4	4	4	4	23.6	.5325	23.6	22.1	.5325	l	m	t	m	l	l	ob	ob	w	w	S ₁ , t	S ₂ , t	
Solomis Robusta.....																							
Lenoir.....	8	4	4	4	4	22.1	.6825	22.1	19	.6825	l	m	t	m	l	l	ob	ob	w	w	S ₁ , t	S ₂ , t	
Olivette Noir:																							
Lenoir.....	8	2	6½	6½	6½	21	.5775	21	24	.5775	l	m	t	m	l	l	ob	ob	b	b	S ₁ , t	S ₂ , t	
Mourvedre X Rupestris, No. 1202.....																							
Riparia X Rupestris, No. 3309.....																							
Solomis X Ohello, No. 1013.....																							
Lenoir.....	20	5½	5½	3	3	20	.7925	20	21.5	.7925	l	m	t	m	l	l	ob	ob	r	r	S ₁ , t	S ₂ , t	
Opiman.....																							
Lenoir.....	3	3	3	4	4	21.5	.7925	21.5	22	.7925	s	c	cy	m	s	s	r	r	w	w	w	w	
Orleans.....																							
Pagadebilo:																							
Own roots.....	3	10	6	10	1	22	.6112	22	23	.6112	m	m	cy	m	m	m	r	r	b	b	w, s, t	w, s, t	
Dog Ridge.....	40	2	2	1	4	23	.8957	23		.8957	m	m	cy	m	m	m	r	r	b	b	w, s, t	w, s, t	

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).						Sugar, Balling scale.	Acid, as tartaric c. c.).	Cluster.			Berry.			Use.
	1909								Size.	Shape.	Compact or loose.	Size.	Shape.	Color.	
	17	18	19	20	21	1913									
Pagadebito—Continued.															
Lenoir.....	40 ⁴	3	7	10	38		P cont	0.7530	m	cy	m	r	b	w, s, t	
Rupestris St. George.....	35	4	11	2	28		23.6	.8035	m	cy	m	r	b	w, s, t	
Taylor Narbonne.....	15	4	5	2	1		23.4	.6945	m	cy	m	r	b	w, s, t	
Own roots.															
Panariti.....		8	20	30			18.6	.8325	l	cy-t	c	ob	w	s, t	
Adobe Giant.....	3						32	.7735	m-l	cy	c	r	b	r, w, s	
Aramon X Rupestris Ganzin, No. 1.....	10		10				32.8	.7050	m-l	cy	c	r	b	r, w, s	
Dog Ridge.....	3		4				33	.6562	m-l	cy	c	r	b	r, w, s	
Herbement.....									m-l	cy	c	r	b	r, w, s	
Lenoir.....			4				29	.5700	m-l	cy	c	r	b	r, w, s	
Mourvedre X Rupestris, No. 1202.....	2		3				29.8	.6502	m-l	cy	c	r	b	r, w, s	
Riparia Gloire.....	2		3				28.1	.7387	m-l	cy	c	r	b	r, w, s	
Riparia X Rupestris, No. 101.....									m-l	cy	c	r	b	r, w, s	
Riparia X Rupestris, No. 3309.....	3		5				34.8	.6375	m-l	cy	c	r	b	r, w, s	
Rupestris Martin.....									m-l	cy	c	r	b	r, w, s	
Rupestris St. George.....	2		5				34.5	.6337	m-l	cy	c	r	b	r, w, s	
Salt Creek.....	3		5				29	.6000	m-l	cy	c	r	b	r, w, s	
Solons X Riparia, No. 1616.....			5				33.5	.6070	m-l	cy	c	r	b	r, w, s	
Taylor Narbonne.....	3		2				31	.6900	m-l	cy	c	r	b	r, w, s	
Parc de Versailles:															
Adobe Giant.....	10	2	2	27	6		21.5	.5225	m	t	m	ob	w	s, t	
Aramon X Rupestris Ganzin, No. 1.....	2	3	2	3	18		21.5	.5353	m	t	m	ob	w	s, t	
Dog Ridge.....	5	4	2 ¹ / ₂	18 ¹ / ₂	13		19.3	.7450	m	t	m	ob	w	s, t	
Lenoir.....	1 ¹ / ₂	3	2	5	3		21.8	.5662	m	t	m	ob	w	s, t	
Mourvedre X Rupestris, No. 1202.....	1	2	2	5	2		21.9	.5431	m	t	m	ob	w	s, t	
Ponroy.....							20	.6150	m	t	m	ob	w	s, t	
Riparia Gloire.....	6	1	4	41	20		22.2	.4875	m	t	m	ob	w	s, t	
Riparia X Rupestris, No. 101.....	1	4	2	12	9		22.1	.5460	m	t	m	ob	w	s, t	
Riparia X Rupestris, No. 3309.....									m	t	m	ob	w	s, t	

TABLE VII. Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).						Sugar, Baling scale.	Acid, as tartaric e. c. (grams per 100	Cluster.			Berry.			Use.
	1909	1910	1911	1912	1913	1943			Size.	Shape.	Compact or loose.	Size.	Shape.	Color.	
I	17	18	19	20	21										30
Peit Syrah—Continued.							<i>P. cent.</i>								
Boursiquot × Rupestris, No. 109-4.			17	3	12		21	1.1850	m	t, cy	c	m	r	b	w
(Boursiquot × Rupestris, No. 601) × Calicola,															
No. 13206.															
Rupestris × Rupestris, No. 3907			9	3 ¹ / ₂	18 ¹ / ₂		21.8	.7800	m	t, cy	c	m	r	b	w
Cabernet × Berlandieri, No. 333			8	7	9		21.6	.8441	m	t, cy	c	m	r	b	w
Cabernet × Rupestris Ganzan, No. 33A			6	6 ¹ / ₂	10 ¹ / ₂		20.3	.8600	m	t, cy	c	m	r	b	w
Cabernet			2	2	16		1.2833		m	t, cy	c	m	r	b	w
Chasselas × Berlandieri, No. 4113			5	1 ¹ / ₂	19		18	.6750	m	t, cy	c	m	r	b	w
Contancia			4	4	8		21.6	.8275	m	t, cy	c	m	r	b	w
Do.			24	1	8		23	.7725	m	t, cy	c	m	r	b	w
Dog Ridge.			11	5	35		20.6	.8150	m	t, cy	c	m	r	b	w
Do.	50	10	4	6	6		20.2	.8670	m	t, cy	c	m	r	b	w
Herbomont.	25	25	14	2	11		21.3	.8700	m	t, cy	c	m	r	b	w
Hoffman.			6	3	13		18.3	.7975	m	t, cy	c	m	r	b	w
Lady.			1	1	5 ¹ / ₂		21.3	.7517	m	t, cy	c	m	r	b	w
Louise.			30	18	6		23.1	.7365	m	t, cy	c	m	r	b	w
Monticola × Riparia, No. 554			3	2	20		21.6	.7917	m	t, cy	c	m	r	b	w
Monticola × Riparia, No. 18808			3	3	4		23.5	1.0087	m	t, cy	c	m	r	b	w
Monticola × Rupestris.			8	3	12 ¹ / ₂		23.3	.7550	m	t, cy	c	m	r	b	w
Motley.			5	3	5		21	.8225	m	t, cy	c	m	r	b	w
Mourvedre × Rupestris, No. 1202			8	6 ¹ / ₂	30		23.6	.8775	m	t, cy	c	m	r	b	w
Do.			20		19		22.5	.8175	m	t, cy	c	m	r	b	w
Mourvedre × Rupestris, No. 1203					1 ¹ / ₂		23	.9750	m	t, cy	c	m	r	b	w
Pardee.				2	22 ¹ / ₂		21.6	.7600	m	t, cy	c	m	r	b	w
Riparia du Colorado			1	2	15 ¹ / ₂		19.6	.7850	m	t, cy	c	m	r	b	w
Riparia × (Cordifolia × Rupestris), No. 106-8			7	4	2 ³ / ₄		19.6	.6508	m	t, cy	c	m	r	b	w
Riparia × Rupestris, No. 101.			20	7	28 ¹ / ₂		22.3	.8600	m	t, cy	c	m	r	b	w
Do.	50	5	11	5 ¹ / ₂	34		22.3	.8725	m	t, cy	c	m	r	b	w
Riparia × Rupestris, No. 101-14.			11	4 ¹ / ₂	28 ¹ / ₂		19.3	.7617	m	t, cy	c	m	r	b	w
Do.									m	t, cy	c	m	r	b	w
Riparia × Rupestris, No. 106-103.			8	2 ¹ / ₂	12 ¹ / ₂		21	.8512	m	t, cy	c	m	r	b	w
Riparia × Rupestris, No. 3306.			34	2 ¹ / ₂	37 ¹ / ₂		20.7	.8325	m	t, cy	c	m	r	b	w

	7	1	3	23.3	m	t, cy	c	m	r	b	w
Do.											
Riparia X Rupestris, No. 3309	26	8	32	23	.9450	t, cy	c	m	r	b	w
Do.	9	3	4	22.2	.6900	t, cy	c	m	r	b	w
Rupestris des Causseottes	7	1	21 $\frac{1}{2}$	21.3	.7853	t, cy	c	m	r	b	w
Rupestris Marth.	16	5	6	19	.9332	t, cy	c	m	r	b	w
Do.	9	1	6	22.4	.9885	t, cy	c	m	r	b	w
Rupestris Metallica.	5	3 $\frac{1}{2}$	12 $\frac{1}{2}$	21	.8020	t, cy	c	m	r	b	w
Do.	9	4 $\frac{1}{2}$	13 $\frac{1}{2}$	22	1.1025	t, cy	c	m	ob	b	w
Rupestris Mission.	8	4	13 $\frac{1}{2}$	20	.8825	t	l	m	ob	b	w
Do.	9	4	12 $\frac{1}{2}$	20.5	.9000	t	l	m	ob	b	w
Rupestris St. George.	16	4	28	22.3	.7650	t	l	m	ob	b	w
Do.	13	6	8	21.5	.8060	t	l	m	ob	b	w
Rupestris X Berlandieri, No. 219A	12	6	29	19.8	.9250	t	l	m	ob	b	w
Rupestris X Berlandieri, No. 301A	8	3 $\frac{1}{2}$	28	17.8	1.0300	t	l	m	ob	b	w
Rupestris X Berlandieri, No. 301B	13	3 $\frac{1}{2}$	32 $\frac{1}{2}$	20	.8125	t	l	m	ob	b	w
Rupestris X (Cordifolia X Rupestris), No. 202-5	10	3 $\frac{1}{2}$	17	20.3	.7696	t	l	m	ob	b	w
Rupestris X Hybrid Azemar, No. 215	2	1 $\frac{1}{2}$	11	22.6	.8275	t	l	m	ob	b	w
Salt Creek	7	3 $\frac{1}{2}$	13	21.6	.7425	t	l	m	ob	b	w
Do.	5	4	2	20.7	.8250	t	l	m	ob	b	w
Solonis Ordinaire.	2	2	3	20	.7900	t	l	m	ob	b	w
Solonis Robusta.	7	5	23 $\frac{1}{2}$	22	.7300	t	l	m	ob	b	w
Do.	8	4	31 $\frac{1}{2}$	27	.8475	t	l	m	ob	b	w
Solonis X Oihello.	15	4	19.9	19.9	.7200	t	l	m	ob	b	w
Solonis X Oihello, No. 1613	4	4	23	23	.8625	t	l	m	ob	b	w
Solonis X Riparia, No. 1616	24	3	28	21.6	.7450	t	l	m	ob	b	w
Do.	15	9	21.1	21.1	.8325	t	l	m	ob	b	w
Taylor Narbonne.	4	2	7	19.6	.8000	t	l	m	ob	b	w
Do.	18	5	8	23.4	.8301	t	l	m	ob	b	w
Petit Verdot:											
Own roots	3	4	2	24.7	.7700	cy	c	m	r	b	w
Dog Ridge	35	7	12	22.5	.8231	cy	c	m	r	b	w
Lenoir	40	10 $\frac{1}{2}$	3 $\frac{1}{2}$	24.2	.7115	cy	c	m	r	b	w
Rupestris St. George	45	5	18	23.7	.7325	cy	c	m	r	b	w
Taylor Narbonne.	30	3	1	23.3	.6844	cy	c	m	r	b	w
Peverell:											
Own roots	2	17	18	26	.7819	cy	c	m	r	b	w
Dog Ridge	40	8	3 $\frac{1}{2}$	22.3	.9175	cy	c	m	r	b	w
Lenoir	50	10	3	22.1	.8715	cy	c	m	r	b	w
Rupestris St. George	50	10	24	22.8	.8520	cy	c	m	r	b	w
Taylor Narbonne.	40	8	2	25	.5130	cy	c	m	r	b	w
Piepol:											
Own roots	8	11	22.5	22.5	.6862	s	l	s	ob	b	w
Piment:											
Lenoir	15	8	12	18.1	.6200	l	l	l	ob	b	w
Pince Muscat:											
Own roots	2	2	21.5	21.5	.6937	l	c	m	ob	b	w
Pineau de Chardonnay:											
Own roots	4	10	25	25	.6137	s	cy	s	r	b	w
Aramon X Rupestris Ganzin, No. 1	8	4	24.3	24.3	.6150	s	cy	s	r	b	w
Dog Ridge	2	2	25.2	25.2	.7281	s	cy	s	r	b	w
Herbement.	8	4	24.9	24.9	.7912	s	cy	s	r	b	w
Lenoir	12	1	2	23.3	.7581	s	cy	s	r	b	w

Origin/Variety	12	20	2	4	16	23	3787	1	t	m	1	0	w	s, t, r
Pizzutella:														
Own roots.....														
Adobe Giant.....	12	20	2	4	16	22.4	.4020	1	t	m	1	0	w	s, t, r
Aramon X Rupestris Ganzin, No. 1.....	8	12	4	4	12	24.5	.4660	1	t	m	1	0	w	s, t, r
Dog Ridge.....	3	3	2			23.2	.4331	1	t	m	1	0	w	s, t, r
Herbement.....						24.8	.3075	1	t	m	1	0	w	s, t, r
Lenoir.....	10	6	4	4	2	34.85	.3485	1	t	m	1	0	w	s, t, r
Mourvedre X Rupestris, No. 1202.....	6	10	5	3	3	23.6	.4005	1	t	m	1	0	w	s, t, r
Riparia Gloire.....	6	3	2	2	7	23.5	.4230	1	t	m	1	0	w	s, t, r
Riparia X Rupestris, No. 101.....	12	25	2	12	2	24.6	.4726	1	t	m	1	0	w	s, t, r
Riparia X Rupestris, No. 3309.....	10	10	3	8	1	33.00	.4300	1	t	m	1	0	w	s, t, r
Rupestris Martin.....	3	0	1			27.8	.3300	1	t	m	1	0	w	s, t, r
Rupestris St. George.....	15	10	4	5	8	24.9	.3336	1	t	m	1	0	w	s, t, r
Salt Creek.....	4	2	1	2	1	23.6	.4635	1	t	m	1	0	w	s, t, r
Solomis X Riparia, No. 1616.....	10	10	1	12	1	51.10	.5110	1	t	m	1	0	w	s, t, r
Taylor Narbonne.....	1	20	1	8	2	23.7	.2950	1	t	m	1	0	w	s, t, r
Ponsard:														
Own roots.....														
Precoce de Courtillet.....	5		13			27	.6466	1	cy	c	1	0	b	s, t, w
Lenoir.....	4			2		22.1	.8006	1	cy	l	m	0	w	s, t, w
Rupestris St. George.....	4					22.1	.8058	1	cy	l	m	0	w	s, t, w
Prune de Capoul:														
Aramon X Rupestris Ganzin, No. 2.....														
Australis.....			25			92	.8162	1	t	c	1	0	b	s, t, st
Berlandieri X Riparia, No. 420A.....			12			18	.8475	1	t	c	1	0	b	s, t, st
Monticola X Riparia, No. 18804.....			12			23.5	.6362	1	t	c	1	0	b	s, t, st
Monticola X Riparia, No. 18808.....			12		6	20.2	.7550	1	t	c	1	0	b	s, t, st
Monticola X Rupestris.....		5				21	.4275	1	t	c	1	0	b	s, t, st
Riparia X Cordifolia X Rupestris, No. 106-8.....			30			22	.7650	1	t	c	1	0	b	s, t, st
Riparia X Rupestris, No. 101-14.....			20			25	.6625	1	t	c	1	0	b	s, t, st
Riparia X Rupestris, No. 3306.....			24			22	.7800	1	t	c	1	0	b	s, t, st
Rupestris Martin.....			15			20	.7950	1	t	c	1	0	b	s, t, st
Purple Damascus:														
Own roots.....														
Adobe Giant.....	40	40	2	12	3	23.2	.6075	m-1	t, cy	m	1	0	b	s, t
Aramon X Rupestris Ganzin, No. 1.....	30	30	3	35	23	75.15	.5630	m-1	t, cy	m	1	0	b	s, t
Dog Ridge.....	13	45	4	10	6	71.80	.7180	m-1	t, cy	m	1	0	b	s, t
Lenoir.....	25	30	3	8	4	61.05	.7065	m-1	t, cy	m	1	0	b	s, t
Mourvedre X Rupestris, No. 1202.....	20	50	10	20	2	70.65	.7065	m-1	t, cy	m	1	0	b	s, t
Riparia Gloire.....	12	29	5	15	14	23.1	.6420	m-1	t, cy	m	1	0	b	s, t
Riparia X Rupestris, No. 101.....	15	25	2	7	6	50.50	.6050	m-1	t, cy	m	1	0	b	s, t
Riparia X Rupestris, No. 3309.....	30	35	3	36	22	23.1	.6795	m-1	t, cy	m	1	0	b	s, t
Rupestris St. George.....	12	30	5	18	12	22.1	.5970	m-1	t, cy	m	1	0	b	s, t
Solomis X Riparia, No. 1616.....	12	30	2	20	8	22.2	.5521	m-1	t, cy	m	1	0	b	s, t
Taylor Narbonne.....	20	50	30	15	2	22.1	.5170	m-1	t, cy	m	1	0	b	s, t
Quagliano:														
Own roots.....														
Dog Ridge.....	10	10		30	7	23.5	.6400	m-1	cy	c	m	0	b	s, t, w
Lenoir.....	60	17		33	6	22.7	.8085	m-1	cy	c	m	0	b	s, t, w
Rupestris St. George.....	40	5		17	1	22.7	.6285	m-1	cy	c	m	0	b	s, t, w
Taylor Narbonne.....	6	9		6	6	21.9	.6744	m-1	cy	c	m	0	b	s, t, w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Ballin's scale.	Acid, as tartaric C.C., (grams per 100)	Berry.			Cluster.			Use.		
	1909	1910	1911	1912	1913			22	23	24	25	26	27		28	29
I	17	18	19	20	21	22	23	24	25	26	27	28	29	30		
Razaki Zolo: Own roots.....	2	4	1	2		P. cent. 22.9	0.5100	m	cy	l	m	ob	r	s, t		
Rupestris St. George.....	12					20.8	.3250	m	cy	l	m	ob	r	s, t		
Riparia X Rupestris, No. 3306.....	12	13	13			20.9	.6252	m	cy	l	m	ob	r	s, t		
Refused:																
Own roots.....				2½		25	.8175	s	cy	c	m	r	b	w		
Dog Ridge.....	30	10	6	6	4	28.7	.6840	s	cy	c	m	r	b	w		
Lenoir.....	50	15	13½	8	28	23.2	.6965	s	cy	c	m	r	b	w		
Rupestris St. George.....	40	15	10	3	32	24.1	.6690	s	cy	c	m	r	b	w		
Taylor Narbonne.....	30	6	1½	1	3	24.5	.7005	s	cy	c	m	r	b	w		
Ribier:																
Lenoir.....	10					22	.6750	s	cy	c	m	r	b	w		
Rupestris St. George.....	10					21.5	.4800	s	cy	c	m	r	b	w		
Robin Noir.....	15	10	3	10		25.5	.7337	l	cy	m	m	ob	b	w		
Own roots.....																
Rodiles.....	25	3	12			21.1	.6641	v, l	t	l	l	ob	r	s, st, l		
Rosa d'Italie:																
Lenoir.....	5	7	½	3		23.2	.7250	m	t	l	m-1	ob	r	s, t, w		
Riparia X Rupestris, No. 3309.....	15	8	4			26	.6100	m	t	l	m-1	ob	r	s, t, w		
Rupestris St. George.....	10	1				26.2	.6750	m	t	l	m-1	ob	r	s, t, w		
Solanis X Riparia, No. 1616.....	15	12	15	2		26.8	.6556	m	t	l	m-1	ob	r	s, t, w		
Taylor Narbonne.....	5		1	½		23.6	.6850	m	t	l	m-1	ob	r	s, t, w		
Rothgraber:																
Own roots.....	20	3	20	8		25.1	.6687	m	t	c	m	o	w	w		
Rousson:																
Own roots.....	10	1		25	4	24.5	.7294	m	t	c	m	ob	w	w		
Dog Ridge.....	15	10	9	2		21.8	.6712	m	t	c	m	ob	w	w		
Lenoir.....	20	15	5	2½	36	22.1	.5900	m	t	c	m	ob	w	w		
Riparia X Rupestris, No. 3306.....	40	12	12	3		24.9	.5655	m	t	c	m	ob	w	w		
Riparia X Rupestris de Jaeger.....	2					23.6	.5437	m	t	c	m	ob	w	w		
Rupestris St. George.....	25	8	11		23	24.4	.5212	m	t	c	m	ob	w	w		

	20	6	3	27	.5175 .3750	.6150	.4625 .4218 .4130 .4170 .4512 .4462 .4555 .4839 .4710 .4287 .4125 .4545 .4594 .4360	23	11 1/2 8 10 16 1 6 12 4 5 12 8 15 10 5 8 12	11 2 9 6 3 1 1/2	24 24 26.2	1.0500 8775 9112	t t r	c c c	m m s	ob ob ro	w w b	w w w	s, t, st	
Solomus X Riparia, No. 1616.																				
Taylor Narbonne.																				
Royal Ascot.	20	6	3	27	.5175	.6150							t	c	m	ob	w	w	w	
Own roots																				
Saint Laurent:																				
Own roots																				
Adobe Giant.	1 1/2	3	1 1/2	26.3	.4625		3						cy	c	m-1	o	w	w	s, t	
Aramon X Rupestris Ganzin, No. 1.	8	14	8	25.4	.4218		3						cy	c	m-1	o	w	w	s, t	
Dog Ridge	8	10	10	25.4	.4130		3						cy	c	m-1	o	w	w	s, t	
Herbmont.	10	16	2	25.6	.4170		5						cy	c	m-1	o	w	w	s, t	
Lenoir	12	1	3	25	.4512		1						cy	c	m-1	o	w	w	s, t	
Mourvedre X Rupestris, No. 1202.	2	6	6	25.3	.4462								cy	c	m-1	o	w	w	s, t	
Riparia Gloire.	12	12	6	24.1	.4555		12	8					cy	c	m-1	o	w	w	s, t	
Riparia X Rupestris, No. 101.	5	4	4	26.2	.4839		2 1/2	11					cy	c	m-1	o	w	w	s, t	
Riparia X Rupestris, No. 3309.	8	12	2	26.5	.4710		3	2					cy	c	m-1	o	w	w	s, t	
Rupestris Martin.	15	3	1	24.1	.4287		1	2					cy	c	m-1	o	w	w	s, t	
Rupestris St. George.	10	10	4	26.2	.4125		4						cy	c	m-1	o	w	w	s, t	
Salt Creek.	5	10	12	25.8	.4545		2	8					cy	c	m-1	o	w	w	s, t	
Solomus X Riparia, No. 1616.	10	5	2	27.2	.4594		1						cy	c	m-1	o	w	w	s, t	
Taylor Narbonne.	8			28	.3975								cy	c	m-1	o	w	w	s, t	
Saint Macaire:	12	1 1/2	1 1/2	29.1	.4360		1 1/2						cy	c	m-1	o	w	w	s, t	
Own roots																				
Aramon X Rupestris Ganzin, No. 1.	12	4	4	22.2	.9056		11						r	c	m	ob	b	w	w	
Dog Ridge	20	5	11	26	.8175		2 1/2						r	c	m	ob	b	w	w	
Lenoir	40	15	13	23.6	.8630		9	2					r	c	m	ob	b	w	w	
Mourvedre X Rupestris, No. 1202.	40	10	11	22	1.0275		9	20					r	c	m	ob	b	w	w	
Rupestris St. George.	40	10	11	24	.8250		3	6					r	c	m	ob	b	w	w	
Solomus X Riparia, No. 1616.	50	10	9	24.8	.9400		3	20					r	c	m	ob	b	w	w	
Taylor Narbonne.	50	10	9	25	.7950		1 1/2	4					r	c	m	ob	b	w	w	
San Gioveco:																				
Dog Ridge	1 1/2			24	1.0500			3					t, cy	l	s	ob	b	w	w	
Lenoir	1 1/2			24	.8775								t, cy	l	s	ob	b	w	w	
Rupestris St. George.	1 1/2			26.2	.9112								t, cy	l	s	ob	b	w	w	
Taylor Narbonne.													t, cy	l	s	ob	b	w	w	
Satin Blanc:																				
Dog Ridge	6			24	.3975								cy	c	m	ob	b	w	w	
Lenoir	3			24.7	.4700			2					cy	c	m	ob	b	w	w	
Rupestris St. George.	8			23.3	.4950			6					cy	c	m	ob	b	w	w	
Sauvignon Blanc:																				
Own roots																				
Dog Ridge	12	10	12	25	.5475		1 1/2	3					cy	c	m	ob	b	w	w	
Lenoir	22	12	15 1/2	21	.8100		3	36					cy	c	m	ob	b	w	w	
Rupestris St. George.	3			22.3	.8635			7					cy	c	m	ob	b	w	w	
Sauvignon Vert:																				
Own roots																				
Aramon X Rupestris Ganzin, No. 1.	10	10	15	25.1	.4785		24	23					t, cy	l	m	ob	b	w	w	
Aramon X Rupestris Ganzin, No. 2.	60	40	19	25.4	.6195		10	15					t, cy	l	m	ob	b	w	w	
Bertandieri X Riparia, No. 420A				27	.5775		6	7					t, cy	l	m	ob	b	w	w	
Constantia				25.3	.6300		10	8					t, cy	l	m	ob	b	w	w	
Dog Ridge	55	12	16	21.5	.6225		12	21					t, cy	l	m	ob	b	w	w	
Herbmont.	60	15	12	22.9	.5970		12	4					t, cy	l	m	ob	b	w	w	

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).					Sugar, Ballings	Acid, as tartaric c. c., per 100	Cluster.			Berry.			Use.	
	1909							Size.	Shape.	Compact or loose.	Size.	Shape.	Color.		
	17	18	19	20	21										
I															
Sauvignon Vert—Continued.						P. cent.									
Lenoir.....	45	15	9	5	5	24.4	.5970	s	t, cy	m	r	w	w	w	w
Monticola X Riparia, No. 18808.....	3	1	3	2	18	24.3	.6530	s	t, cy	m	r	w	w	w	w
Mourvedre X Rupestris, No. 1202.....	50	12	13	2	18	22.8	.6825	s	t, cy	m	r	w	w	w	w
Riparia X Rupestris, No. 101.....	50	12	13	6	18	22.4	.6915	s	t, cy	m	r	w	w	w	w
Riparia X Rupestris, No. 101-14.....	2	2	2	2	24.5	24.5	.5137	s	t, cy	m	r	w	w	w	w
Riparia X Rupestris, No. 3306.....	60	10	10	8	7	24.5	.6937	s	t, cy	m	r	w	w	w	w
Riparia X Rupestris, No. 3309.....	60	10	10	8	10	25.6	.6013	s	t, cy	m	r	w	w	w	w
Rupestris Martin.....	60	12	8	23	23	23.1	.5130	s	t, cy	m	r	w	w	w	w
Rupestris St. George.....	55	8	5	2	4	25.1	.6325	s	t, cy	m	r	w	w	w	w
Salt Creek.....	50	12	9	3	7	24.1	.5905	s	t, cy	m	r	w	w	w	w
Solomis Robusta.....	2	2	2	2	2	27	.6262	s	t, cy	m	r	w	w	w	w
Solomis X Othello, No. 1013.....	55	45	13	15	4	25	.6466	s	t, cy	m	r	w	w	w	w
Solomis X Riparia, No. 1016.....	60	10	20	2	12	24.4	.5390	s	t, cy	m	r	w	w	w	w
Taylor Narbonne.....	2	2	9	24.4	.6185	s	t, cy	m	r	w	w	w	w
Schiradzouli Blanc.....	2	2	9	23	.4625	m	t	m	o	w	s, t, st		
Schiradzouli Violet.....	2	2	5	21	.6150	m	t	m	o	r	s, t, st		
Lenoir.....	8	3	2	15	4	24.6	.5670	m	cy	m	r	w	w	w	w
Own roots.....	30	10	12	3	4	24.2	.7556	m	cy	m	r	w	w	w	w
Aramon X Rupestris Ganzin, No. 1.....	25.5	.7680	m	cy	m	r	w	w	w	w
Berlandieri X Riparia, No. 420A.....	30	3	13	8	10	24.5	.7635	m	cy	m	r	w	w	w	w
Dog Ridge.....	40	15	15	5	12	22.3	.7200	m	cy	m	r	w	w	w	w
Herbmont.....	20	5	5	3	4	21.2	.7275	m	cy	m	r	w	w	w	w
Lenoir.....	20	5	5	3	4	27	.5400	m	cy	m	r	w	w	w	w
Monticola X Riparia, No. 18808.....	20	2	24	.5850	m	cy	m	r	w	w	w	w
Mourvedre X Rupestris, No. 1202.....	20	8	9	2	5	22.9	.6360	m	cy	m	r	w	w	w	w
Riparia X Rupestris, No. 101.....	35	5	3	3	4	25.6	.8075	m	cy	m	r	w	w	w	w
Riparia X Rupestris, No. 3306.....	35	5	2	2	3	21.8	.6921	m	cy	m	r	w	w	w	w
Riparia X Rupestris, No. 3309.....	20	2	2	25	.5819	m	cy	m	r	w	w	w	w
Rupestris Martin.....	20	2	7	25	.5819	m	cy	m	r	w	w	w	w
Rupestris St. George.....	20	8	22.6	.6015	m	cy	m	r	w	w	w	w

Salt Creek.....	25	5	6	1	4	25.2	.7020	m	cy	l	m	r	w	w
Solonis X Othello, No. 1613.....					3	25	.9675	m	cy	l	m	r	w	w
Solomis X Riparia, No. 1616.....	25	5			2	27.2	.5275	m	cy	l	m	r	w	w
Taylor Narbonne.....	25	4	4	4	2	25.4	.6900	m	cy	l	m	r	w	w
Semillon Blanc:														
Own roots.....	3	4		4	8	23.8	.6206	m	cy	l	m	r	w	w
Setine:														
Own roots.....	2		1	3	2	25.5	.7312	s	cy	c	s	r	b	b
Dog Ridge.....	15	5	9	9	3	26.9	.8460	s	cy	c	s	r	b	b
Lenoir.....	25	8	7	6	14	25	.6662	s	cy	c	s	r	b	b
Rupestris St. George.....	35	15	19	2	10	25.9	.8145	s	cy	c	s	r	b	b
Taylor Narbonne.....		10		3		26	.7387	s	cy	c	s	r	b	b
Servan Rose:														
Lenoir.....	8		6	6	6	20	.7150	m	r	l	l	ob	r	s, t, st
Sicilien:														
Own roots.....		1		3		20.5	.7162	l	l	l	l	ob	w	s, t, st
Aramon X Rupestris Ganzin, No. 1.....	16					17.4	.6937	l	l	l	l	ob	w	s, t, st
Lenoir.....	4		5½		½	24.3	.4875	l	l	l	l	ob	w	s, t, st
Slankamenka:														
Own roots.....	2	1	2	10		25	.7181	l	l	l	l	ob	w	s, t, st
Souvenir du Congrès:														
Dog Ridge.....	11		1			23.6	.5550	s	cy	l	s	r	w	w, t
Lenoir.....	2					23	.5325	s	cy	l	s	r	w	w, t
Rupestris St. George.....	3		½			22	.6825	s	cy	l	s	r	w	w, t
Spartanosa:														
Own roots.....	6	3	1	15		21.1	.7237	m	r	l	m	r	b	w
Sucre de Marseille:														
Dog Ridge.....	20	2	2	10	3	24.7	.5830	s	r	l	m	r	w	w
Lenoir.....	20	5	9		3	24	.6137	s	r	l	m	r	w	w
Rupestris St. George.....	10	3	1½		8	24.8	.6075	s	r	l	m	r	w	w
Taylor Narbonne.....	5					26.8	.4875	s	r	l	m	r	w	w
Suleicho:														
Own roots.....	25					15.9	1.1475	s	cy	c	m	ob	w	st, s, t
Sultana:														
Own roots.....			½		4	23.5	.8887	l	l	l	m	r	w	s, t, r, w
Sultamina:														
Own roots.....		½			4	25.5	.6187	vl	l	l	m	ob	w	s, t, r, w
Do.....	20	10				24	.5425	vl	l	l	m	ob	w	s, t, r, w
Aramon X Rupestris Ganzin, No. 1.....	12	12			25.5	25.5	.6112	vl	l	l	m	ob	w	s, t, r, w
Do.....			4	46		25.5	.5850	vl	l	l	m	ob	w	s, t, r, w
Aramon X Rupestris Ganzin, No. 2.....				4		24	4.500	vl	l	l	m	ob	w	s, t, r, w
Australis.....				9	13	25	5.100	vl	l	l	m	ob	w	s, t, r, w
Dog Ridge.....	25					23	4.950	vl	l	l	m	ob	w	s, t, r, w
Hotporup.....				6		22	5.025	vl	l	l	m	ob	w	s, t, r, w
Lenoir.....	8		4			23.5	.6070	vl	l	l	m	ob	w	s, t, r, w
Monticola X Riparia, No. 554.....		3	2	9		24.1	.5500	vl	l	l	m	ob	w	s, t, r, w
Monticola X Riparia, No. 18804.....						23	.5775	vl	l	l	m	ob	w	s, t, r, w
Monticola X Riparia, No. 18808.....			8			23	.5325	vl	l	l	m	ob	w	s, t, r, w
Monticola X Riparia, No. 18815.....		5				25.1	.5100	vl	l	l	m	ob	w	s, t, r, w
Monticola X Rupestris.....						23	.6850	vl	l	l	m	ob	w	s, t, r, w
Ponroy.....				4		23	.6150	vl	l	l	m	ob	w	s, t, r, w

TABLE VII.—Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).						Sugar, Ballings scale.	Acid, as tartaric (grams per 100 c.c.).	Cluster.			Berry.		Use.			
	1909	1910	1911	1912	1913	21			22	23	24	25	26		27	28	29
Sultana—Continued.																	
Ramsey.....							24.7										
Riparia X (Cordifolia Rupestris, No. 106-8).....		2	8	15			25	0.5470	v1	1	1	m	ob	w	s, t, w		
Riparia X Rupestris, No. 101.....							25	.5475	v1	1	1	m	ob	w	s, t, w		
Riparia X Rupestris, No. 101-14.....	15	2		5			26.6	.5850	v1	1	1	m	ob	w	s, t, w		
Riparia X Rupestris, No. 3306.....				4			23	.5962	v1	1	1	m	ob	w	s, t, w		
Riparia X Rupestris, No. 3309.....	20	3	5				23	.4950	v1	1	1	m	ob	w	s, t, w		
Riparia X Rupestris de Jaeger, No. 201.....				16			22	.6025	v1	1	1	m	ob	w	s, t, w		
Rupestris des Caussettes.....				7			20	.5625	v1	1	1	m	ob	w	s, t, w		
Rupestris Martin.....				3			25	1.2175	v1	1	1	m	ob	w	s, t, w		
Rupestris Mission.....				10			26	.5025	v1	1	1	m	ob	w	s, t, w		
Rupestris X Berlandieri, No. 219 A.....			6	14			24.8	.5645	v1	1	1	m	ob	w	s, t, w		
Salt Creek.....			10				25	.5400	v1	1	1	m	ob	w	s, t, w		
Solonis Robusta.....				5			29	.5775	v1	1	1	m	ob	w	s, t, w		
Solonis X Othello, No. 1613.....				11			24	.5625	v1	1	1	m	ob	w	s, t, w		
Solonis X Riparia, No. 1616.....	25	50					27.8	.5325	v1	1	1	m	ob	w	s, t, w		
Vermorel.....			3				23	.6000	v1	1	1	m	ob	w	s, t, w		
Viala.....			5	20			25	.5041	v1	1	1	m	ob	w	s, t, w		
Sultana Rosaee.....							25.2	.6675	1	1	1 to c	m	ob	r	s, st, t, 1		
Lenoir.....	1	2															
Sylvaner.....	3	2	4	18			25.5	.5868	s	cy	c	s	r	w	w		
Own roots.....	20	10	11	3			24.4	.6450	s	cy	c	s	r	w	w		
Aramon X Rupestris Ganzin, No. 1.....	20	16	9	4		6	22.6	.7365	s	cy	c	s	r	w	w		
Dog Ridge.....	20	12	6	2			24.6	.7312	s	cy	c	s	r	w	w		
Herbement.....	20	10	10	1			24.7	.6974	s	cy	c	s	r	w	w		
Riparia.....	25	3	3	2		7	25.2	.7475	s	cy	c	s	r	w	w		
Riparia X Rupestris, No. 101.....	25	9	4	1			26.5	.7069	s	cy	c	s	r	w	w		
Riparia X Rupestris, No. 3309.....	15	10	2	8		6	26.5	.7100	s	cy	c	s	r	w	w		
Rupestris Martin.....	15	2	2	1		2	27.7	.7500	s	cy	c	s	r	w	w		
Rupestris St. George.....	24	2	7	1													
Own roots.....		13	10	5			23	.6650	1	1	1	1	ob	w	s, st, t		

Valdepenas:	2	5	2	10	23.8	.7294	m	cy	m	m	ob	b	w
Own roots.....	30	6	10	10	24.5	.6445	m	cy	1	1	ob	b	w
Aramon X Rupestris Ganzin, No. 1.....	30	9	10	18	24.5	.6705	m	cy	1	1	ob	b	w
Dog Ridge.....	45	5	11	18	23.2	.6705	m	cy	1	1	ob	b	w
Herbement.....	40	5	4	18	23.8	.7530	m	cy	1	1	ob	b	w
Lenoir.....	30	3	3	3	26.1	.6763	m	cy	1	1	ob	b	w
Mourvedre X Rupestris, No. 1202.....	30	8	2	15	24.6	.6437	m	cy	1	1	ob	b	w
Riparia X Rupestris, No. 101.....	30	5	7	29	25.7	.5975	m	cy	1	1	ob	b	w
Riparia X Rupestris, No. 3309.....	45	7	3	12	24.9	.6074	m	cy	1	1	ob	b	w
Rupestris Martin.....	30	7	3	5	24.1	.7065	m	cy	1	1	ob	b	w
Rupestris St. George.....	30	7	11	4	25.7	.6131	m	cy	1	1	ob	b	w
Salt Creek.....	25	15	2	3	24.7	.6420	m	cy	1	1	ob	b	w
Solonia X Riparia, No. 1616.....	30	2	13	3	26.7	.6345	m	t	1	1	ob	r	w
Taylor Narbonne.....	30	4	4	7	27.2	.6415	m	t	1	1	ob	r	w
Veltheimer:	3	5	6	18	27.2	.6415	m	t	1	1	ob	r	w
Own roots.....	50	4	2	3	25.6	.7305	m	t	1	1	ob	r	w
Dog Ridge.....	35	15	3	3	23.9	.6356	m	t	1	1	ob	r	w
Herbement.....	40	3	1	2	25	.7425	m	t	1	1	ob	r	w
Lenoir.....	40	3	3	3	23.6	.6925	m	t	1	1	ob	r	w
Riparia X Rupestris, No. 3306.....	30	1	1	1	25	.7069	m	t	1	1	ob	r	w
Rupestris Martin.....	40	3	1	1	24	.6850	m	t	1	1	ob	r	w
Rupestris St. George.....	40	3	1	1	21.3	.6606	m	t	1	1	ob	w	s, st, t
Salt Creek.....	40	8	6	3	19.4	.5998	m	t	1	1	ob	w	s, st, t
Verdel:	20	5	20	12	21.3	.6606	m	t	1	1	ob	w	s, st, t
Own roots.....	40	3	18	4	21.3	.5739	m	t	1	1	ob	w	s, st, t
Dog Ridge.....	60	70	60	10	20	.8310	m	t	1	1	ob	w	s, st, t
Lenoir.....	60	17	22	10	20.1	.8300	m	t	1	1	ob	w	s, st, t
Rupestris St. George.....	25	5	14	7	21.9	.6860	m	t	1	1	ob	w	s, st, t
Verdel:	50	50	8	40	20	.8300	m	t	1	1	ob	w	s, st, t
Own roots.....	1	8	10	8	21.9	.7815	s	cy	c	s	r	d	w
Adobe Giant.....	30	30	4	12	24.3	.5820	s	cy	c	s	r	d	w
Aramon X Rupestris Ganzin, No. 1.....	15	3	15	7	24.2	.6035	s	cy	c	s	r	d	w
Dog Ridge.....	20	30	6	10	22.9	.6975	s	cy	c	s	r	d	w
Mourvedre X Rupestris, No. 1202.....	30	35	12	25	24.8	.6537	s	cy	c	s	r	d	w
Riparia X Rupestris, No. 101.....	16	10	5	14	25.7	.4640	s	cy	c	s	r	d	w
Riparia X Rupestris, No. 3309.....	40	50	5	38	22	.6180	s	cy	c	s	r	d	w
Rupestris St. George.....	30	40	5	12	24.8	.6723	s	cy	c	s	r	d	w
Solonia X Riparia, No. 1616.....	15	5	2	8	25.7	.4530	s	cy	c	s	r	d	w
Taylor Narbonne.....	15	20	6	15	24.5	.4935	s	cy	c	s	r	d	w
Verdelho:	8	1	3	3	25.6	.8006	m	t	1	1	ob	w	w
Own roots.....	3	3	6	2	26.6	.6781	m	t	1	1	ob	w	w
Verdelho de Madere:	2	5	5	9	28.9	.7275	m	t	1	1	ob	w	w
Aramon X Rupestris Ganzin, No. 1.....	8	9	5	5	26.7	.7165	m	t	1	1	ob	w	w
Dog Ridge.....	10	8	5	12	27.6	.7245	m	t	1	1	ob	w	w
Riparia X Rupestris, No. 3309.....	8	5	1	4	27.6	.7144	m	t	1	1	ob	w	w
Rupestris St. George.....	10	8	5	12	27.6	.7144	m	t	1	1	ob	w	w
Solonia X Riparia, No. 1616.....	8	4	1	3	27.6	.7144	m	t	1	1	ob	w	w

TABLE VII. Relative behavior and value for different purposes of grape varieties tested by grafting on resistant stocks and by growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and stock (on own roots, if so stated).	Weight of fruit per vine (pounds).						Sugar, Balling scale.	Acid, as tartaric (C. C.).	Cluster.			Berry.			Use.			
	1909	1910	1911	1912	1913	1914			22	23	24	25	26	27		28	29	
																		17
Vermentino:																		
Own roots.....	10	20	20	8			22.1	0.5720	m-1	t	m	l	ob	W	W			
Adobe Giant.....	50	35	5	25	18		22.4	.5750	m-1	t	m	l	ob	W	W			
Aramon X Rupestris Ganzin, No. 1.....	25	30	8	5	16		22.7	.4755	m-1	t	m	l	ob	W	W			
Dog Ridge.....	40	10	10	8	8		22.7	.4227	m-1	t	m	l	ob	W	W			
Hesperiont.....	25	15	3	3	6		22.3	.5517	m-1	t	m	l	ob	W	W			
Leonard.....	40	40	5	15	12		22.3	.4935	m-1	t	m	l	ob	W	W			
Mourvedre X Rupestris, No. 1202.....	20	30	5	10	7		22.9	.4865	m-1	t	m	l	ob	W	W			
Riparia Gloire.....	15	50	3	38	17		22.9	.4890	m-1	t	m	l	ob	W	W			
Riparia X Rupestris, No. 101.....	50	50	3	25	30		22.7	.5670	m-1	t	m	l	ob	W	W			
Riparia X Rupestris, No. 3300.....	55	4	10	25	11		21.3	.5370	m-1	t	m	l	ob	W	W			
Rupestris Marzemino.....	5	5	1 ¹	1 ¹	0		21.3	.5370	m-1	t	m	l	ob	W	W			
Rupestris St. George.....	50	30	15 ¹	20	12		22.4	.5057	m-1	t	m	l	ob	W	W			
Salt Creek.....	25	30	1	1	5		22.0	.4694	m-1	t	m	l	ob	W	W			
Solonis X Riparia, No. 1616.....	30	30	6	35	12		21.9	.4890	m-1	t	m	l	ob	W	W			
Taylor Northome.....	12	50	12	35	24		23.2	.4455	m-1	t	m	l	ob	W	W			
Vernaccia Sarda:																		
Own roots.....	12	6	3	32	16		25.6	.8265	m	t	l	m	ob	W	S, st, t, w			
Vigne de Zericho:																		
Lenoir.....	4	11	13	13			19.2	.7363	l	t	c	l	ob	W	S, st, t			
Mourvedre X Rupestris, No. 1202.....	4	11	13	13			22	.6188	l	t	c	l	ob	W	S, st, t			
Wälschriesling:																		
Own roots.....	6	3	3	8	6		25.8	.6165	s	cy	c	s	f	W	W			
Wernre:																		
Own roots.....	4	2	2	3	1		26.1	.6120	m	t	t	m	f	W	W			
White Hanepoot:																		
Own roots.....	10	10	6	6	7		27.4	.5812	m	t	l	l	ob	W	S, st, t, f			
Adobe Giant.....	10	10	6	6	7		27.4	.5812	m	t	l	l	ob	W	S, st, t, f			
Aramon X Rupestris Ganzin, No. 2.....	10	10	6	6	7		24	.6900	m	t	l	l	ob	W	S, st, t, f			
Australis.....	15	15	15	15	15		29	.4425	m	t	l	l	ob	W	S, st, t, f			
Berlandieri X Riparia, No. 420A.....	15	15	15	15	15		31	.5025	m	t	l	l	ob	W	S, st, t, f			
Dog Ridge.....	15	15	15	15	15		25.5	.5565	m	t	l	l	ob	W	S, st, t, f			

Table VIII gives an alphabetical list of additional Vinifera varieties grafted on resistant stocks which are under test at the Chico, Colfax, Fresno, Geyserville, Lodi, and Oakville experiment vineyards. Arabic figures are used to indicate the number of different stocks upon which each is grafted and the location of the same. The tests are too young to permit the drawing of conclusions.

TABLE VIII.—Additional Vinifera varieties of grapes under test on resistant stocks at six experiment vineyards in California.

[Arabic figures are used to indicate the number of different stocks upon which each variety is grafted and the location of the same.]

Varieties.	Chico.	Colfax.	Fresno.	Geyserville.	Lodi.	Oakville.	Varieties.	Chico.	Colfax.	Fresno.	Geyserville.	Lodi.	Oakville.
Ach-I-Soum						1	Child of Hall	2	5			5	4
Actoni Maceron	1						Cinsaut	1					
Actoniky	1	2	3		4	4	Clairette à Gros Grain	1	6			6	9
Agadia						1	Coarna Neagra	2	5				6
Agra-Ash						3	Corinth Blanc				9		
Ahmeur bon Ahmeur	2	4			4	5	Corinth à Gros Grain	1	5	6	2	1	6
Ah Sûibe						1	Corinth Rose	2	6	12		4	7
Ajaki-Odia	1						Coristano	1	8			6	3
Ajmi	1					5	Cornichon	2	4	3		6	4
Ak-Uzum						2	Damas Rose			17			
Albardiens	1						Damugue	1					
Aldara						1	Dattier de Beyrouth					1	3
Alexandria			4				Deis-al-A'anze	1					6
Alicante	2						Dizmar	1					
Almeria	1				7	1	Dnrelabi	1					
Amlachu						1	Dronkusa	2					5
Aneb el Cadi	1	4			4	5	Emathia	1	6	2		10	6
Angelina	2	5			5	7	Erz Emperor	1	3	6		3	1
Aramon	1						Erz Roumilli	1					9
Arecussia							Fajoumi Jaune	1	5			6	3
Ash Khuta						2	Fajphy		8	1		9	16
Askaree	1					4	Fehher Som	1	7			7	5
Asmi						6	Fehher Zagos	1					
Atch Gau						2	Ferrara	1	3	25			6
Aswad Kari						2	Fintendo	2					
Atch Kiek						2	Flame Tokay	1	9	4		4	3
Aturk-Ash						2	Frankenthal Precoce	2		2			
Angulata	1	7	1		5	3	Fredericton	1	5	1		6	5
Awasarghua						2	Ghulabi						2
Bokator						1	Golden Champion						6
Barducci	2	4	2		5	4	Golden Hamburg						5
Bermestia Bianca	1						Golden Queen	1				3	11
Bicane	1	5	1		6	5	Goolabie	1	5	24		2	2
Black Hamburg	2	2			2	2	Green Hungarian	1					3
Black Monukka	1		18		1	8	Grenache	1					
Black Morocco	1	7			2		Gros Blanc de Lausanne	1	5	1		7	5
Black Prince	1						Gros Guillaume			24			5
Black Seedless						3	Hebron	1	8			6	10
Black Shahanee	1	4			5	5	Huasco			10			
Blanc d'Ambre	1	4			5	5	Hunisa	1	6	14			1
Blaney White	1	3			6	9	Hyeales	1	12	1			8
Blauer Portugieser							Insolia Bianca	2	6	2		6	16
Bowood Muscat	1						Jauzani						5
Brustiana	2	5			7	8	Jbâ'i						2
Buecleugh	1	5			5	8	Jerisiana						1
Buckland	1					8	Joannenc						2
Calabrian	1	7			7	6	Jubelli	1					6
Calmette	1		3			2	Kabbajuk						2
Carignane	1					6	Kadarka	1					
Castiza		4			4	5	Kandihar						2
Chadeh Arabieh	2				6	6	Kara-sarma						1
Chaouch Blanche	1	6			6	5	Kara-uzum						1
Chaouch Rose	1	9			8	8	Kharashani						1
Chasselas Rose de Faloux			1				Kâstûfi-dakar						2
Chasselas Rouge	1	2				3	Kâstûfi-inti						5
Chasselas St. Bernard	1	2	5		5	4	Katchich						1
Chavoosche	1						Kechmish-Aly-Blanc	2	5	2		7	3
Chauche Gris						3	Keropodia	1	7	2		7	8
Chaweesh	1					5	Key	1	2			2	6

TABLE VIII.—Additional *Vinifera* varieties of grapes under test on resistant stocks at six experiment vineyards in California—Continued.

Varieties.	Chico.	Collax.	Fresno.	Geyserville.	Lodi.	Oakville.	Varieties.	Chico.	Collax.	Fresno.	Geyserville.	Lodi.	Oakville.
Khudud-ul-Banat.....					5	3	Prune de Cazouls.....	2	5	23		8	5
Kishmish.....						2	Purple Damascus.....	2	6				2
Kishmish Daba.....						1	Quagliano.....	1					
Kishmish Red.....						1	Red Hanepoot.....			17			1
Kishmishi.....	1					1	Ribier.....	1					
Kordash.....						2	Rodites.....	1	11	25		7	14
Korsa Kishmish.....						1	Ronde Weisse.....						1
Ksil-Isjum.....						1	Rose d'Italie.....	1	6			3	4
Kurdi.....	1					1	Roussaou.....			2			
Kuristi Mici.....	1	5	2		7	4	Saibe.....						2
Kurtelaska.....	1						St. Macaire.....	1					
Leani Zolo.....	2					5	Satin Blanc.....	1	8			6	8
Lignan.....	1	2	2			3	Schiradzouli Violet.....	1	7	22		7	8
Listan.....	1		6			1	Semillon.....						9
Madeleine Royale.....						1	Servan Rose.....	1	6	2		7	7
Malvasia.....	1	6			4	5	Sev-ursa.....						1
Malvasia Rosaria.....						1	Sgtoruk.....						
Mantua de Pilas.....	1						Shahmani.....						5
Maraville de Malaga.....	1	6	23		6	5	Shakafi.....						2
Marmora.....	1						Shakaribira.....						1
Melon.....	1	7				1	Shanzi.....						1
Meyer No. 64.....						1	Shatawi.....						5
Meyer No. 114.....						2	Shirshira.....						1
Meyer No. 115.....						1	Sicilien.....	1	9	1		6	11
Meyer No. 801.....						3	Slankamenka.....	2					
Meyer No. 803.....						1	S. P. I. No. 601.....						1
Meyer No. 832.....						2	S. P. I. No. 611.....						1
Meyer No. 866.....						1	S. P. I. No. 614.....						1
Miksasi.....						5	S. P. I. No. 6914.....						1
Millenium.....	1	3	5			8	S. P. I. No. 30042.....						1
Molinera Gordo.....	2	8	2		5	10	Sultana.....			1			2
Monake.....						1	Sultanina.....			14			4
Mondeuse.....	1						Sultanina Rosea.....	3	38			6	22
Mourestel.....	1					4	Suri.....						6
Mukhkh-ul-Baghl.....						2	Sylvaner.....						5
Muscat Albardiens.....		6					Syrian.....					3	
Muscat Bonod.....	1	6	2		9	10	Tavris.....						1
Muscat Capusines.....	1					4	Tavris.....						1
Muscat Gros Noir Hatif.....	2		1				Teneron.....	1	9	3			12
Muscat Hamburg.....			5				Terret Monstre.....	1	8	2		8	6
Muscat Noir Precoce.....	1		2		1		Tiffihi Ahmer.....	1					3
Nasa Valentiana.....	1						Tinta Amerella.....	1					
Nebbiolo.....						1	Trifere du Japon.....	1	13	2		5	15
Ohanez.....	1	13	48	79	10	6	Triomphe.....	2	8				5
Olivette Blanche.....	1	15	5		9	17	Trojka.....	1					
Olivette Chaptal.....	1	9	1		6	10	Ubeide.....						6
Olivette Noir.....	2	10			6	10	Uva de Casta.....						6
Olivette Rose.....			1				Uva de Embarque.....						6
Opiman.....	2	3			4	3	Valandova.....			25			
Pagadebito.....	1						Valdenpasas.....	1					
Palarusa.....	2						Veltliner.....						11
Panariti.....	1		25				Velusa.....						1
Panse de Roquevaire.....						5	Verdal.....		2			1	1
Paykanee Razuki.....	1						Verdelho de Madere.....			1			
Pearl de Casaba.....	1				3	3	Vigne de Zericho.....	1	6	22		7	6
Pedro Ximines.....	2						West Prolific.....						4
Perle Imperial Blanche.....	2	7			3	11	White Hanepoot.....			10			
Peru.....	1	1			1	2	White Kapadulari.....						2
Petit Syrah.....	1			1		3	White Luglienga.....	1	12	2		10	14
Philipi.....						1	White Tokay.....	1	1				2
Piment.....	1	5	29		10	6	Zabalkanski.....	1					3
Pis des Chevre Rouge.....	1	6	3		6	5	Zeine.....						6
Pizzutella.....	2						Zinfandel.....	1					4

The following is an alphabetically arranged list of *Vinifera* varieties in the Brawley Experiment Vineyard (Pl. II, fig. 2), the plantings of which are as yet too young to permit the drawing of conclusions relative to their adaptability:

Ach-I-Soum, Actoni, Actoniky, Agadia, Ahmeur bon Ahmeur, Ajmi, Aleatico, Alexandria, Almeria, Aneb-el-Cadi, Angelina, Augulata, Askari, Aspiran Noir, Atch Kiek, Awasarghua.

Barducci, Beclan, Bermestia Bianca, Boal de Madera, Bowood Muscat, Black Morocco, Black St. Peter, Black Shahanee, Black Zante, Blanc d'Ambre, Blaney White, Blauer Portugieser, Buccleuch, Brustiana.

Carignane, Calabrian, Catarratto a la Porta, Castiza, Chadieh Arabieh, Chavooschee, Chaouch, Chaouch Rose, Chasselas Bouches du Rhone, Chasselas Dore, Chasselas Fontainebleau, Chasselas Florence, Chasselas Montauban, Chasselas Musque Vrai, Chasselas Negrepoint, Chasselas Rose, Chasselas Rose de Falloux, Chasselas Rouge, Chasselas St. Bernard, Child of Hall, Cinsaut, Clairette à Gros Grain, Clairette Blanche, Coarna Neagra, Commandeur, Corinthe Rose, Corinthe à Gros Grain, Cornichon, Coristana, Crujidero.

Damascus, Danugue, Dattier de Beyrouth, Dizmar, Dodrelabi, Downing, Drnekusa, Dronkane, Duc de Magenta.

Emperor.

Fajaumi Jaune, Faphly, Feher Goher Noir, Feher Som, Feher Zagos, Ferrara, Fintendo, Flame Tokay, Foster, Frankenthal Precoce, Frederickton.

Golden Champion, Golden Hamburg, Golden Queen, Goolabie, Gradiska, Green Hungarian, Grenache, Gros Blanc de Lausanne, Gros Maroc.

Hebron Hycals, Hunisa, Huasco, Hutab.

Insolia Bianca.

Jura Muscat.

Kadarka, Kahallillee, Kakour, Keropodia, Kishmish, Kuristi Mici, Kurtelaska.

Lahntraube, Lampasas, Leani Zolo, Luglienga.

Madeleine Angevine, Madeleine Blanche, Madeleine Rose, Madeleine Royale, Mamelon, Malaga, Malvasia, Malvasia Rosario, Maraville de Malaga, Marascina, Marzamina, Meslier, Millenium, Mission, Molinera Gordo, Mondeuse, Mourisco Bianca, Muscateller, Muscatelle Fino, Muscat Albardiens, Muscat Bonod, Muscat Capusines, Muscat Gros Noir Hatif, Muscat Hamburg, Muscat Madera Rose, Muscat Noir de Hongrie, Muscat Noir Precoce, Muscat Talabot.

Napoleon, Nasa Valentiana, Negrara di Gattinara, Neiretta di Costillo.

Oeru di Boe, Ohanez, Olivette Blanche, Olivette de Cadenet, Oliver de Serres, Olivette Noir.

Pagadebito, Parc de Versailles, Palarusa, Panariti, Paykane Razuki, Pearl de Casaba, Perruno, Persian, No. 20; Persian, No. 21; Persian, No. 23; Persian, No. 25; Persian, No. 26; Persian, No Number; Persian, No Tag; Philipi, Piment, Pince Muscat, Prune de Cazouls.

Rose de Italie, Royal Ascot, Rozaki Zolo.

Schach-I-Soum, Schiradzouli Blanc, Schiradzouli Violet, Servan Blanc, Servan Rose, Sgotoruk, Shiraz, Sicilian, Slankamenka, St. Laurient, Sultana, Sultanina, Sultanina Rosea, Syrian.

Terret Monstre, Tinta Amerilla, Torok Goher Noir, Trentham Black, Trifere du Japon, Triomphe, Trivoti, Trojka, Trousseau, Tsien Tsien.

Valdepenas, Veltliner, Verdal, Vernaccia Sarda, Vigne de Zericho.

Wilmot Hamburg, Wilmot No. 16, White Corinth, White Tokay.

Zante.

Table IX shows additional tests of improved American native and Franco-American grape varieties grafted on resistant stocks which are being made at the Fresno and Oakville experiment vineyards. (See Pl. I, figs. 1 and 2.) Arabic figures are used to indicate the number of different stocks upon which each is grafted and the location of the same. The tests are too young to permit the drawing of conclusions.

TABLE IX.—Varieties of American native and Franco-American grapes under test on resistant stocks at the Fresno and Oakville experiment vineyards, showing the number of different stocks included in the test.

Varieties.	Fresno.	Oakville.	Varieties.	Fresno.	Oakville.
Agawam.....stocks..	1	4	Concord.....stocks..	5	10
America × Malaga No. 2:			Concord Improved.....do.....		1
No. 1.....stocks..		1	Continental.....do.....		1
No. 2.....do.....		2	Lindley.....do.....	4	11
No. 16.....do.....		1	Lucile.....do.....		6
Armistead Seedling.....do.....		1	Massasoit.....do.....		1
Barry.....do.....	1	4	Moore.....do.....	1	7
Brighton.....do.....	4	6	Niagara.....do.....	5	6
Brilliant.....do.....		4	Perkins.....do.....		1
Campbell.....do.....		5	Rebecca.....do.....	1	4
Catawba.....do.....	1	9	Riggs No. 16.....do.....		1
Champion.....do.....		2			

The explanation of columns already given for Table VII will apply for the most part to Table X, showing the relative behavior and value for different purposes of improved American native and Franco-American grape varieties, except that the parentage of the varieties displaces the designation of the stock, the column for the "Year grafted" is omitted, and the congeniality column is changed to a growth-rating column, because all the varieties are on their own roots. The column for weight of pruning is also omitted. The abbreviations used to designate the parent species are as follows: Aest. for Aestivalis, Bourq. for Bourquiniana, Champ. for Champini, Lab. for Labrusca, Lins. for Linsecomii, Rip. for Riparia, Rup. for Rupestris, Vin. for Vinifera.

Do.	1904	93	S	3 to 5	Mar. 30	Apr. 8	June 9	June 18	May 26	June 23	Sept. 26
Do.	1903	87	S	3 to 5	Mar. 20	Apr. 10	May 21	June 8	May 26	June 23	Sept. 24
Do.	1904	89	S	3 to 5	Mar. 16	Apr. 10	May 26	June 25	May 30	June 29	Sept. 24
Blondin:	1903	53	S	1 to 4	Mar. 14	Apr. 1	May 18	June 5	May 23	June 20	Sept. 17
(Bourq. X Aest.) X (Lhs. X Lab.)	1906	86	S	1 to 4	Mar. 20	Apr. 8	May 19	June 5	May 23	June 20	Sept. 12
Bourisquon X Rupestris, No. 601:	1903	58	S	1 to 4	Mar. 14	Apr. 1	May 18	June 5	May 23	June 20	Sept. 5
Vin. X Rup.	1906	86	S	1 to 4	Mar. 20	Apr. 8	May 19	June 5	May 23	June 20	Sept. 5
Do.	1905	90	S	1 to 4	Mar. 18	Apr. 10	May 3	June 7	May 9	June 11	Sept. 5
Do.	1905	87	S	1 to 4	Mar. 18	Apr. 10	May 3	June 7	May 9	June 11	Sept. 5
Do.	1903	93	C, S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1907	90	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1906	90	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	84	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	79	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	92	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	87	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Bourisquon X Rupestris, No. 603:	1906	90	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Vin. X Rup.	1904	84	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	91	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	94	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	97	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	84	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	88	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	98	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Bourisquon X Rupestris, No. 109-4:	1907	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Vin. X Rup.	1907	89	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	94	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	97	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	84	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	88	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1905	98	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Bourisquon X Rupestris, No. 3907:	1906	91	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Vin. X Rup.	1907	91	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	94	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	93	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	97	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	87	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Bourisquon X Rupestris, No. 4306:	1907	94	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Vin. X Rup.	1904	99	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1907	94	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	94	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	98	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Bourisquon X Rupestris, No. 4308:	1906	92	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Vin. X Rup.	1907	89	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	96	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	85	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	85	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Do.	1904	95	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Brighton:	1905	87	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25
Lab. X Vin.	1905	87	S	1 to 4	Mar. 18	Apr. 1	May 8	June 15	May 12	May 28	Sept. 25

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Experiment vineyard.	Year planted.	Growth rating.	How pruned.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
						Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
I		3	4	5	6	7	8	9	10	11	12	13	14
Brilliant: Lab. X (Vin. X Bourq.)	Gi	1905	Per cl. 77	S	Mar. 6	Apr. 1	May 26	May 30	June 3	Sept. 5	Sept. 25
Do.	F	1904	80	S	Mar. 12do.	May 15	May 20	May 15	May 24	Aug. 20	Sept. 4
Do.	G	1905	73	S	Mar. 2	Mar. 20	May 22	May 18	May 26	June 23	Sept. 2	Sept. 23
Do.	M	1904	84	S	Mar. 26	Mar. 31	May 30	June 7	June 10	June 11	Oct. 4	Sept. 26
Canada: (Rup. X Lab.) X Vin.	F	1903	87	S	Mar. 10	Mar. 25	May 10	May 25	May 15	May 30	Sept. 5	Sept. 19
Do.	L	1906	88	S	Mar. 17	Apr. 15	May 18	May 30	May 12	June 4	Sept. 20	Sept. 22
Do.	O	1903	96	S	1 to 3	Mar. 20	Apr. 4	May 15	May 25	May 25	June 10	Sept. 10	Sept. 23
Do.	S	1906	93	S	Mar. 10	Apr. 12	May 25	June 15	June 19	Sept. 24	Sept. 26
Carignane X Rupestris, No. 404: Vin. X Rup.	F	1904	97	S	Mar. 7	Mar. 30	May 10	May 20	May 15	May 26	Sept. 9	Sept. 17
Do.	C	1904	95	S	Mar. 16	Mar. 20	May 19	June 7	May 24	June 12	Sept. 26	Sept. 23
Do.	Li	1905	84	S	Mar. 13	Apr. 1	May 23	June 10	June 8	June 14	Sept. 23	Oct. 1
Do.	M	1904	87	S	Mar. 23do.	May 28	June 8	June 2	June 13	Sept. 26	Oct. 8
Do.	S	1904	86	S	1 to 3	Mar. 18do.	May 22	May 27	May 26	June 1do.	Oct. 7
Do.	S	1907	89	S	Mar. 12do.	May 29	June 5	June 4	June 20	Sept. 24
Carignane X Rupestris, No. 501: Vin. X Rup.	L	1904	92	S	Mar. 7	Mar. 30	May 10	June 3	May 14	June 5	Oct. 26
Do.	M	1904	90	S	Mar. 26	Mar. 29	June 13	June 12	June 5	June 10	Oct. 10	Oct. 1
Do.	O	1904	95	S	1 to 3	Mar. 16	Apr. 1	May 14	May 18	June 1	June 10	Sept. 27	Oct. 17
Do.	S	1907	90	S	Mar. 14	Mar. 30	May 27	June 10do.	Oct. 17
Carmen: Lins. X (Vin. X Lab. X Bourq.)	F	1904	52	Sdo.	Mar. 20	May 21	May 26	May 25	June 3	Aug. 23	Sept. 19
Do.	O	1903	74	S	1 to 3	Mar. 16	Apr. 3	May 22	May 31	May 27	June 23	Sept. 10	Sept. 25
Castel, No. 1028: Rup. X Vin.	O	1907	84	S	2 to 4	Mar. 20	Apr. 6	May 28	May 29	June 2	June 13	Sept. 9	Oct. 6
Castel, No. 10002: (Lab. X Rup.) X Vin.	O	1907	87	S	2 to 4	Mar. 19	Apr. 1	May 7	May 25	May 21	May 27	Sept. 22	Oct. 9

Catwaba:	1906	68	s	Mar. 20	Apr. 8	May 26	June 3	May 30	June 8	Sept. 24
Lab. X Vin.....	1905	68	s	Mar. 7	Apr. 8	May 26	June 3	May 30	June 8	Sept. 24
Do.....	1905	68	s	Mar. 7	Apr. 8	May 26	June 3	May 30	June 8	Sept. 24
Do.....	1905	91	s	Mar. 10	Apr. 31	May 24	June 5	May 28	May 29	Sept. 11
Do.....	1905	88	s	Mar. 11	Apr. 3	May 12	June 5	May 28	May 29	Sept. 23
Do.....	1905	87	s	Mar. 26	Apr. 10	May 30	June 3	June 3	June 16	Oct. 10
Do.....	1905	77	s	Mar. 22	Apr. 1	May 17	May 25	May 22	June 11	Sept. 9
Do.....	1905	90	c, s	1 to 4	Mar. 12	Apr. 2	May 20	June 10	May 30	June 14	Oct. 14
Centennial:											
Lab. X Aest.....	1905	87	s	Mar. 15	Apr. 9	May 9	May 17	May 24	May 25	Sept. 25
Do.....	1904	86	s	Mar. 13	Apr. 25	May 10	May 28	May 24	May 25	Sept. 5
Do.....	1905	79	s	2 to 3	Mar. 14	Apr. 1	May 16	May 28	May 21	June 1	Sept. 9
Champagne:											
Chateau X Lab.....	1903	94	s	2 to 3	Mar. 22	Apr. 3	May 19	May 24	May 24	June 11	Sept. 27
Chaselas X Rupestris, No. 901:											
Vin. X Rup.....	1904	91	s	Mar. 6	Mar. 15	May 6	May 16	May 15	June 1	Sept. 8
Do.....	1904	95	s	1 to 2	Mar. 17	Mar. 25	May 12	May 25	May 24	June 1	Sept. 27
Clainette Doré Ganzin:											
Vin. X Rup.....	1903	88	c, s	Mar. 9	Apr. 5	May 12	May 20	May 20	May 26	Sept. 25
Do.....	1905	90	s	do.....	Mar. 27	May 26	June 3	May 30	June 9	Sept. 27
Do.....	1907	73	s	Mar. 6	Mar. 21	May 13	May 22	May 18	May 26	Sept. 27
Do.....	1903	94	s	2 to 4	Mar. 16	Mar. 22	May 20	May 30	May 24	June 4	Oct. 15
Clevener:											
Lab. X (Rip. X Aest.).....	1907	83	s	Mar. 22	Mar. 30	May 8	May 22	May 12	May 29	Oct. 9
Cloëta:											
(Lins. X Rup.) X (Lab. X Vin.).....	1903	90	s	2 to 4	Mar. 21	Apr. 9	May 19	May 29	May 24	June 3	Sept. 24
Columbaud X Rupestris:											
Vin. X Rup.....	1903	96	s	2 to 4	Mar. 17	Mar. 25	May 22	May 28	June 1	June 5	Sept. 26
Concord:											
Lab.....	1906	88	s	Mar. 23	Apr. 6	June 1	June 7	June 5	June 12	Oct. 4
Do.....	1905	68	s	Mar. 18	Apr. 1	May 26	Sept. 25
Do.....	1904	76	s	2 to 4	Mar. 17	Mar. 30	May 12	May 28	May 17	June 2	Oct. 7
Do.....	1904	83	s	Mar. 12	Apr. 10	May 10	June 1	June 5	June 7	Sept. 20
Cornucopia:											
Vin. X Rip.....	1903	56	s	Mar. 1	Apr. 1	May 1	May 10	May 15	Sept. 6
Do.....	1907	88	s	Mar. 12	Mar. 27	May 22	June 8	May 27	Oct. 15
Couderc, No. 101:											
Vin. X Rip.....	1907	94	s	Mar. 15	Apr. 4	May 28	June 4	June 8	June 12	Sept. 15
Do.....	1904	93	c, s	Mar. 8	Mar. 15	May 6	May 24	May 25	May 29	Sept. 17
Do.....	1904	93	s	Mar. 4	Mar. 14	do.....	May 16	May 12	May 21	Sept. 10
Do.....	1904	94	s	Mar. 9	Mar. 30	May 12	June 8	May 16	May 13	Sept. 24
Do.....	1904	86	s	Mar. 26	Apr. 4	May 29	June 6	June 3	June 10	Sept. 26
Do.....	1904	93	s	1 to 5	Mar. 18	Mar. 25	May 17	May 25	May 23	May 29	Sept. 25
Do.....	1904	91	s	Mar. 10	Apr. 8	May 20	June 15	May 24	June 19	Sept. 22
Couderc, No. 201:											
Rip. X (Rup. X Vin.).....	1907	90	s	Mar. 17	Mar. 28	May 10	May 25	May 25	Sept. 20
Do.....	1904	98	c, s	Mar. 10	Mar. 25	do.....	May 20	May 17	May 25	Sept. 4
Do.....	1904	95	s	Mar. 2	Mar. 22	May 17	June 17	May 21	June 16	Sept. 10
Do.....	1905	96	s	Mar. 12	Apr. 6	May 23	June 8	May 29	June 12	Sept. 23
Do.....	1904	87	s	Mar. 26	do.....	do.....	June 10	May 31	June 14	Oct. 1
Do.....	1904	94	s	2 to 5	Mar. 20	Apr. 7	May 18	May 24	May 18	May 31	Sept. 26
Do.....	1904	90	s	Mar. 10	Apr. 5	May 27	June 8	May 31	June 13	Sept. 27
Do.....	1904	90	s	Mar. 10	Apr. 5	May 27	June 8	May 31	June 13	Sept. 27

Do.	1904	2 to 4	Mar. 21	Apr. 1	May 17	May 26	May 23	May 29	Sept. 25	Oct. 5
Do.	1904	Mar. 10do.	May 31	June 15	June 5	June 20	Sept. 24	Oct. 14
Couderc, No. 71-06: Rup. X (Lins. X Vin.)	1907	2 to 3	Mar. 23	Mar. 27	May 17	June 8	May 21	June 14	Sept. 25	Oct. 10
Couderc, No. 71-20: Rup. X (Lins. X Vin.)	1907	2 to 5do.	Mar. 25do.	June 6do.	June 11	Sept. 21	Oct. 8
Couderc, No. 74-17: Complex Hybrid.	1907	1 to 3	Mar. 24	Apr. 1	May 23	May 28	June 1	June 5	Oct. 6	Oct. 10
Couderc, No. 82 X 32: Complex Hybrid.	1907	Mar. 25	Apr. 10	May 2	June 12	May 9	June 17	Sept. 25	Do.
Do.	1904	Mar. 15	Mar. 22	May 15	May 25	May 21	June 1	Sept. 1	Sept. 19
Do.	1904	Mar. 2	Mar. 25	May 23	June 1	May 27	June 1	Sept. 20
Do.	1904	Mar. 15	Apr. 5	May 24	June 10	May 22	June 15	Sept. 15
Do.	1904	Mar. 26do.	June 9	June 18	May 29	June 22	Sept. 28	Oct. 4
Do.	1904	Mar. 16do.	May 28	June 10	May 26	June 16	Sept. 22	Oct. 10
Do.	1904	Mar. 12	Apr. 7	May 28	June 30	June 2	July 3	Sept. 30	Oct. 14
Couderc, No. 84 X 61: (Vin. X Rup.) X Vin.	1907	Mar. 17	Apr. 2	May 25	June 4	June 8	June 11	Sept. 20	Oct. 4
Do.	1904	Mar. 10	Apr. 8	May 12	May 26	May 30	May 31	Sept. 15	Sept. 23
Do.	1904	Mar. 11	Mar. 20	May 12	May 26	May 20	May 31	Sept. 8	Sept. 20
Do.	1904	Mar. 5	Mar. 16	May 16	June 8do.	June 12	Sept. 20	Oct. 8
Do.	1904	Mar. 10	Mar. 25	May 26	June 10	May 30	June 14	Sept. 20	Oct. 1
Do.	1907	Mar. 3	Mar. 26	May 12	June 29	May 17	June 4do.
Do.	1904	Mar. 29	Apr. 1	Apr. 26	June 9	Apr. 30	June 19	Sept. 24	Oct. 4
Do.	1904	2 to 4	Mar. 20	Mar. 30	May 18	June 10	May 23	June 16	Sept. 26	Oct. 9
Do.	1904	Mar. 13	May 26	June 1	June 18	May 23	June 16	Sept. 27	Oct. 2
Couderc, No. 85 X 113: (Vin. X Rup.) X Vin.	1907	2 to 4	Mar. 24	Apr. 1	May 31	June 1	June 3	June 20	Sept. 23	Oct. 10
Couderc, No. 87 X 115: Vin. X (Rup. X Vin.)	1907	Mar. 16	Apr. 5	June 1	June 18	June 5	June 23	Sept. 15	Sept. 23
Do.	1904	Mar. 8	Apr. 1	May 6	May 11	May 10	June 5	Sept. 5	Sept. 19
Do.	1904	Mar. 15	Mar. 20	May 15	May 21	May 25	May 29	Aug. 21	Sept. 19
Do.	1907	Mar. 2	Mar. 25	May 16	June 3	May 20	June 12	Sept. 23	Oct. 10
Do.	1907	Mar. 9	Mar. 30	May 26	June 18	May 29	June 22do.
Do.	1904	Mar. 27	Apr. 3	May 25	June 9	Apr. 29	June 22do.
Do.	1904	1 to 2	Mar. 17	Apr. 1	May 17	June 7	May 22	June 12	Sept. 24	Oct. 7
Do.	1904	Mar. 18do.	May 17	June 7	May 22	June 12	Sept. 26	Oct. 17
Couderc, No. 124 X 30: Vin. X (Rup. X Vin.)	1907	Mar. 7	Mar. 28	May 10	May 25	Sept. 25
Do.	1904	Mar. 10	Mar. 25	May 14	May 24	May 20	May 29	Sept. 9	Sept. 17
Do.	1904	Mar. 2	Mar. 30	May 21	June 15	May 25	June 30	Sept. 25	Oct. 5
Do.	1904	Mar. 13	Apr. 3	May 12	June 14	May 18do.	Sept. 22
Do.	1904	Mar. 14do.	May 16	June 3	Sept. 24	Do.
Do.	1904	Mar. 14	Apr. 1	May 29	Apr. 29	June 20	Sept. 25	Oct. 8
Do.	1904	1 to 3	Mar. 28	Apr. 2	May 17	June 5	Apr. 24	June 10	Sept. 27	Oct. 10
Do.	1904	Mar. 18	Mar. 27	May 17	June 5	May 24	June 17	Sept. 20	Sept. 25
Do.	1904	Mar. 12	Apr. 1	May 22	June 12	May 27	June 17	Sept. 20	Oct. 30
Couderc, No. 132-11: Complex Hybrid.	1907	1 to 3	May 24	Mar. 25	May 26	June 3	May 30	June 9	Sept. 27	Oct. 30
Couderc, No. 199-88: Vin. X (Rup. X Vin.)	1907	1 to 2	Mar. 25do.do.	May 31	May 24	June 4	Sept. 25	Oct. 5
Couderc, No. 241-125: Vin. X (Rup. X Vin.)	1907	2 to 4	Mar. 27	Mar. 28do.	June 6	May 30	June 13	Sept. 24	Oct. 8

Gold Coin: Aest. X Lab.	1904	F	60	s	Mar. 23	Apr. 8	May 12	May 30	May 20	June 6	Sept. 5	Sept. 19	
	Do.	G	79	s	Mar. 7	Apr. 5	May 18	June 14	June 18	Sept. 20	Oct. 7	
	Do.	O	67	s	3 to 4	Mar. 16	Apr. 3	May 30	June 1	June 3	June 15	Sept. 25	Sept. 25	
	Do.	S	85	s	Mar. 10	Apr. 12	May 23	June 18	June 1	June 23	Sept. 24	Oct. 16	
Governor Ross: Lab. X Vin.	1904	GI	88	s	Mar. 20	Apr. 8	May 20	June 3	May 25	June 10	Sept. 18		
	Herbement:	1906	Cx	92	c, s	Mar. 18	Apr. 3	Apr. 21	June 18	Apr. 28	June 24	Sept. 25	Oct. 4
		1905	GI	84	s	Mar. 10	Apr. 5	May 26	May 31	May 31	May 31	Sept. 19	Sept. 30
		1904	M	87	s	Mar. 9	Apr. 1	Apr. 30	May 13	May 4	May 27	Sept. 26	Oct. 8
1903		O	84	s	2 to 3	Mar. 22	do.	do.	do.	do.	June 9	do.	Oct. 10	
Herbert:	1904	S	78	s	Mar. 12	Mar. 25	May 25do.	June 8	June 11		
	1905	F	70	s	Mar. 5	Apr. 10	May 1	May 28	May 20	June 1	Aug. 25	Sept. 10	
	Do.	O	87	s	Mar. 14	Apr. 1	May 18	Oct. 9		
	1905	F	85	s	Mar. 9	Apr. 5	May 15	May 20	May 19	May 26	Sept. 9	Sept. 17	
Hexamer:	1905	F	85	s	Mar. 15	Apr. 6	May 17	June 12	May 22	June 18	Sept. 8	Oct. 15	
	1905	O	89	s	2 to 4	
Husmann:	1906	O	73	s	2 to 3	
	1905	GI	91	c, s	
Isabella: Lab. X Lab.	1906	Cx	91	s	Mar. 19	Mar. 31	Apr. 30	June 5	May 6	June 10	Sept. 15	Oct. 2	
	1905	GI	73	s	Mar. 10	Apr. 15	May 5	May 23	May 14	May 26	Sept. 12	Sept. 20	
Jaeger: Lins. X Bourq.	1905	F	90	s	Mar. 6	Mar. 18	May 5	June 14	May 15	May 19	Sept. 12		
	Do.	L	88	s	Mar. 9	Apr. 1	May 8	May 25	May 11	June 9	Sept. 25		
	Do.	M	81	s	Mar. 25	Apr. 3	May 28	June 3	June 11	June 10		
	Do.	O	91	s	1 to 5	Mar. 16	Apr. 1	May 15	May 26	May 21	May 30		
	Do.	S	97	c, s	Mar. 10	Apr. 30	May 20	June 10	May 21	May 30		
	Do.	F	68	s	Mar. 15	Apr. 3	May 14	May 22	May 20	May 27	Sept. 24	Oct. 17	
	Do.	G	87	s	Mar. 4	Apr. 5	May 29	June 3	May 28	June 12	Sept. 5	Sept. 19	
	Do.	L	70	s	Mar. 20	Apr. 6	May 12	May 20	May 10	May 24	Sept. 20	Sept. 25	
	Do.	O	79	s	1 to 3	Mar. 25	Apr. 2	May 28	June 5	June 3	June 17	Sept. 25		
Jefferson: Lab. X Vin.	1906	C	80	s	Mar. 31	Apr. 16	May 30	June 3do.	June 8	Oct. 4	
	1905	O	69	s	1 to 3	Mar. 24	Apr. 1	May 16	May 28	May 22	May 31	Sept. 23		
Kiowa: Lins. X Bourq.	1903	F	55	s	Mar. 10	Mar. 12	Apr. 18	May 3	May 5	May 12		
	1903	O	86	s	2 to 3	Mar. 24	Apr. 5	May 28	June 7	June 3	June 12	Sept. 27		
Lampasas: Lab. X Aest.	1908	F	91	c, s	Mar. 10	Mar. 18	May 8	May 15	May 15	Sept. 10	
	1904	F	27	s	Mar. 12	Mar. 25	May 5	May 17	May 23		
Lenoir: Bourq.	1906	Cx	93	c, s	Mar. 23	Apr. 2	June 9	June 12	June 14	June 17	Sept. 10	Sept. 23	
	1905	GI	98	s	Mar. 18	Mar. 28	May 25	May 30	May 30	June 5	Sept. 29	Sept. 29	
	1904	F	79	s	Mar. 14	Mar. 20	May 20	May 30	May 20	May 20	Sept. 8	Sept. 12	

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Experiment vineyard.	Year planted.	Growth rating.	How pruned.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
						Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
1	2	3	4	5	6	7	8	9	10	11	12	13	14
Lenoir—Continued.													
Bourq.	G	1904	<i>Per ct.</i> 95	S	Mar. 2	Mar. 27	May 24	June 5	May 29	June 29	Sept. 22	Oct. 7
Do.	L	1904	94	S	Mar. 8	Mar. 28	May 23	do.....	May 28	June 29	Sept. 25	Oct. 28
Do.	L	1904	96	S	Mar. 12	Mar. 31	May 18	May 20	May 25	May 31	Sept. 24	Oct. 3
Do.	M	1904	82	S	Mar. 27	Apr. 5	May 26	June 15	May 30	June 20	Sept. 26	Oct. 8
Do.	S	1903	98	S	3 to 5	Mar. 16	Apr. 3	May 19	June 1	May 23	June 5	Sept. 29	Oct. 3
Do.	S	1904	93	S	Mar. 4	Apr. 2	May 27	June 15	June 1	June 29	Sept. 24	Oct. 16
Lindley:													
Lab. X Vin.	CX	1906	89	S	Mar. 22	Apr. 1	Apr. 30	June 5	May 7	June 20	Sept. 10	Oct. 4
Do.	GI	1905	83	S	Mar. 10	Mar. 28	May 15	May 24	May 20	May 28	Sept. 20	Oct. 21
Do.	F	1904	88	S	Mar. 12	Apr. 1	May 9	May 15	May 12	May 21	Aug. 24	Sept. 21
Do.	O	1905	93	S	2 to 4	Mar. 16	Mar. 26	May 10	May 27	May 14	June 1	Sept. 9	Oct. 7
Louisiana:													
Bourq.	O	1903	57	S	2 to 4	Mar. 20	Apr. 15	May 22	June 10	May 27	June 23	do.....	Do.
Do.	S	1907	86	S	Mar. 14	Apr. 5	May 30	June 12	June 4	do.....	do.
Lukfafa:													
Champ. X Lab.	F	1903	83	S	Mar. 10	Apr. 1	Apr. 3	May 10	May 6	May 14	Sept. 23	Sept. 26
Do.	L	1904	88	S	Mar. 5	Apr. 5	May 12	May 25	May 26	May 30
Do.	O	1903	95	S	Mar. 19	Apr. 4	do.....	May 23	May 16	May 28
Do.	O	1903	93	S	2 to 3	Mar. 16	do.....	May 18	June 1	May 23	June 6	Sept. 10
Manito:													
(Lab. X Vin. X Bourq.) X (Lins. X Rup.)	O	1903	86	S	Mar. 6	Apr. 3	June 5	Sept. 19
Marguerite:													
Lins. X Bourq.	G	1904	76	S	Mar. 18	do.....	May 25	June 5	May 29
Do.	L	1904	70	S	Mar. 19	do.....	June 4	June 10	June 7	June 23
Do.	O	1903	91	S	Mar. 19	Apr. 9	May 29	June 15	June 2	June 30
Do.	S	1904	92	S	Mar. 12	Apr. 7	Oct. 14
Martha:													
Lab. X Vin.	GI	1905	60	S	Mar. 20	Apr. 15	May 25	June 3	Sept. 25

Missouri Riesling: Rip. X Lab.	1904 L	83	s	Mar. 13	Apr. 1	May 8	May 17	May 18	May 22	Sept. 5	Sept. 19
Do.	1904 O	82	s	Mar. 14	Apr. 3	May 12	May 28	May 12	June 1	Sept. 24	Sept. 25
Do.	1904	70	s	1 to 3	Mar. 14	Apr. 3	May 12	May 27	May 17	May 31	Sept. 20	Oct. 5
Mrs. Munson: Lins. X Bourq.	1905	80	s	3 to 5	Mar. 12	Apr. 6	May 19	June 15	May 23	June 22	Sept. 22	Oct. 1
Muench: Lins. X Bourq.	1904 F	87	s	3 to 4	Mar. 22	Apr. 1	May 18	May 30	May 23	June 6	Sept. 6	Sept. 19
Do.	1903	82	s	Mar. 20	Apr. 1	May 18	June 1	May 25do.	Sept. 25	Sept. 27
Niagara: Lab. X Vin.	1906 Cx	88	s	3 to 7	Mar. 19	Apr. 5	May 3do.	May 12	June 7	Sept. 5	Oct. 4
Do.	1905 F	71	s	1 to 3	Mar. 7	Apr. 7	May 6	May 14do.	May 19	Aug. 24	Oct. 26
Do.	1905 O	89	s	1 to 3	Mar. 13	Mar. 25	May 16	May 27	May 20	June 21	Sept. 10	Sept. 21
Oliatafo: (Vin. X Lab.) X (Lins. X Lab.)	1907	81	s	2 to 4	Mar. 20	Mar. 30	June 8	June 10	June 12	June 16	Sept. 8	Oct. 2
Pardes: Vin. X Rup.	1907 Cx	94	s	Mar. 19	Apr. 3	Apr. 27	June 5	May 2	June 11	Sept. 25	Oct. 4
Do.	1905 Gl	89	s	Mar. 8	Mar. 25	May 20	May 21	May 15	May 25	Sept. 19	Sept. 26
Do.	1904 F	89	s	Mar. 5	Mar. 28	May 5	May 15	May 15	May 21	Sept. 8	Sept. 20
Do.	1904 Li	73	s	Mar. 7	Mar. 25	May 10	May 28	May 16	June 16	Sept. 15	Sept. 28
Do.	1904 M	53	s	Mar. 20	Mar. 29	June 6	June 12	June 11	Sept. 25	Oct. 10
Do.	1904 O	93	s	2 to 4	Mar. 18	Mar. 25	May 15	May 28	May 20	June 3	Sept. 26	Oct. 9
Do.	1904 S	95	s	Mar. 14	Mar. 27	May 26	June 10	May 31	June 14	Sept. 24	Oct. 14
Pierce: Lab. X Vin.	1905 L	88	s	Mar. 11	Apr. 3	May 4	May 18	May 7	May 23do.	Oct. 2
Do.	1905 M	85	s	Mar. 27	Apr. 1	May 21	June 8	May 26	June 13	Oct. 6
Do.	1905 S	92	s	Mar. 14	Apr. 5	May 23	June 3	May 27	June 9	Sept. 30
Rebecca: Lab. X Vin.	1905 F	63	s	Mar. 9	Apr. 1	May 6	May 18	May 18	May 23	Aug. 25	Aug. 30
Do.	1905 M	81	s	Mar. 28	Apr. 6	June 3	June 8	June 11	June 16	Sept. 4
Do.	1905 O	89	s	1 to 3	Mar. 19	Apr. 3	May 12	May 24	May 16	May 29	Sept. 9	Oct. 10
Do.	1905 S	84	s	Mar. 13	Apr. 5	May 28	June 15	June 1	June 18	Sept. 24	Oct. 16
Rommel: Lab. X (Rip. X Vin.)	1904 F	84	s	Mar. 9	Mar. 25	May 7	May 18	May 15	May 23	Aug. 20	Sept. 20
Do.	1904 L	86	s	Mar. 18	Apr. 3	May 10	May 27	May 14	June 1	Sept. 24	Oct. 5
Do.	1907 S	89	s	Mar. 12	Apr. 1	May 24	June 28	May 28	July 1	Oct. 17
R. W. Munson: Lins. X (Lab. X Vin.)	1906 C	90	s	Mar. 27	Apr. 8	May 8	May 20	May 9	June 4
Do.	1906 Cx	80	s	Mar. 6	Apr. 5	May 1	June 10	May 9	June 15
Do.	1907 Gl	83	s	Mar. 6	Apr. 1	May 18	May 26	May 18	May 31	Aug. 24	Sept. 15
Do.	1904 F	83	s	Mar. 6	Apr. 3	May 22do.	May 18	May 29
Do.	1904 C	71	s	Mar. 4	Apr. 3	May 2	June 13	June 8	June 20
Do.	1904 Li	71	s	Mar. 10	Apr. 3	June 2	June 15	June 8	June 20
Do.	1904 L	78	s	Mar. 11	Apr. 3	June 2	June 18	May 17	May 23
Do.	1904 M	92	s	Mar. 29do.	June 8	June 14	May 17	June 19
Do.	1903 O	91	s	2 to 5	Mar. 22do.	May 18	May 30	May 24	June 4	Sept. 26
Do.	1904 S	85	s	Mar. 13	Apr. 12	May 26	June 20	May 31	June 25

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Experiment vineyard.	Year planted.	Growth rating.	How pruned.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-selling date.		Fruit-ripening date.		
						Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	
Seibel, No. 1: Rup. X (Lms. X Vin.)	C	1907	<i>Per et.</i>			7	8	9	10	11	12	13	14	
	CX	1907	93	s		Mar. 29	Apr. 13	May 2	May 20	May 27	June 8	Sept. 26	Oct. 15	
	GI	1905	91	s		Mar. 19	Apr. 1	May 30	June 3	June 5	June 2	Sept. 16	Sept. 28	
	Do.	1904	94	s		Mar. 15	Apr. 1	May 20	May 25	May 15	May 31	Aug. 20	Sept. 15	
	F	1904	74	s		Mar. 11	do.	do.	May 5	May 18	June 18	Sept. 24	Sept. 30	
	G	1904	90	s		Mar. 4	do.	May 21	June 15	May 25	June 21	Sept. 21	Sept. 30	
	L	1904	48	s		Mar. 14	Apr. 5	May 15	June 4	May 17	June 20	Sept. 20	Oct. 3	
	M	1904	89	s		Mar. 13	Apr. 1	May 13	June 15	May 17	June 20	Sept. 26	Oct. 8	
	O	1904	63	s		Mar. 27	Apr. 4	May 15	June 5	May 20	June 10	Sept. 28	Oct. 9	
	S	1904	86	s		Mar. 21	Mar. 30	May 15	June 18	May 25	June 23	Sept. 24	Oct. 12	
	Do.	1904	86	s		Mar. 20	Apr. 1	May 15	June 1	May 20	June 5	Sept. 24	Oct. 12	
	Seibel, No. 2: Rup. X (Lms. X Vin.)	C	1910	93	s		Mar. 28	Apr. 6	May 16	May 27	May 21	June 1	Sept. 24	Oct. 5
		CX	1907	90	s		Mar. 24	do.	June 5	June 9	June 10	June 14	Sept. 18	Sept. 25
		GI	1905	92	s		Mar. 20	Mar. 20	May 13	May 27	May 24	June 4	Aug. 23	Do.
F		1904	78	s		Mar. 10	Mar. 20	May 10	May 25	May 30	May 30	Aug. 23	Do.	
Do.		1904	82	s		Mar. 6	Apr. 1	May 28	June 10	May 20	June 14	Sept. 25	Oct. 5	
G		1904	41	s		Mar. 14	do.	May 16	June 20	May 20	June 25	Sept. 25	Oct. 5	
L		1904	82	s		Mar. 12	Mar. 30	May 19	June 3	May 22	June 10	Sept. 25	Oct. 8	
M		1904	88	s		Mar. 28	Apr. 6	May 26	do.	June 1	June 9	Sept. 28	Oct. 9	
O		1904	82	s		Mar. 21	Mar. 29	do.	June 31	May 30	June 9	Sept. 25	Oct. 12	
Do.		1904	82	s		Mar. 17	Apr. 1	do.	June 18	May 30	June 27	Sept. 25	Oct. 12	
S		1907	87	s		Mar. 15	Mar. 25	May 15	June 2	May 20	June 7	Sept. 25	Oct. 12	
Seibel, No. 14: Rup. X (Lms. X Vin.)		C	1906	92	s		Mar. 31	Apr. 10	May 20	June 1	May 30	June 6	Sept. 5	Sept. 23
		CX	1906	95	s		Mar. 20	Apr. 5	May 4	June 12	May 17	June 17	Sept. 19	Sept. 25
		GI	1905	63	s		Mar. 9	Mar. 25	May 9	May 27	May 13	May 30	Sept. 19	Sept. 25
	Do.	1905	63	s		Mar. 9	Mar. 25	May 9	May 27	May 13	May 30	Sept. 19	Sept. 25	
	O	1904	84	s		Mar. 17	Apr. 1	May 19	May 29	May 24	June 3	Sept. 20	Oct. 9	
	Do.	1904	84	s		Mar. 17	Apr. 1	May 19	May 29	May 24	June 3	Sept. 20	Oct. 9	

Seibel, No. 29: Rup. × (Lins. × Vin.)	0	1907	88	s	2 to 4	Mar. 25	Apr. 12	June 1	June 15	June 5	June 20	Sept. 25	Oct. 15
Seibel, No. 38: Rup. × (Lins. × Vin.)	C	1906	93	s	Mar. 27	Apr. 21	May 20	June 1	June 7
Seibel, No. 60: Rup. × (Lins. × Vin.)	0	1907	70	s	Apr. 5	May 29	June 8	June 2	June 12
Seibel, No. 70: Rup. × (Lins. × Vin. × Rup.)	0	1907	81	s	2 to 5	Mar. 19	Apr. 1	May 26	June 5	June 1	June 10	Sept. 25	Oct. 10
Seibel, No. 78: Rup. × (Lins. × Vin.)	0	1907	80	s	2 to 5	Mar. 26	Mar. 28	June 8	June 14	Sept. 23	Oct. 2
Seibel, No. 80: Rup. × (Lins. × Vin.)	0	1907	89	s	2 to 5	Mar. 24	May 21	May 30	May 25	June 5	Sept. 24	Oct. 7
Seibel, No. 128: Rup. × (Lins. × Vin.)	0	1907	83	s	2 to 4	Apr. 1	May 26	June 11	June 1	June 16	Sept. 21	Oct. 8
Seibel, No. 156: Rup. × (Lins. × Vin.)	0	1907	84	s	2 to 3	Mar. 21	Apr. 3	May 27	June 7	June 13	Sept. 23	Oct. 15
Seibel, No. 209: Rup. × (Lins. × Vin.)	0	1907	82	s	2 to 4	Mar. 25	Mar. 28	May 31	June 12	June 4	June 18	Sept. 24	Oct. 8
Seibel, No. 215: Rup. × (Lins. × Vin.)	C	1906	93	s	Mar. 30	Apr. 10	May 16	June 3	May 21	June 9	Sept. 20
Do.	G1	1905	76	s	Mar. 20	Mar. 30	May 25	June 9	June 14	Sept. 22	Oct. 27
Do.	0	1907	86	s	2 to 4	Mar. 21	Apr. 3	May 26	June 9	June 1
Seibel, No. 334: Rup. × (Lins. × Vin. × Rup.)	0	1907	78	s	2 to 5	Mar. 24	Mar. 28	May 5	June 8	May 30	Sept. 23	Do.
Seibel, No. 1004: Rup. × (Lins. × Vin.)	0	1907	82	s	2 to 5	Mar. 16	Apr. 10	May 28	June 10	June 2	Sept. 22	Oct. 6
Seibel, No. 1070: Rup. × (Lins. × Vin.)	0	1907	86	s	2 to 5	Mar. 22	Apr. 2	May 27	June 1	Sept. 20	Oct. 9
Seibel, No. 1077: Rup. × (Lins. × Vin.)	0	1907	81	s	3 to 5	Mar. 26	Apr. 10	May 29	June 15	June 2	June 21	Sept. 29	Oct. 7
Seibel, No. 2010: Rup. × (Lins. × Vin.)	0	1907	82	s	2 to 5	Mar. 24	Mar. 30	May 24	June 3	May 28	June 8	Sept. 30	Oct. 15
Seibel, No. 2029: Rup. × (Lins. × Vin.)	0	1907	86	s	2 to 5	Apr. 8	May 28	June 18	June 1	June 23	Sept. 24
Seibel, No. 2033: Rup. × (Lins. × Vin.)	0	1907	85	s	2 to 3	Mar. 22	Mar. 24	May 24	June 5	May 28	June 10	Sept. 28	Oct. 10
Seibel, No. 2043: Rup. × (Lins. × Vin. × Rup.)	St	1910	69	s	Mar. 5	Mar. 29	May 17	May 27	May 20	May 31
Seibel, No. 2044: Rup. × (Lins. × Vin. × Rup.)	0	1907	88	s	1 to 5	Mar. 24	Apr. 1	May 22	May 30	May 26	June 6	Sept. 23	Oct. 15
Seibel, No. 2056: Rup. × (Lins. × Vin. × Rup.)	0	1907	89	s	3 to 5	Mar. 26	May 26	May 30	June 3	Sept. 30	Oct. 9
Shala: Lins. × (Rup. × Lab.)	C	1903	90	s	Mar. 31	Apr. 17	May 5	May 20	May 9	May 25
Do.	0	1903	90	s	Mar. 19	Apr. 5	May 19	June 2	May 23	June 7
Winchell: Lab. × (Aest. × Vin.)	0	1905	80	s	Mar. 21	Apr. 2	June 3	June 7	June 5	June 12
Wine King: (Aest. × Lab.) × (Lins. × Rup.)	C	1907	97	s	Mar. 31	Apr. 8	May 4	May 30	May 9	June 4
Do.	0	1903	93	s	2 to 4	Mar. 16	Apr. 3	May 19	May 31	May 25	June 3

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Experiment vineyard.	Year planted.	Growth rating.	How pruned.	Nodes bearing fruit.	Growth-starting date.		Blossoming date.		Fruit-setting date.		Fruit-ripening date.	
						Early season.	Late season.	Early season.	Late season.	Early season.	Late season.	Early season.	Late season.
1		3	4	5	6	7	8	9	10	11	12	13	14
Worden:													
Lab.													
Do.	C	1906	72			Mar. 30	Apr. 20	May 8	May 20	June 5	May 26	Sept. 20	Sept. 30
Do.	Cx	1906	81	s		Mar. 25	Apr. 5	June 1	June 5	June 5	June 4	Aug. 29	Sept. 15
Do.	F	1905	67	s		Mar. 15	do.	May 15	May 30	June 1	June 6	June 6	June 6
Do.	G	1905	56	s		Mar. 5	Apr. 3	May 1	May 23	May 14	May 14	May 13	Sept. 9
Do.	L	1905	72	s		Mar. 11	Apr. 5	May 10	May 23	May 30	June 28	June 5	do.
Do.	M	1905	74	s		Mar. 26	Apr. 6	May 25	June 8	May 30	June 13	June 5	do.
Do.	O	1905	47	s	1 to 2	Mar. 18	Apr. 5	May 17	May 30	June 8	June 12	June 5	do.
Do.	S	1905	80	s		Mar. 11	Apr. 8	May 25	June 9	June 8	June 12	June 5	do.
Do.	F	1905	71	s		Mar. 7	Mar. 25	May 8	May 20	May 12	May 24	Aug. 25	Sept. 5
Do.	G	1905	82	s		Mar. 2	do.	May 24	June 5	May 28	June 10	June 5	do.
Do.	L1	1905	45	s		Mar. 11	Apr. 5	May 25	June 5	June 4	June 5	June 5	do.
Do.	L	1905	66	s		do.	Apr. 1	May 10	May 27	May 14	May 29	June 2	do.
Do.	M	1905	70	s		Mar. 30	Apr. 5	June 5	June 17	June 10	June 22	June 2	do.
Do.	70	1905	85	s	1 to 3	Mar. 18	Mar. 30	May 16	May 25	May 23	June 2	June 2	do.
Do.	S	1905	82	s		Mar. 16	Apr. 5	May 24	June 18	May 27	June 22	June 2	do.
Xhita:													
(Lins. × Rup.) × (Vin. × Lab.)	C	1907	80	s		Mar. 29	Apr. 17	May 2	May 10	May 7	May 20	do.	do.
Do.	F	1903	61	s		Mar. 5	Mar. 25	May 10	May 15	May 15	do.	do.	do.
Do.	O	1903	76	s	2 to 3	Mar. 16	Apr. 1	May 18	June 5	May 24	June 18	Sept. 20	Oct. 1

Wyoming Reed:

Variety and parentage.	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tar- tronic grams per 100 c. c.	Cluster.			Berry.		Use.	
	1909	1910	1911	1912	1913			Size.	Shape.	Com- pact or loose.	Size.	Shape.		Color.
1	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Agawam: Lab. X Vin.....	2	4			1½	<i>Per et.</i> 24.6	0.655	m	cy	c	m, l	r	r	s, st, t
Do.....	3		½			22.5	.6175	m	cy	c	m, l	r	r	s, st, t
Albania: Lins. X (Aest. X Lab. X Bourq.).....	4					24.6	.4425	l	t	m	m	r	w	w, s, st, t
Alicante Ganzin: Vin. X Rup.....	6	3			16	24.2	.8920	l	t	c	m	r	b	w
Alicante X Rupestris Terrace, No. 20: Vin. X Rup.....	50	8				25.2	.6913	s	cy	c	s	r	b	w
Do.....	20	5	15			24.7	.5425	s	cy	c	s	r	b	w
Do.....		1½	8	10		25.6	.86	s	cy	c	s	r	b	w
Amerbonte: Bourq. X (Lins. X Rup.).....								l	t	m	s	r	r	w, s, t
Do.....								l	t	m	s	r	r	w, s, t
Do.....								l	t	m	s	r	r	w, s, t
Do.....								l	t	m	s	r	r	w, s, t
Atoka: (Lins. X Rup.) X (Bourq. X Lab.).....	10	6		2	2	28.8	.8022	l	t	m	s	r	r	w, s, t
Do.....	8					24.9	.9155	l	t	m	s	r	r	w, s, t
Barry: Lab. X Vin.....			1	1		25.5	.6525	m	cy	m	l	r	b	s, st, t
Do.....						25.2	.7875	m	cy	m	l	r	b	s, st, t
Berekmans: Rip. X (Lab. X Bourq.).....			½	½		25.5	.8588	m	cy	c	m	r	b	w, st, t
Do.....			1			23	.825	l	cy	c	m	r	b	w, st, t
Big Extra: Lins. X (Lab. X Vin.).....		½		2		25.8	.8362	l	cy	c	m	r	b	w, st, t
Do.....	8		2	16	1	22.4	.6631	l	cy	c	m	r	b	w, st, t
Do.....			1		2½	22	.9825	l	cy	c	m	r	b	w, st, t
Do.....			1			21	.8025	l	cy	c	m	r	b	w, st, t
Do.....				1		20.6	.9358	l	cy	c	m	r	b	w, st, t
Do.....		2½				23.2	1.2006	l	cy	c	m	r	b	w, st, t
Do.....	10	5				22.9	.8075	l	cy	c	m	r	b	w, st, t
Do.....		4			1	23.8	.8581	l	cy	c	m	r	b	w, st, t
Blondin: Bourq. X Aest.) X (Lins. X Lab.).....	4	3	½			22.6	.6617	l	cy	c	m	r	w	w, t
Do.....	6		5			24.3	.5737	l	cy	c	m	r	w	w, t

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tartaric, grams per 100 C. C.	Cluster.			Berry.			Use.
	1909	1910	1911	1912	1913			Size.	Shape.	Com- pact or loose.	Size.	Shape.	Color.	
I	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Bourisquon X Rupestris, No. 601: Vin. X Rup.			2	7½		Per cent 25.5	1.037	1	t	m	s	r	b	w
Do.	15	25	10	13		23.9	.737	1	t	m	s	r	b	w
Do.	15	25	7	2	16	25.5	.8031	1	t	m	s	r	b	w
Do.			3	3		22.5	1.117	1	t	m	s	r	b	w
Do.			5	5		22	.9825	1	t	m	s	r	b	w
Do.	30	15	10	5		23.2	1.1381	1	t	m	s	r	b	w
Bourisquon X Rupestris, No. 603: Vin. X Rup.			2	4		21.5	.737	1	t	m	s	r	b	w
Do.	15	18	10	4	23	21.3	.8155	1	t	m	s	r	b	w
Do.			4			21	1.2225	1	t	m	s	r	b	w
Do.			6	6		22	.9112	1	t	m	s	r	b	w
Do.		3	6	6		20.1	.94	1	t	m	s	r	b	w
Do.	20		3	6	24	21	.9187	1	t	m	s	r	b	w
Do.		4	3	16	16	23.6	.7631	1	t	m	s	r	b	w
Bourisquon X Rupestris, No. 109-4: Vin. X Rup.			1			27	.93	1	t	m	s	r	b	w
Do.			4	6		20	.8437	1	t	m	s	r	b	w
Do.	15	20	10	13		27.1	.9693	1	t	m	s	r	b	w
Do.		11	5	1		25	1.0858	1	t	m	s	r	b	w
Do.						26	.9525	1	t	m	s	r	b	w
Do.		3				23.2	1.1570	1	t	m	s	r	b	w
Do.	15					23.2	.6975	1	t	m	s	r	b	w
Do.		3	2		8	24.9	.9350	1	t	m	s	r	b	w
Bourisquon X Rupestris, No. 3907: Vin. X Rup.			1½	4½		25	1.012	1	t	m	s	r	b	w
Do.			10	8		24	.8962	1	t	m	s	r	b	w
Do.	20	20	10	8	8	25.2	.8846	1	t	m	s	r	b	w
Do.					9½	32	1.1250	1	t	m	s	r	b	w
Do.								1	t	m	s	r	b	w
Do.		5	2		5	24.4	.7631	1	t	m	s	r	b	w
Bourisquon X Rupestris, No. 4306: Vin. X Rup.			2	2		25	1.02	m	cy	m	s	r	b	w
Do.	6		20	20		23.6	.9295	m	cy	m	s	r	b	w
Do.			1	½		22	1.005	m	cy	m	s	r	b	w
Do.						23.8	.75	m	cy	m	s	r	b	w

Bourisquon X Rupestris, No. 4308: Vin. X Rup.....	2	4½	24.5	1.113	m	m	m	m	m	s	r	r	b	w
Do.....	5	27	.8625	m	m	m	m	m	s	r	r	b	w
Do.....	12	25.6	.8625	m	m	m	m	m	s	r	r	b	w
Do.....	4	21.4	1.4275	m	m	m	m	m	s	r	r	b	w
Do.....	12	25.2	.78	m	m	m	m	m	s	r	r	b	w
Brighton: Lab. X Vin.....	1	7	22.7	.705	m	m	m	m	m	m	r	r	b	s, st, t
Brilliant: Lab. X (Vin. X Bourq.).....	1	7	26.1	.6355	m	m	m	m	m	m	r	r	b	s, st, t
Do.....	2	2½	26.1	.5825	m	m	m	m	m	m	r	r	b	s, st, t
Do.....	2	2	21	.6	m	m	m	m	m	m	r	r	b	s, st, t
Do.....	½	26	.975	m	m	m	m	m	m	r	r	b	s, st, t
Canada: (Rip. X Lab.) X Vin.....	15	15	23.1	.7031	m	m	m	m	m	m	r	r	b	s, st, t
Do.....	1	2	28.4	.7791	m	m	m	m	m	m	r	r	b	s, st, t
Do.....	9	25	.7800	m	m	m	m	m	m	r	r	b	s, st, t
Do.....	6	23.4	.8337	m	m	m	m	m	m	r	r	b	s, st, t
Carignane X Rupestris, No. 404: Vin. X Rup.....	7	12	22.9	.8033	m	m	m	m	m	m	r	r	b	w
Do.....	12	20½	20.8	.7775	m	m	m	m	m	m	r	r	b	w
Do.....	2	2½	24.1	.8572	m	m	m	m	m	m	r	r	b	w
Do.....	6	8	22.2	.887	m	m	m	m	m	m	r	r	b	w
Do.....	5	23.2	.6337	m	m	m	m	m	m	r	r	b	w
Do.....	6	23	.712	m	m	m	m	m	m	r	r	b	w
Carignane X Rupestris, No. 501: Vin. X Rup.....	15	3½	26	.645	m	m	m	m	m	m	r	r	b	w
Do.....	3	22	.9012	m	m	m	m	m	m	r	r	b	w
Do.....	3	24.3	.7462	m	m	m	m	m	m	r	r	b	w
Do.....	7	23	.6	m	m	m	m	m	m	r	r	b	w
Carman: Lins. X (Vin. X Lab. X Bourq.).....	3	1	28.6	.5669	m	m	m	m	m	m	r	r	b	w, t
Do.....	4	4	24.8	.6275	m	m	m	m	m	m	r	r	b	w, t
Castel, No. 1028: Rip. X Vin.....	3	25.2	.6100	m	m	m	m	m	m	r	r	b	w
Castel, No. 10002: (Lab. X Rup.) X Vin.....	5	7	25.2	.6100	m	m	m	m	m	m	r	r	b	w
Catawba: Lab. X Vin.....	1	25.9	.5970	m	m	m	m	m	m	r	r	b	w
Do.....	3	28	.57	m	m	m	m	m	m	r	r	b	w, st, t
Do.....	½	21.7	.397	m	m	m	m	m	m	r	r	b	w, st, t
Do.....	½	50	.6	m	m	m	m	m	m	r	r	b	w, st, t
Do.....	1	4	25.2	.8925	m	m	m	m	m	m	r	r	b	w, st, t
Do.....	½	21	1.005	m	m	m	m	m	m	r	r	b	w, st, t
Do.....	2	2	24.3	.8255	m	m	m	m	m	m	r	r	b	w, st, t
Do.....	5	4	23.5	.697	m	m	m	m	m	m	r	r	b	w, st, t
Centennial: Lab. X Aest.....	2	3	23	.9825	m	m	m	m	m	m	r	r	b	w, t
Do.....	3	5	24.3	.6002	m	m	m	m	m	m	r	r	b	w, t
Do.....	3½	23.4	.7912	m	m	m	m	m	m	r	r	b	w, t
Champnel: Champ. X Lab.....	5	22.8	.78	m	m	m	m	m	m	r	r	b	w, t

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Weight of fruit per vine (pounds).					Sugar, Balling scale.	Acid, as tar- taric, grams per 100 c. c.	Cluster.			Berry.		Use.	
								Shape.	Size.	Com- pact or loose.	Shape.	Color.		
	1909	1910	1911	1912	1913									Size.
I	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Chasselas X Rupestris, No. 901:						<i>Per cl.</i>								
Vin, X Rup.....	5		10			26.5	.6337	m	ey	c	m	I	b	w
Do.....	8					25	.7275	m	ey	c	m	I	b	w
Chairette Doré Ganzin:														
Vin, X Rup.....	4	2		29		20.4	1.1437	1	t	1	m	I	w	w
Do.....				2		25.1	.9825	1	t	1	m	I	w	w
Do.....	60	20	13	8	42	18.3	1.2255	1	t	1	m	I	w	w
Clevener:														
Lab, X (Rip, X Aest).....						25	.7875	s	ey	c	s	I	b	w
Cloeta:														
(Lins, X Rup.) X (Lab, X Vin.).....		4				22.9	.66	1	ey	1	m	I	b	w, t
Columbaud X Rupestris:														
Vin, X Rup.....	10					21.9	.78	m	t	m	m	I	b	w
Lab.....						24	.3975	m	ey	m	m, 1	I	b	w, st, t
Do.....				4		23	.3525	m	ey	m	m, 1	I	b	w, st, t
Do.....						22.4	.57	m	ey	m, 1	m, 1	I	b	w, st, t
Do.....						26	1.132	m	ey	m	m, 1	I	b	w, st, t
Cornucopia:														
Vin, X Rip.....	4					25	.825	1	ey	c	m	I	b	st, t
Do.....						25	.8625	1	ey	c	m	I	b	st, t
Coudere, No. 101:														
Vin, X Rip.....	15	1	3	3	5	28.3	1.1275	s	ey	c	s	I	b	w
Do.....		20	5	5	24	24.4	.8344	s	ey	c	s	I	b	w
Do.....			4	4	24	21	.8675	s	ey	c	s	I	b	w
Do.....			8	8	124	25.1	1.0225	s	ey	c	s	I	b	w
Do.....		2	4	4	4	22.3	1.0916	s	ey	c	s	I	b	w
Do.....	40	8	10	5	44	22	1.0355	s	ey	c	s	I	b	w
Do.....		4	3	3	9	26.1	.9693	s	ey	c	s	I	b	w
Coudere, No. 201:														
Rip, X (Rup, X Vin.).....	15	20	8	15	23	21	.8818	s	ey	c	s	I	b	w
Do.....			1	3	3	19	.7935	s	ey	c	s	I	b	w
Do.....			7	3	3	19	1.1300	s	ey	c	s	I	b	w
Do.....			7	3	3	24	1.1193	s	ey	c	s	I	b	w

Do.....	6	4	4 ¹ / ₂	21.0	1.0716	s	cy	c	r	b	w
Do.....	6	11	5	26	.9570	s	cy	c	r	b	w
Do.....	8	8	21	23	1.0423	s	cy	c	r	b	w
Condere, No. 503: Rup. X Vin.											
Do.....	3	3	2	23.1	.8437	l	t	l	r	b	w
Do.....	25	10	8	21.8	.8190	l	t	l	r	b	w
Do.....	20	2	37	18	.8190	l	t	l	r	b	w
Do.....	31	2	16	19.5	1.042	l	t	l	r	b	w
Do.....	32	10	3	23.3	.795	l	t	l	r	b	w
Do.....	60	16	18	20.3	.8731	l	t	l	r	b	w
Do.....	10	14	13	20.3	.766	l	t	l	r	b	w
Do.....	10	6	28 ¹ / ₂	23.1	.8137	l	t	l	r	b	w
Do.....	8	8	4	21	.8100	s	cy	s	r	b	w
Do.....	8	4	20	22.3	.9383	s	cy	s	r	b	w
Do.....	40	11	3	20	.7065	s	cy	s	r	b	w
Do.....	8	6	3	22.8	.8493	s	cy	s	r	b	w
Condere, No. 3701: Vin. X Rup.											
Do.....	3	2	4	24.6	1.055	m	t	c	r	b	w
Do.....	3	5	3	21	.9000	m	t	c	r	b	w
Do.....	3	12	2	24.4	.9575	m	t	c	r	b	w
Do.....	4	9	9	22.1	1.0162	m	t	c	r	b	w
Do.....	25	6 ¹ / ₂	5	22.1	.8879	m	t	c	r	b	w
Do.....	5	6	8	23.6	.9031	m	t	c	r	b	w
Condere, No. 4401: Rup. X Vin.											
Do.....	5	2	2	26	.7075	m	cy	m	r	b	w
Do.....	3	4	3	23.8	1.1125	m	cy	m	r	b	w
Do.....	10	10	33	24	.9513	m	cy	m	r	b	w
Do.....	2	8	18 ¹ / ₂	21.2	.9712	m	cy	m	r	b	w
Do.....	2	8	1 ¹ / ₂	21.5	1.0856	m	cy	m	r	b	w
Do.....	10	12	5 ¹ / ₂	25.2	.975	m	cy	m	r	b	w
Do.....	10	4	5	22	.9825	m	cy	m	r	b	w
Do.....	3	4	13	24.5	1.0341	m	cy	m	r	b	w
Condere, No. 71-06: Rup. X (Lins. X Vin.)											
Do.....	4	2	3 ¹ / ₂	25	.6543	m	cy	m	r	b	w
Condere, No. 71-20: Rup. X (Lins. X Vin.)											
Do.....	3	4	4	26.6	.8580	m	cy	m	r	b	w
Condere, No. 74-17: Complex Hybrid											
Do.....	3	4	1 ¹ / ₂	24.5	.6575	m	cy	m	r	b	w
Condere, No. 82 X 32: Complex Hybrid											
Do.....	6	8 ¹ / ₂	6	22.1	.65	m	cy	l	0	w	w, t, st
Do.....	2	1 ¹ / ₂	9	22.6	.8819	m	cy	l	0	w	w, t, st
Do.....	3	2	1 ¹ / ₂	19	.9458	m	cy	l	0	w	w, t, st
Do.....	1	2	2	24.5	.9837	m	cy	l	0	w	w, t, st
Do.....	3	6	3 ¹ / ₂	23	.727	m	cy	l	0	w	w, t, st
Do.....	15	3	3	21.1	.7800	m	cy	l	0	w	w, t, st
Do.....	1	4	1 ¹ / ₂	21	.907	m	cy	l	0	w	w, t, st
Condere, No. 84 X 61: (Vin. X Rup.) X Vin.											
Do.....	1	4	1 ¹ / ₂	25.3	.9425	m	t	m	r	b	w
Do.....	20	1	3	24.5	.499	m	t	m	r	b	w
Do.....	20	5	3	23.3	.6032	m	t	m	r	b	w

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Weight of fruit per vine (pounds).						Sugar, Balling scale.	Acid, as tar- taric, grams per 100 C. C.	Cluster.		Berry.			Use.
	1909								Shape.	Com- pact or loose.	Size.	Shape.	Color.	
	1900	1910	1911	1912	1913	1914								
I	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Couderc, No. 84 × 61—Continued. (Vin. × Rup.) × Vin.	2 3	1 10	3	<i>P. et al.</i> 20.6	.9275	m	t	m	m	r	b	w
Do.	3	2	24	.7969	m	t	m	m	r	b	w
Do.	4	16	23.8	.615	m	t	m	m	r	b	w
Do.	6	21.5	1.077	m	t	m	m	r	b	w
Do.	4	22.5	.715	m	t	m	m	r	b	w
Do.	4	6	26.5	.8468	m	t	m	m	r	b	w
Couderc, No. 85 × 113; Vin. × (Rup.) × Vin.	6	4	2½	2½	23.7	.8156	m	cy	m	m	r	b	w
Couderc, No. 87 × 115; Vin. × (Rup. × Vin.)	1	3	24	.57	l.	cy t	l	s	r	b	w
Do.	2	3	26	.6562	l.	cy t	l	s	r	b	w
Do.	20	4	34	21	23.9	.372	l.	cy t	l	s	r	b	w
Do.	2	2	9	20	.7875	l.	cy t	l	s	r	b	w
Do.	6	10½	23.8	.7875	l.	cy t	l	s	r	b	w
Do.	6	24.7	.8095	l.	cy t	l	s	r	b	w
Do.	35	10	13	6	21.8	.6224	l.	cy t	l	s	r	b	w
Do.	6	22	21.7	.7612	l.	cy t	l	s	r	b	w
Couderc, No. 124 × 30; Vin. × (Rup. × Vin.)	25	15	8	7	12	21.7	.877	m	cy	m	l	r	b	w
Do.	8	8	22.3	.9105	m	cy	m	l	r	b	w
Do.	7	4	11½	19.5	.8927	m	cy	m	l	r	b	w
Do.	4	23.5	1.0744	m	cy	m	l	r	b	w
Do.	6	6	23.3	1.995	m	cy	m	l	r	b	w
Do.	14	17	21.8	1.1555	m	cy	m	l	r	b	w
Do.	40	5	4	28	21.8	.8868	m	cy	m	l	r	b	w
Do.	4	12	22.3	.9702	m	cy	m	l	r	b	w
Couderc, No. 132-11; Complex Hybrid	6	2	2	2	18	22.7	.5106	l	t	m	m	r	b	w
Couderc, No. 199-88; Vin. × (Rup. × Vin.)	3	26	.6632	m	t	m	m	r	w	w, t
Couderc, No. 241-125; Vin. × (Rup. × Vin.)	1	3	26	.7162	m	cy	m	s	r	b	w
Couderc, No. 267-27; Complex Hybrid	3	2	10	29	.6262	m	cy	m	m	r	b	w

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Weight of fruit per vine (pounds).						Sugar, Balling scale.	Acid, as tar- taric, grams per 100 C. C.	Cluster.			Berry.			Use.
	1900								Shape.	Size.	Com- pact or loose.	Shape.	Size.	Color.	
	1909	1910	1911	1912	1913	1914									
1	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
Isabella: Lab. × Vin.						<i>Per ct.</i> 26	1.658	m	cy	c	m	f	b	st, t	
Do.	2	2	1					m	cy	c	m	f	b	st, t	
Do.						18	.587	m	cy	c	m	f	b	st, t	
Do.								m	cy	c	m	f	b	st, t	
Do.				1½		28	.72	m	cy	c	m	f	b	st, t	
Do.			1½			23	.3037	m	cy	c	m	f	b	st, t	
Do.	3	6	6		22	.7837		m	cy	c	m	f	b	st, t	
Jaeger: Lins. × Bourq.	6		½		1½	25	.6921	m	cy	c	s	f	b	w	
Do.					3	23	1.027	m	cy	c	s	f	b	w	
Do.								m	cy	c	s	f	b	w	
Do.	3					24.9	.81	m	cy	c	s	f	b	w	
Jefferson: Lab. × Vin.								m	cy	c	l	f	f	st, t	
Do.	3	1			1	29	.3777	m	cy	m	l	f	f	w, t	
Kiowa: Lins. × Bourq.	3							m	cy	m	l	f	f	w, t	
Do.								s	cy	m	l	f	f	w, t	
Lamparas: Lab. × Aest.						20.8	.66	m	cy	m	m	f	f	t	
Laussel: Lins. × (Lab. × Aest.)								m	cy	c	m	f	f	t	
Lenoir: Bourq.						23	.5625	l	t	l	s	f	b	w	
Do.	5	2	2			29.2	.977	l	t	l	s	f	b	w	
Do.	15	5	5	30	10	23.6	1.3234	l	t	l	s	f	b	w	
Do.		6	1			29.2	.7375	l	t	l	s	f	b	w	
Do.			2		9	27	.995	l	t	l	s	f	b	w	
Do.			1½			24	1.1212	l	t	l	s	f	b	w	
Do.						24.9	1.1141	l	t	l	s	f	b	w	
Do.		1½				22	1.295	l	t	l	s	f	b	w	
Do.		5		5½		24	1.0575	l	t	l	s	f	b	w	
Do.	15	2	2		4	23.8	.8493	l	t	l	s	f	b	w	

Lingley: Lab. X Vin.....	8	1	2	1½	1	26.4	.5000	m	cy	l	m	r	r	w, t
Do.....	4	5	8	22.5	.5512	m	cy	l	m	r	r	w, t
Louisiana: Do.....	12	5	4	23.1	.4500	m	cy	l	m	r	r	w, t
Bourq.....	23.3	.5300	m	cy	l	m	r	r	w, t
Lakfala: Do.....	25.5	.7350	s	cy	c	s	r	b	w, t
Champ. X Lab.....	30	.6325	s	cy	c	s	r	b	w, t
Do.....	20	.7275	m	cy	c	l	r	b	t
Do.....	m	cy	c	l	r	b	t
Manito: (Lab. X Vin. X Bourq.) X (Lins. X Rup.).....	3	23.6	.7012	m	cy	c	m	r	r	t
Marguerite: Lins. X Bourq.....	22	.6075	m	cy	c	m	r	b	w, t
Do.....	26.5	.90	m	cy	c	m	r	b	w, t
Do.....9	m	cy	c	m	r	b	w, t
Do.....	26	m	cy	c	m	r	b	w, t
Martha: Lab. X Vin.....	23	.51	m	t	l	m	r	w	t
Missouri Riesling: Rip. X Lab.....	2	4	1	4	6	23.9	.5791	s	cy	c	s	r	r	w
Do.....	1½	25.6	.6300	s	cy	c	s	r	r	w
Do.....	3	1	1½	25.5	.4900	s	cy	c	s	r	r	w
Mrs. Munson: Lins. X Bourq.....	4	½	1	23.6	.6775	m	cy	l	m	r	b	t
Muench: Lins. X Bourq.....	4	6	2	8	4	25.7	.6700	m	cy	c	s	r	b	w
Do.....	2	4	4	23.9	.8662	m	cy	c	s	r	b	w
Niagara: Lab. X Vin.....	26	.51	m	cy	m	l	r	w	w, st, t
Do.....	23.8	.485	m	cy	m	l	r	w	w, st, t
Do.....	25.6	.7275	m	cy	m	l	r	w	w, st, t
Oliatao: (Vin. X Lab.) X (Lins. X (Vin. X Lab.)).....	25.6	.8625	m	t	l	m	r	w	t
Pardes: Vin. X Rup.....	23.5	.8925	m	cy	m	s	r	b	w
Do.....	20	5	3	3½	22.9	.825	m	cy	m	s	r	b	w
Do.....	15	15	5	15	22.4	.65	m	cy	m	s	r	b	w
Do.....	25.3	.9262	m	cy	m	s	r	b	w
Do.....	22.5	.8612	m	cy	m	s	r	b	w
Do.....	23.8	.9725	m	cy	m	s	r	b	w
Do.....	25	3	11	4	10	23.8	.9725	m	cy	m	s	r	b	w
Do.....	25.1	.8775	m	cy	m	s	r	b	w
Pierce: Lab. X Vin.....	24.1	.6208	m	cy	c	l	o	b	st, t
Do.....	21	.9825	m	cy	c	l	o	b	st, t
Do.....	19.7	.9187	m	cy	c	l	o	b	st, t
Rebecca: Lab. X Vin.....	1½	24.6	.5137	m	cy	m	s	o	w	t
Do.....	26	.75	m	cy	m	s	o	w	t
Do.....	6	22.2	.4012	m	cy	m	s	o	w	t
Do.....	23	.6244	m	cy	m	s	o	w	t

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Weight of fruit per vine (pounds).					Sugar, Baling scale.	Acid, as tartaric, grams per 100 c. c.	Cluster.			Berry.			Use.		
	1							20	21	22	23	24	25		26	27
	1909	1910	1911	1912	1913											
Rommel: Lab. X (Rip. X Vin.)	6	3	5	6	6	Par ct. 95	.4095	m	cy	1	1	F	w	w, t		
Do.	21	21	21	2	8	23.1	.5062	m	cy	1	1	F	w	w, t		
Do.						23	.3975	m	cy	1	1	F	w	w, t		
R. W. Mission: Lins. X (Lab. X Vin.)								m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Do.	3	4	4			26.2	.5075	m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Do.	2					26.6	.7513	m-l	cy	1	1	F	b	w, t		
Do.								m-l	cy	1	1	F	b	w, t		
Seibel, No. 1: Rup. X (Lins. X Vin.)								m	cy	1	1	F	b	w		
Do.		2	1	6½		37	.7018	m	cy	1	1	F	b	w		
Do.	18	15	15			22.8	.9081	m	cy	1	1	F	b	w		
Do.	30	7	2	5	1	26	.5580	m	cy	1	1	F	b	w		
Do.			4		14	22	.9037	m	cy	1	1	F	b	w		
Do.						24.7	.705	m	cy	1	1	F	b	w		
Do.				4		28.1	.765	m	cy	1	1	F	b	w		
Do.		3	3	7		22.2	.9886	m	cy	1	1	F	b	w		
Do.		2	3	7		25.1	.7437	m	cy	1	1	F	b	w		
Do.	50	5	9	9	6	25.4	.8418	m	cy	1	1	F	b	w		
Do.		10	3			25.4	.8418	m	cy	1	1	F	b	w		
Do.								m	cy	1	1	F	b	w		
Seibel, No. 2: Rup. X (Lins. X Vin.)								m	cy	1	1	F	b	w		
Do.		2	3	5½		26	1.1766	m	cy	1	1	F	b	w		
Do.	25	18	18			24	1.2825	m	cy	1	1	F	b	w		
Do.	30	6	2	18	3	26	.7290	m	cy	1	1	F	b	w		
Do.			2	4½	23½	20	.7925	m	cy	1	1	F	b	w		
Do.						25	.9487	m	cy	1	1	F	b	w		

Do.....	3	3	7	25.5	1.1737	m	cy	l	l	r	b	w
Do.....	3	4	6	23.1	1.0991	m	cy	l	l	r	b	w
Do.....	2	6	3	23.5	1.115	m	cy	l	l	r	b	w
Do.....	50	1	3	24	1.0177	m	cy	l	l	r	b	w
Do.....						m	cy	l	l	r	b	w
Seibel, No. 14: Rup. X (Lins. X Vin.)						m	cy	l	l	r	b	w
Do.....	1	1	8	29	.960	m	cy	l	l	r	b	w
Do.....	6	10	10	23	.7750	m	cy	l	l	r	b	w
Do.....	15			24	1.3325	m	cy	l	l	r	b	w
Seibel, No. 20: Rup. X (Lins. X Vin.)	2	3	3	24	.8895	s	cy	c	s	r	b	w
Seibel, No. 38: Rup. X (Lins. X Vin.)						s	cy	c	s	r	b	w
Seibel, No. 60: Rup. X (Lins. X Vin. X Rup.)						s	cy	c	s	r	b	w
Seibel, No. 70: Rup. X (Lins. X Vin. X Rup.)						s	cy	c	s	r	b	w
Seibel, No. 78: Rup. X (Lins. X Vin.)	5	3	4	13	.9144	s	cy	c	s	r	b	w
Seibel, No. 80: Rup. X (Lins. X Vin.)	4	4		6	.6212	m	cy	c	m	r	b	w
Seibel, No. 128: Rup. X (Lins. X Vin.)	4	4		18	.6725	m	cy	c	m	r	b	w
Seibel, No. 136: Rup. X (Lins. X Vin.)	1½	2		8	.9335	m	t	m	m	r	b	w
Seibel, No. 209: Rup. X (Lins. X Vin.)	2	1	1½	31	.9518	m	t	m	m	r	b	w
Seibel, No. 215: Rup. X (Lins. X Vin.)		1		9	.7912	m	t	m	m	r	b	w
Do.....	6	10		22.3	.8375	m	cy	m	m	r	b	w
Do.....	½	4	25	24	.8250	m	cy	m	m	r	b	w
Seibel, No. 334: Rup. X (Lins. X Vin. X Rup.)	2	1		6	1.150	l	cy	c	m	r	b	w
Seibel, No. 1004: Rup. X (Lins. X Vin.)	4	4	3	11	.9555	l	cy	m	m	r	b	w
Seibel, No. 1070: Rup. X (Lins. X Vin.)	5	5	3	9	.9593	l	cy	m	m	o	b	w
Seibel, No. 1077: Rup. X (Lins. X Vin.)	1			6	.7275	m	cy	c	m	r	b	w
Seibel, No. 2010: Rup. X (Lins. X Vin.)		½	1	2	.8487	s	cy	c	m	r	b	w
Seibel, No. 2029: Rup. X (Lins. X Vin.)		1		3	1.155	s	cy	c	m	r	b	w
Seibel, No. 2033: Rup. X (Lins. X Vin.)	3	2	3	12	.8812	l	cy	c	m	r	b	w
Seibel, No. 2043: Rup. X (Lins. X Vin. X Rup.)						s	cy	c	m	r	b	w
Seibel, No. 2044: Rup. X (Lins. X Vin. X Rup.)	3	1½	2	5	1.1535	s	cy	c	m	r	b	w
Seibel, No. 2056: Rup. X (Lins. X Vin. X Rup.)		1		6	.9750	s	cy	c	m	r	b	w

TABLE X.—Relative behavior and value for different purposes of improved native American and Franco-American varieties of grapes growing on their own roots in eleven experiment vineyards in California—Continued.

Variety and parentage.	Weight of fruit per vino (pounds).					Sugar, Balling scale.	Acid, as tartaric, grams per 100 c. c.	Cluster.			Berry.			Use.
	1909							Size.	Shape.	Com- pact or loose.	Size.	Shape.	Color.	
	1909	1910	1911	1912	1913									
1	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Shaha: Lins. × (Rup. × Lab.).....						<i>Per ct.</i>		1	cy	c	1	r	b	w, t
Do.....								1	cy	m	1	r	b	w, t
Winchell: Lab. × (Aest. × Vin.).....								1	t	1	s	r	w	w, t
Wine King: (Aest. × Lab.) × (Lins. × Rup.).....								1	cy	c	m	r	b	w
Do.....						22.8	0.9000	1	cy	c	m	r	b	w
Worden: Lab.....								m	cy	1	m	r	b	t
Do.....						27	.4850	m	cy	1	m	r	b	t
Do.....	3					23.3	.5325	m	cy	1	m	r	b	t
Do.....								m	cy	1	m	r	b	t
Do.....								m	cy	1	m	r	b	t
Do.....								m	cy	1	m	r	b	t
Do.....						20.9	.5775	m	cy	1	m	r	b	t
Do.....								m	cy	1	m	r	b	t
Do.....								m	cy	1	m	r	r	t
Wyoming Red: Lab.....	1½	1	1			24.9	.5300	m	cy	1	m	r	r	t
Do.....								m	cy	1	m	r	r	t
Do.....								m	cy	1	m	r	r	t
Do.....								m	cy	1	m	r	r	t
Do.....								m	cy	1	m	r	r	t
Do.....						24.8	.6150	m	cy	1	m	r	r	t
Do.....								m	cy	1	m	r	r	t
Do.....								m	cy	1	m	r	r	t
Xlnbr: (Lins. × Rup.) × (Vin. × Lab.).....								1	cy	m	m	r	b	w
Do.....								1	cy	m	m	r	b	w
Do.....						23	.9375	1	cy	m	m	r	b	w

CONCLUSIONS AND SUGGESTIONS.

Varying soil, climatic, and other conditions complicate the successful establishment of vineyards on resistant stocks.

The adaptability of varieties to soil, climatic, and other conditions can be closely forecasted, but congeniality has to be determined by actual test.

The best results are obtained where the scion and stock are congenial and both are suited to all the conditions of the environment.

When both scion and stock varieties are suited to the conditions but do not thrive, congeniality is probably lacking.

The ideal vine is one having a most resistant root which is congenial to a top that produces the best fruit abundantly.

Different species used as stocks with the same variety grafted on them (Table VII) may increase or diminish its vigor and productivity; increase or diminish the quality, size, and appearance of the fruit; cause it to ripen earlier or later; and bring about results varying from perfect success to almost complete failure.

Extensive saccharine and acid determinations made (Table VII) of varieties grafted on resistant stocks and contrasted with the congeniality and growth ratings made of the same vines the same season show a close correspondence between these important chemical constituents of the fruit and the congeniality of graft and stock. Similar growth ratings of a variety grafted on various stocks are found to be accompanied by fairly definite percentages of sugar and acid. The congeniality of the variety to the stock materially affects the resistant qualities of the stock.

The quantity and quality of the fruit are usually in opposition on the soils and vines producing most abundantly; the fruit is usually not of as much value per given unit as it is on vines that are relatively less productive.

Most vine varieties making perfect growth on resistant stocks are found to yield heavier crops than the same variety when grown on its own roots.

The best results can be obtained only when the varieties are placed under soil, climatic, and other conditions to which they are adapted and by using the methods of pruning, training, and culture best suited to each one.

A number of the new grape introductions of the Department of Agriculture are proving to be superior to the varieties that are now commercially grown for certain purposes.

The relative rooting qualities of resistant varieties are an important consideration in the cost of establishing resistant vineyards.

Some stocks are suited for bench grafting, while others are especially valuable for vineyard grafting.

Cuttings of many hybrids root easily, although the cuttings from one of the parents may be hard to root.

Where conditions are not suited to a given species, they are often well adapted to hybrids of that species with some other species.

As cuttings of *Monticola*, *Berlandieri*, *Aestivalis*, *Linsecomii*, *Bicolor*, and *Candicans* (Pls. VI, VII, and VIII) are hard to root, they should be rooted in the nursery and grafted there, or planted in the vineyard and grafted afterwards.

Riparia cuttings root easily and are excellent stocks well suited for vineyard and bench grafting, but they are adapted to but few California soils. Soils in which *Riparia* varieties thrive usually produce large crops of only fair quality.

Rupestris cuttings root and graft easily and are best adapted to bench grafting. When so used the dormant eyes should be cut out of the stock. Many varieties are not congenial to *Rupestris*, and their fruit is usually somewhat later in ripening than when grown on some other stock.

In most instances *Riparia*, *Berlandieri*, *Champini*, and *Aestivalis* stocks (Pl. VI, VII, and VIII) are congenial to *Vinifera* varieties. Their fruitfulness is increased and the time of ripening hastened in comparison with the same varieties grown on other stocks.

Some of the hybrid resistant-stock varieties are making enviable records as stocks under California conditions.

Where all the qualities desired can not be found in a hybrid, a complex hybrid—that is, a hybrid of hybrids—may yield the desired results.

A grower of *Vinifera* grapes should decide before locating his vineyard what varieties he desires to grow, and then choose soil and other conditions suited to such varieties. He should know whether stocks are to be established in the vineyard and grafted afterwards or whether the plantings are to be of bench or nursery grafts. He should then select the resistant varieties best suited to the purpose and conditions and which at the same time are congenial to the varieties he intends to grow. He should familiarize himself with all the operations necessary in establishing a resistant vineyard.

The amount of money practically thrown away in the reestablishment of *Vinifera* vineyards in this country since the first appearance of phylloxera in them can not be even approximately estimated.

The direct causes of this waste of money have been due to lack of information and the fact that there was no source from which data could be obtained. This has resulted in the taking of chances by the growers in planting nonresistants, or in using the wrong resistants, or in using resistants which were not congenial to the varieties they were growing. Other causes for this waste have been the purchase of bench grafts on resistant stocks not true to

name and the lack of proper care and management of resistant vineyards, such as allowing roots to grow from Vinifera tops grown on resistant stocks. These mistakes have delayed the general use of resistant. There should be no further delay of this kind. The Department of Agriculture is now prepared and will be glad to give information of value along all these lines.



